

**ACETAMINOPHEN DEXTROMETHORPHAN HYDROBROMIDE PHENYLEPHRINE
HYDROCHLORIDE- acetaminophen dextromethorphan hydrobromide phenylephrine
hydrochloride capsule, liquid filled
Granules India Limited**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

**Acetaminophen 325 mg
Dextromethorphan Hydrobromide 10 mg
Phenylephrine Hydrochloride 5 mg**

Active ingredient

(in each Softgel)

Acetaminophen 325 mg

Dextromethorphan hydrobromide 10 mg

Phenylephrine hydrochloride 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves these symptoms due to cold and flu:
sneezing
itching of the nose, throat or watery eyes due to hay fever
cough
nasal congestion
sinus congestion and pressure
sore throat
head ache
minor aches and pains
- helps clear nasal passages and shrinks swollen membranes
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 10 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin or severe allergic reactions. Symptoms may

include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you ever had an allergic reaction to this product or any of its ingredients.
- in children 12 years of age.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough with excess phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dosage

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or head ache that lasts for 1 week, these could be the signs of

serious condition

- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding

ask a health professional before use

Keep out of the reach of children

In case of overdose, get medical help or contact Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than the recommended dose

adults & children under 12 years and over

- take 2 softgels with water every 4 hours.
- do not exceed 10 softgels in 24 hours or as directed by a doctor

children under 12 years

- do not use

Other information

- store in a cool and dry place.
- protect from sunlight.
- **Parents: Learn about teen medicine abuse, WWW.StopMedicineAbuse.org**

Inactive ingredients

FD&C Blue 1, FD&C Red 40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sodium hydroxide, sorbitol sorbitan, titanium dioxide


Questions or comments?

call **1-877-770-3183** Mon-Fri 9:00 AM to 4:30 PM EST

Daytime Softgels

80mm

105mm



nm Note: Ensure that the unvarnished area is not less than the minimum area specified o

JOB INFORMATION	HISTORY	COLORS
ITEM#: DFT_BUSTER_80mmX105mm_SevereCoughSoreThroatDay_R1 BLISTER TEMPLATE	Version# 1 Printed	Black
CAD#:	Rev-1: Rev-6:	
	Rev-2: Rev-7:	
	Rev-3: Rev-8:	
	Rev-4: Rev-9:	
	Rev-5: Rev-10:	Q.C. _____

GRANULES ARTWORK APPROVAL (THIS PROOF MUST BE SIGNED AND RETURNED)	
Name (print)	Signature

ATTENTION: PLEASE PROOFREAD CAREFULLY.
 ed by a third party for Granules and is intended to represent the image prepared for you according to the instructions provided. Neither supplier nor G
 either in format or content. This proof does not accurately represent colors that will print in the finished product. Any colors referenced on this proof
 ed product will be printed to match the referenced Pantone® color or any other approved match standard established or provided by the supplier or
 By signing this proof you are agreeing to these terms and authorizing its release into production under these conditions.

ACETAMINOPHEN DEXTROMETHORPHAN HYDROBROMIDE

PHENYLEPHRINE HYDROCHLORIDE

acetaminophen dextromethorphan hydrobromide phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62207-912
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POVIDONE (UNII: FZ989GH94E)	
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	G02
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62207-912-69	12 in 1 BLISTER PACK; Type 0: Not a Combination Product	07/07/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/07/2017	

