

**MUCINEX FAST-MAX CONGESTION AND HEADACHE- acetaminophen,  
guaifenesin, and phenylephrine hydrochloride solution  
RB Health (US) LLC**

-----  
**Mucinex® Fast-Max®  
Congestion & Headache**

**Drug Facts**

<b>Active ingredients (in each 20 mL)</b>	<b>Purposes</b>
<b>Acetaminophen 650 mg</b>	<b>Pain reliever/fever reducer</b>
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses**

- temporarily relieves these common cold and flu symptoms:
  - nasal congestion
  - headache
  - minor aches and pains
  - sinus congestion and pressure
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### **When using this product do not use more than directed**

#### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours

- children under 12 years of age: do not use

### **Other information**

- each 20 mL contains: **sodium 12 mg**
- store between 20-25°C (68-77°F)
- do not refrigerate

### **Inactive ingredients**

anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate <sup>1</sup>, xanthan gum

---

<sup>1</sup> may contain this ingredient

### **Questions?**

**1-866-MUCINEX (1-866-682-4639)**

You may also report side effects to this phone number.

Dist. by: Reckitt Benckiser  
Parsippany, NJ 07054-0224

### **PRINCIPAL DISPLAY PANEL - 180 mL Bottle Label**

#### **MAXIMUM STRENGTH\***

NDC 63824-536-66

060616 3036245

**Mucinex®**  
**FAST-MAX®**

#### **CONGESTION & HEADACHE**

**Acetaminophen**– Pain Reliever/Fever Reducer  
Guaifenesin – Expectorant  
Phenylephrine HCl – Nasal Decongestant

- ✓ **Relieves Headache & Fever**
- ✓ **Relieves Nasal & Chest Congestion**
- ✓ **Thins & Loosens Mucus**

**6 FL OZ (180mL)**

**FOR AGES 12+**

MAXIMUM STRENGTH\*

NDC 63824-536-66

060616 3036245



## CONGESTION & HEADACHE

**Acetaminophen** – Pain Reliever/Fever Reducer  
**Guaifenesin** – Expectorant  
**Phenylephrine HCl** – Nasal Decongestant

- ✓ Relieves Headache & Fever
- ✓ Relieves Nasal & Chest Congestion
- ✓ Thins & Loosens Mucus

6 FL OZ (180mL)

FOR AGES 12+

### Drug Facts

#### Active ingredients (in each 20 mL) Purposes

Acetaminophen 650 mg.....	Pain reliever/fever reducer
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg....	Nasal decongestant

#### Uses

- temporarily relieves these common cold and flu symptoms:
  - nasal congestion
  - headache
  - minor aches and pains
  - sinus congestion and pressure
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

### Drug Facts (continued)

- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

PEEL HERE

### Drug Facts (continued)

- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not use more than directed**

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not take more than directed (see Overdose warning)

**Drug Facts** (continued)

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

**Other information**

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F)
- do not refrigerate

**Inactive ingredients** anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate†, xanthan gum †may contain this ingredient

**Questions?**

**1-866-MUCINEX (1-866-682-4639)**  
You may also report side effects to this phone number.



HEALTH • HYGIENE • HOME

Please visit our website  
[www.mucinex.com](http://www.mucinex.com)

**PEEL CORNER TO READ COMPLETE  
DRUG FACTS AND INFORMATION**

\*Per 4-hour dose

**Tamper evident: Do not use if neckband  
on bottle cap is broken or missing.**



3 63824 01665 8

Dist. by: Reckitt Benckiser  
Parsippany, NJ 07054-0224  
©2018 RB

011018

LOT:  
EXP:  
MADE IN:  
3055005

**MUCINEX FAST-MAX CONGESTION AND HEADACHE**

acetaminophen, guaifenesin, and phenylephrine hydrochloride solution

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63824-536
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

**Product Characteristics**

<b>Color</b>	blue	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63824-536-66	180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)	10/01/2016	04/01/2025

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/01/2016	04/01/2025

**Labeler** - RB Health (US) LLC (081049410)