

**CORICIDIN HBP NIGHTTIME MULTI-SYMPTOM COLD-
acetaminophen, dextromethorphan hydrobromide, doxylamine
succinate liquid
Bayer HealthCare LLC.**

Coricidin HBP Nighttime Multi-Symptom Cold Liquid

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

Purpose

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- headache
- sore throat
- runny nose
- minor aches and pains
- cough due to minor throat and bronchial irritation
- sneezing
- fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 120 mL (4 doses) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin or severe allergic reactions. Symptoms may

include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- glaucoma
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- redness or swelling is present

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help

or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over: take 30 mL in the dosing cup provided every 6 hours while symptoms last
- do not exceed 120 mL (4 doses) in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- **each 15 mL contains:** sodium 18 mg
- store at 20°-25°C (68°-77°F)

Inactive ingredients

alcohol 10%, anhydrous citric acid, FD&C blue no.1, FD&C red no.40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments

Questions or comments? 1-800-317-2165 (Mon – Fri 9AM – 5PM EST)

Drug Facts (continued)

Ask a doctor before use if you have

- liver disease
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- glaucoma
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- redness or swelling is present

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- adults and children 12 years and over: take 30 mL in the dosing cup provided every 6 hours while symptoms last
- do not exceed 120 mL (4 doses) in 24 hours, unless directed by a doctor
- children under 12: ask a doctor

Other information

- each 15 mL contains: sodium 18 mg
- store at 20°-25°C (68°-77°F)

Inactive ingredients alcohol 10%, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)

Coricidin[®]

HBP DECONGESTANT-FREE COLD SYMPTOM RELIEF for people with **HIGH BLOOD PRESSURE**

NIGHTTIME[®]
MULTI-SYMPTOM COLD

RELIEVES:

- Sore Throat
- Cough
- Sneezing
- Runny Nose
- Body Aches, Pains, Headache and Fever

Alcohol 10%

Cherry FLAVORED

12 FL OZ (355 mL)

Dextromethorphan HBr - Cough suppressant, Doxylamine succinate - Antihistamine, Acetaminophen - Pain reliever/fever reducer

See New Warnings Information

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

PARENTS: Learn about teen medicine abuse www.StopMedicineAbuse.org

DECONGESTANT-FREE COLD SYMPTOM RELIEF for people with **HIGH BLOOD PRESSURE**

Drug Facts

Active ingredients (in each 15 mL)

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

Uses

- temporarily relieves common cold/flu symptoms:
- headache
- sore throat
- runny nose
- minor aches and pains
- cough due to minor throat and bronchial irritation
- sneezing
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 120 mL (4 doses) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin redness
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

CORICIDIN HBP NIGHTTIME MULTI-SYMPTOM COLD

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11523-4358
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg in 15 mL	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg in 15 mL	

Inactive Ingredients		
Ingredient Name		Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523-4358-1	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/20/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/20/2015	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 11/2023

Bayer HealthCare LLC.