ALKA-SELTZER PLUS MAXIMUM STRENGTH NIGHT COLD AND FLUacetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled Bayer HealthCare LLC.

Alka-Seltzer Plus® Maximum Strength Night Cold & Flu Liquid Gels

Drug Facts

Active ingredients (in each capsule)

Acetaminophen 325 mg

Dextromethorphan hydrobromide 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine hydrochloride 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

Uses

- \cdot temporarily relieves these symptoms due to a cold or flu:
- \cdot minor aches and pains
- headache
- \cdot nasal and sinus congestion
- · cough
- sore throat
- · runny nose
- sneezing
- \cdot temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- \cdot more than 4000 mg of acetaminophen in 24 hours
- \cdot with other drugs containing acetaminophen

 \cdot 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe

allergic reactions. Symptoms may include:

 \cdot skin reddening \cdot blisters \cdot rash \cdot hives

 \cdot facial swelling \cdot asthma (wheezing) \cdot shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than

2 days, is accompanied or followed by fever, headache, rash, nausea,

or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use

 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

• if you have ever had an allergic reaction to this product or any of its ingredients

• in children under 12 years of age

Ask a doctor before use if you have

- liver disease heart disease high blood pressure
- thyroid disease diabetes glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

 persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

 \cdot pain, cough, or nasal congestion gets worse or lasts more than

7 days

- · fever gets worse or lasts more than 3 days
- \cdot redness or swelling is present
- new symptoms occur
- \cdot cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

· nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help

or contact a Poison Control Center right away. Quick medical attention

is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

 \cdot do not take more than the recommended dose

 \cdot adults and children 12 years and over: take 2 capsules with water

every 4 hours. Do not exceed 10 capsules in 24 hours or as

directed by a doctor.

 \cdot children under 12 years: do not use

Other information

Other information

 \cdot store at room temperature. Avoid excessive heat above 40°C

(104ºF).

Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol

sorbitan solution, titanium dioxide

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)

Alka-Selter PLUS[®]

MAXIMUM STRENGTH Cold & Flu

NIGHT

ACETAMINOPHEN /Pain Reliever-Fever Reducer

Dextromethorphan Hydrobromide / Cough Suppresant

Doxylamine Succinate / Antihistamine

Phenylephrine HCl / Nasal Decongestant

- Cough
- Nasal Congestion
- Runny Nose
- Headache & Body Ache
- Sore Throat

20 LIQUID GELS (Liquid Filled Capsules)



ALKA-SELTZER PLUS MAXIMUM STRENGTH NIGHT COLD AND FLU acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0280-1600	
Route of Administration	ORAL			

Active Ingredient/	Active Moiety					
	Ingredient Name		Basis of S	trength	Strength	
DEXTROMETHORPHAN H (DEXTROMETHORPHAN - U		9D2RTI9KYH)		DEXTROMETHORPHAN HYDROBROMIDE		
ACETAMINOPHEN (UNII:	36209ITL9D) (ACETAMIN	IOPHEN - UNII:36209ITL	D) ACETAMINOPHE	N	325 mg	
DOXYLAMINE SUCCINAT UNII:95QB77JKPL)	E (UNII: V9BI9B5YI2) (D	OXYLAMINE -	DOXYLAMINE SU	JCCINATE	6.25 mg	
PHENYLEPHRINE HYDRO UNII:1WS297W6MV)	OCHLORIDE (UNII: 04JA5	59TNSJ) (PHENYLEPHRINE	- PHENYLEPHRINE HYDROCHLORID		5 mg	
Inactive Ingredien	ts					
	Ingredie	nt Name		St	trength	
TITANIUM DIOXIDE (UNII	: 15FIX9V2JP)					
FD&C BLUE NO. 1 (UNII:	H3R47K3TBD)					
GLYCERIN (UNII: PDC6A30	C0OX)					
WATER (UNII: 059QF0KO0	IR)					
POLYETHYLENE GLYCO	L, UNSPECIFIED (UNII:	3WJQ0SDW1A)				
POVIDONE (UNII: FZ989G	iH94E)					
D&C YELLOW NO. 10 (U	NII: 35SW5USQ3G)					
GELATIN (UNII: 2G86QN32	27L)					
PROPYLENE GLYCOL (UN	NII: 6DC9Q167V3)					
SHELLAC (UNII: 46N107B	710)					
SODIUM HYDROXIDE (UN	NII: 55X04QC32I)					
SORBITAN (UNII: 6092IC)	/9RU)					
SORBITOL (UNII: 506T60A	425R)					
Product Character	ristics					
Color	green	Score no s		no score	score	
Shape	OVAL	Size		20mm		
Flavor				AS;NITE		
Contains						
Packaging						
# Item Code	Package Desc	ription	Marketing Start Date		ting End ate	

l	π	item coue	i dekage beschption	Date	Date
		NDC:0280- 1600-20	2 in 1 CARTON	06/19/2017	
	1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
I					

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
Category	Citation	Date	Date
OTC Monograph Drug	M012	06/19/2017	

Revised: 12/2023

Bayer HealthCare LLC.