

**ALKA-SELTZER PLUS SEVERE COUGH MUCUS AND CONGESTION DAY AND NIGHT- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride, guaifenesin  
Bayer HealthCare LLC.**

-----

**Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Day & Night Liquid Gels**

**Do not take these products at the same time.**

**Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Day Liquid Gels**

***Drug Facts***

**Active ingredients (in each capsule)**

Acetaminophen 250 mg

Dextromethorphan hydrobromide 10 mg

Guaifenesin 200 mg

Phenylephrine hydrochloride 5 mg

***Purposes***

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

***Uses***

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves these symptoms due to a cold or flu:
  - nasal congestion · sinus congestion and pressure
  - minor aches and pains · headache
  - cough · sore throat
  - temporarily reduces fever

***Warnings***

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

· 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

· skin reddening · blisters · rash · hives  
· facial swelling · asthma (wheezing) · shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

***Do not use***

● with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

● if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

● if you have ever had an allergic reaction to this product or any of its ingredients

● in children under 12 years of age

**Ask a doctor before use if you have**

● liver disease ● heart disease ● high blood pressure

● thyroid disease ● diabetes

● difficulty in urination due to enlargement of the prostate gland

● persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

● cough with excessive phlegm (mucus)

**Ask a doctor or pharmacist before use if** you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

**Stop use and ask a doctor if**

· pain, cough, or nasal congestion gets worse or lasts more than 7 days

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with fever, rash or headache that lasts.

These could be signs of a serious condition.

- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

### ***Directions***

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

## **Other information**

### ***Other information***

store at 15-25°C (59-77°F)

***Inactive ingredients*** FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol-sorbitan solution, titanium dioxide

## **Questions or comments?**

***Questions or comments?*** 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)

**Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Night Liquid Gels**

### ***Drug Facts***

#### ***Active ingredients (in each capsule)***

Acetaminophen 325 mg

Dextromethorphan hydrobromide 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine hydrochloride 5 mg

## **Purposes**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

## **Uses**

- temporarily relieves these symptoms due to a cold or flu:
- minor aches and pains · headache
- nasal and sinus congestion · cough
- sore throat · runny nose · sneezing
- temporarily reduces fever

## **Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening · blisters · rash · hives
- facial swelling · asthma (wheezing) · shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use to sedate children.**

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional

conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

- in children under 12 years of age

**Ask a doctor before use if you have**

- liver disease ● heart disease ● high blood pressure
- thyroid disease ● diabetes ● glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help

or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- do not take more than the recommended dose

- adults and children 12 years and over: take 2 capsules with water

every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.

- children under 12 years: do not use

***Other information***

- store at 15-25 C 59-77 F

***Inactive ingredients*** FD&C blue #1, D&C yellow #10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

***Questions or comments?*** 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)



**Do not take these products at the same time.**

**Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Day Liquid Gels**

**Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Night Liquid Gels**

**Drug Facts**

**Active ingredients (in each capsule) Purposes**

Acetaminophen 250 mg..... Pain reliever/fever reducer  
 Dextromethorphan hydrobromide 10 mg..... Cough suppressant  
 Guaifenesin 200 mg..... Expectorant  
 Phenylephrine hydrochloride 5 mg..... Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves these symptoms due to a cold or flu:
  - nasal congestion
  - sinus congestion and pressure
  - minor aches and pains
  - headache
  - cough
  - sore throat
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

**Drug Facts**

**Active ingredients (in each capsule) Purposes**

Acetaminophen 325 mg..... Pain reliever/fever reducer  
 Dextromethorphan hydrobromide 10 mg..... Cough suppressant  
 Doxylamine succinate 6.25 mg..... Antihistamine  
 Phenylephrine hydrochloride 5 mg..... Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - nasal and sinus congestion
  - cough
  - sore throat
  - runny nose
  - sneezing
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take
 

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product



● with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

## Cut Edge

8 1/2 E

### Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Day Liquid Gels

#### Drug Facts (continued)

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

#### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough with excessive phlegm (mucus)

### Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Night Liquid Gels

#### Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use to sedate children.**

#### Do not use


- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

#### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

#### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers



## ALKA-SELTZER PLUS SEVERE COUGH MUCUS AND CONGESTION DAY AND NIGHT

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride, guaifenesin kit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0280-0506
---------------------	----------------	---------------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-0506-20	1 in 1 CARTON; Type 0: Not a Combination Product	05/30/2017	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 BLISTER PACK	12
Part 2	2 BLISTER PACK	8

## Part 1 of 2

# ALKA-SELTZER PLUS SEVERE COUGH, MUCUS AND CONGESTION DAY

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, guaifenesin capsule, liquid filled

## Product Information

Route of Administration ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

## Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	AS;M
Contains			

## Packaging

		Marketing Start	Marketing End
--	--	-----------------	---------------

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 CARTON		
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/12/2017	

## Part 2 of 2

### ALKA-SELTZER PLUS SEVERE COUGH MUCUS AND CONGESTION NIGHT

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled

## Product Information

**Route of Administration** ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1VS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SHELLAC</b> (UNII: 46N107B71O)	

**D&C YELLOW NO. 10** (UNII: 35SW5USQ3G)

**POVIDONE** (UNII: FZ989GH94E)

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	AS;Nite
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 CARTON		
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/12/2017	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/30/2017	

**Labeler** - Bayer HealthCare LLC. (112117283)

Revised: 12/2023

Bayer HealthCare LLC.