

**MEDIQUE MEDI SELTZER- aspirin, citric acid, sodium bicarbonate tablet, effervescent**  
**Unifirst First Aid Corporation**

-----  
**Medique Medi Seltzer**

***Drug Facts***

***Active ingredient (in each tablet)***

Aspirin (NSAID\*) 325 mg

Citric Acid 1000 mg

Sodium bicarbonate (heat-treated) 1916 mg

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever /fever reducer

Antacid

Antacid

***Uses***

**For the temporary relief of**

- heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin,

ibuprofen, naproxen, or others)

- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/ fever reducer
- if you are taking a prescription drug for gout, diabetes or arthritis

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you are on a sodium restricted diet
- you reached age 60 or older

### **Ask a doctor or pharmacist before use if you are**

- taking any other drugs (prescription or non-prescription). Antacids may interact with certain drugs.
- under a doctor's care for any serious condition

### **When using this product do not exceed recommended dosage.**

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - you feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin (**\*NSAIDS**) **at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- do not exceed recommended dosage
- to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking

**Adults and children 12 to 60 years of age:** 2 tablets every 4 hours. Do not exceed 8 tablets in 24 hours or do not use the maximum dosage for more than 10 days.

**Children under 12 years of age:** Do not give to children under 12 years of age.

60 years and older: Do not exceed 4 tablets in 24 hours or do not use the maximum dosage for more than 10 days.

**Other information**

- each tablet contains: Sodium 524 mg
- store at room temperature 59°-86° F (15°-30° C)
- read all product information before using
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

none

**Questions or comments?** 1-800-634-7680

**Medique Medi-Seltzer Label**

Medique®

Medi-Seltzer

Effervescent Antacid/Pain Reliever

Pull to Open

This Package is for Households without Young Children.

Aches, Fever ● Aspirin (NSAID) 325 mg

Antacid ● Citric Acid 1000 mg Sodium Bicarbonate 1916 mg

72 Tablets

(36 x 2)

Tamper Evident Unit Dose Packets

**Medique®**  
**Medi-Seltzer**  
Effervescent Antacid/  
Pain Reliever

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

Collect Medibucks from certain large size Medique Brand products and redeem them for great gifts & savings.  
www.mediqueproducts.com

15 MEDIBUCKS 15

**Medique®**  
**Medi-Seltzer**  
Effervescent Antacid/  
Pain Reliever

Pull to Open

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

72 Tablets (36 x 2)

Tamper Evident Unit Dose Packets

**Drug Facts**

**Active ingredient (in each tablet)** Purpose  
Aspirin (NSAID) 325 mg \_\_\_\_\_ Pain reliever/fever reducer  
Citric Acid 1000 mg \_\_\_\_\_ Antacid  
Sodium bicarbonate (heat-treated) 1916 mg \_\_\_\_\_ Antacid  
Nonsteroidal anti-inflammatory drug

**Uses**  
For the temporary relief of:  
● heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains  
● upset stomach with headache from overindulgence in food or drink  
● headache, body aches, and pain alone

**Warnings**  
Riley's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Riley's syndrome, a rare but serious illness.

**Allergy Alert:** Aspirin may cause a severe allergic reaction which may include:  
● hives ● facial swelling  
● asthma (wheezing) ● shock  
If an allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Stomach Bleeding Warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is higher if you:  
● are age 60 or older  
● have had stomach ulcers or bleeding problems  
● take a blood thinning (anticoagulant) or steroid drug  
● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
● have 3 or more alcohol drinks every day while using this product  
● take more or for a longer time than directed

**Do not use**  
● if you have ever had an allergic reaction to any other pain reliever/fever reducer  
● if you are taking a prescription drug for gout, diabetes or arthritis

**Ask a doctor before use if**  
● stomach bleeding warning applies to you  
● you have a history of stomach problems such as heartburn  
● you have high blood pressure, heart disease, liver disease, or kidney disease  
● you are taking a diuretic  
● you are on a sodium restricted diet  
● you reached age 60 or older  
(continued on the next panel)

**Directions**  
2 tablets every 4 hours. Do not exceed 8 tablets in 24 hours or do not use the maximum dosage for more than 10 days.  
Children under 12 years of age: Do not give to children under 12 years of age.  
60 years and older: Do not exceed 4 tablets in 24 hours or do not use the maximum dosage for more than 10 days.

**Other information**  
● Each tablet contains: Sodium 524mg  
● Store at room temperature 59°-86° F (15°-30° C)  
● Read all product information before using  
● Avoid excessive heat and humidity  
● Tamper evident sealed packets  
● Do not use any opened or torn packets

**Inactive ingredients** None

**Questions or comments:** 1-800-634-7680

**Retain carton for complete product information**

Reorder #13524  
Manufactured for  
Medique Products  
Fort Myers, FL 33967  
1-800-634-7680  
www.mediqueproducts.com

3 47682 13524 9

# MEDIQUE MEDI SELTZER

aspirin, citric acid, sodium bicarbonate tablet, effervescent

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-135
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1ZRA3Q20)	SODIUM BICARBONATE	1916 mg

## Product Characteristics

<b>Color</b>	white (white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	25mm
<b>Flavor</b>		<b>Imprint Code</b>	L421
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-135-20	18 in 1 BOX	12/30/2008	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-135-24	36 in 1 BOX	12/30/2008	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-135-60	300 in 1 BOX	12/30/2008	
3		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/30/2008	

**Labeler** - Unifirst First Aid Corporation (832947092)

