

MECLIZINE HYDROCHLORIDE- meclizine hydrochloride tablet

NCS HealthCare of KY, LLC dba Vanguard Labs

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use MECLIZINE HYDROCHLORIDE TABLETS safely and effectively. See full prescribing information for MECLIZINE HYDROCHLORIDE TABLETS.

MECLIZINE hydrochloride tablets, for oral use
Initial U.S. Approval: 1957

INDICATIONS AND USAGE

Meclizine hydrochloride tablets are indicated for the treatment of vertigo associated with diseases affecting the vestibular system in adults (1). (1)

DOSAGE AND ADMINISTRATION

- Recommended dosage: 25 mg to 100 mg daily, in divided doses (2.1).
- Tablets: Swallow whole (2.2).

DOSAGE FORMS AND STRENGTHS

- Tablets: 12.5 mg and 25 mg (3).

CONTRAINDICATIONS

Meclizine hydrochloride tablets are contraindicated in patients with hypersensitivity to meclizine or any of the inactive ingredients (4). (4)

WARNINGS AND PRECAUTIONS

- May cause drowsiness: Use caution when driving a car or operating dangerous machinery (5.1).
- Potential anticholinergic action: this drug should be prescribed with care to patients with a history of asthma, glaucoma, or enlargement of the prostate gland (5.2).

ADVERSE REACTIONS

Common adverse reactions are anaphylactic reaction, drowsiness, dry mouth, headache, fatigue, and vomiting. On rare occasions blurred vision has been reported (6). (6)

To report SUSPECTED ADVERSE REACTIONS, contact Amneal Pharmaceuticals at 1-877-835-5472 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. (6)

DRUG INTERACTIONS

- Co-administration of meclizine hydrochloride with other CNS depressants, including alcohol, may result in increased CNS depression (7.1).
- CYP2D6 inhibitors: As meclizine is metabolized by CYP2D6, there is a potential for drug-drug interactions between meclizine hydrochloride and CYP2D6 inhibitors (7.2).

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 5/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

2.2 Administration Instructions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Drowsiness
- 5.2 Concurrent Medical Conditions

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

- 7.1 CNS Depressants
- 7.2 CYP2D6 Inhibitors

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment
- 8.7 Renal Impairment
- 8.8 Genetic CYP2D6 Polymorphism

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Meclizine hydrochloride tablets are indicated for the treatment of vertigo associated with diseases affecting the vestibular system in adults.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

The recommended dosage is 25 mg to 100 mg daily administered orally, in divided doses, depending upon clinical response.

2.2 Administration Instructions

Tablets

Meclizine hydrochloride tablets must be swallowed whole.

3 DOSAGE FORMS AND STRENGTHS

- Meclizine hydrochloride tablets USP, 12.5 mg are light blue colored, oval shaped tablets with “AN 441” debossed on one side and plain on the other side.
- Meclizine hydrochloride tablets USP, 25 mg are light yellow colored, oval shaped tablets with “AN 442” debossed on one side and plain on the other side.

4 CONTRAINDICATIONS

Meclizine hydrochloride tablets are contraindicated in patients with a hypersensitivity to meclizine or any of the inactive ingredients [see *Adverse Reactions (6)* and *Description (11)*].

5 WARNINGS AND PRECAUTIONS

5.1 Drowsiness

Since drowsiness may occur with use of meclizine hydrochloride, patients should be warned of this possibility and cautioned against driving a car or operating dangerous machinery.

Patients should avoid alcoholic beverages while taking meclizine hydrochloride [see *Drug Interactions (7.1)*].

5.2 Concurrent Medical Conditions

Because of its potential anticholinergic action, meclizine hydrochloride should be used with caution in patients with asthma, glaucoma, or enlargement of the prostate gland.

6 ADVERSE REACTIONS

The following adverse reactions associated with the use of meclizine hydrochloride were identified in clinical studies or post-marketing reports. Because some of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Anaphylactic reaction, drowsiness, dry mouth, headache, fatigue, and vomiting. On rare occasions blurred vision has been reported.

7 DRUG INTERACTIONS

7.1 CNS Depressants

There may be increased CNS depression when meclizine hydrochloride is administered concurrently with other CNS depressants, including alcohol [see *Warnings and Precautions (5.1)*].

7.2 CYP2D6 Inhibitors

Based on *in-vitro* evaluation, meclizine is metabolized by CYP2D6. Therefore, there is a

possibility for a drug interaction between meclizine hydrochloride and CYP2D6 inhibitors. Therefore, monitor for adverse reactions and clinical effect accordingly.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Data from epidemiological studies have not generally indicated a drug-associated risk of major birth defects with meclizine during pregnancy. However, in a published study, an increased incidence of fetal malformations was observed following oral administration of meclizine to pregnant rats during the period of organogenesis, at doses similar to those used clinically. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. The background risk of major birth defects and miscarriage for the indicated population is unknown.

Data

Human Data

Epidemiological studies reporting on pregnancies exposed to meclizine have not identified an association between the use of meclizine during pregnancy and an increased risk of major birth defects.

Animal Data

In a published study, oral administration of meclizine (25 mg/kg to 250 mg/kg) to pregnant rats during the period of organogenesis resulted in a high incidence of fetal malformations. These effects occurred at doses as low as 25 mg/kg, which is approximately 2 times the maximum recommended human dose (100 mg) on a body surface area (mg/m²) basis.

8.2 Lactation

Risk Summary

There are no data on the presence of meclizine in human milk, the effects on the breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for meclizine hydrochloride and any potential adverse effects on the breastfed infant from meclizine hydrochloride or from the underlying maternal condition.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

8.5 Geriatric Use

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

8.6 Hepatic Impairment

The effect of hepatic impairment on the pharmacokinetics of meclizine has not been evaluated. As meclizine hydrochloride undergoes metabolism, hepatic impairment may result in increased systemic exposure of meclizine. Treatment with meclizine hydrochloride should be administered with caution in patients with hepatic impairment.

8.7 Renal Impairment


The effect of renal impairment on the pharmacokinetics of meclizine has not been evaluated. Because of a potential for drug/metabolite accumulation, meclizine hydrochloride should be administered with caution in patients with renal impairment and in the elderly, as renal function generally declines with age.

8.8 Genetic CYP2D6 Polymorphism

The genetic polymorphism of CYP2D6 that results in poor-, intermediate-, extensive-, and ultrarapid metabolizer phenotypes could contribute to large inter-individual variability in meclizine exposure. Therefore, when meclizine hydrochloride is administered to patients with CYP2D6 polymorphism, monitor for adverse reactions and clinical effect accordingly.

11 DESCRIPTION

Meclizine hydrochloride, a histamine (H₁) receptor antagonist, is a white or slightly yellowish, crystalline powder. Its molecular formula is C₂₅H₂₇ClN₂·2HCl·H₂O and its molecular weight is 481.88. It has the following structural formula:



Chemically, meclizine hydrochloride is 1-(*p*-chloro- α -phenylbenzyl)-4-(*m*-methylbenzyl) piperazine dihydrochloride monohydrate.

Each meclizine hydrochloride 12.5 mg tablet contains 12.5 mg of meclizine dihydrochloride equivalent to 10.53 mg of meclizine free base.

Each meclizine hydrochloride 25 mg tablet contains 25 mg of meclizine dihydrochloride equivalent to 21.07 mg of meclizine free base.

Inactive ingredients for the tablets are: colloidal silicon dioxide, lactose monohydrate, magnesium stearate, microcrystalline cellulose, sodium starch glycolate and talc. The 12.5 mg tablets also contain FD&C Blue #1 Aluminum Lake. The 25 mg tablets also

contain D&C Yellow #10 Aluminum Lake.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The precise mechanism by which meclizine exerts its therapeutic effect is unknown but is presumed to involve antagonism of the histamine H1 receptor.

12.2 Pharmacodynamics

There are no relevant pharmacodynamic data regarding meclizine.

12.3 Pharmacokinetics

The available pharmacokinetic information for meclizine following oral administration has been summarized from published literature.

Absorption

Meclizine is absorbed after oral administration with maximum plasma concentrations reaching at a median T_{max} value of 3 hours post-dose (range: 1.5 to 6 hours) for the tablet dosage form.

Distribution

Drug distribution characteristics for meclizine in humans are unknown.

Elimination

Meclizine has a plasma elimination half-life of about 5 to 6 hours in humans.

Metabolism In an *in vitro* metabolic study using human hepatic microsome and recombinant CYP enzyme, CYP2D6 was found to be the dominant enzyme for metabolism of meclizine.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Animal studies to assess the carcinogenic potential of meclizine have not been conducted.

Mutagenesis

Genetic toxicology studies of meclizine have not been conducted.

Impairment of Fertility

Animal studies to assess the effects of meclizine on fertility and early embryonic development have not been conducted.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Meclizine Hydrochloride Tablets USP, **12.5 mg** are supplied as light blue colored, oval shaped tablets with “AN 441” debossed on one side and plain on the other side.

Meclizine Hydrochloride Tablets USP, **25 mg** are supplied as light yellow colored, oval shaped tablets with “AN 442” debossed on one side and plain on the other side.

They are available as follows:

Blistercards of 30: NDC 0615-8462-39

16.2 Storage and Handling

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP.

Keep this and all medication out of the reach of children.

17 PATIENT COUNSELING INFORMATION

Administration Instructions

Advise patients that the tablets must be swallowed whole [see *Dosage and Administration (2.1)*].

Adverse Reactions

Advise patients that meclizine hydrochloride may cause anaphylactic reaction, drowsiness, dry mouth, headache, fatigue, vomiting and, on rare occasions, blurred vision [see *Warnings and Precautions (5.1)*, *Adverse Reactions (6)*].

Inform patients that meclizine hydrochloride may impair their ability to engage in potentially dangerous activities, such as operating machinery or vehicles.

Concomitant Drug Interactions

Advise patients regarding medications that should not be taken in combination with meclizine hydrochloride or that may necessitate increased monitoring [see *Drug Interactions (7.1, 7.2)*]. Inform patients that alcohol may increase adverse reactions.

Concurrent Medical Conditions

Advise patients to notify their healthcare provider about all of their medical conditions, including if they are pregnant or plan to become pregnant or if they are breastfeeding [see *Warnings and Precautions (5.2)*, *Use in Specific Populations (8.1, 8.2)*].

Manufactured by:

Amneal Pharmaceuticals Pvt. Ltd.

Oral Solid Dosage Unit

Ahmedabad 382213, INDIA

Distributed by:

Amneal Pharmaceuticals LLC

Bridgewater, NJ 08807

Rev. 08-2021-00

PRINCIPAL DISPLAY PANEL

VLI NDC 0615-8462-39
Meclizine Hydrochloride
Tabs USP 25 mg*
LOT 8462-
EXP
8462-AA-39 v00
QTY
30
Mfg By Amneal
(NDC 53746-442-10)
PKG BY VANGARD
GLASGOW, KY 42141

(Amneal NDC 53746-442-10)
Meclizine Hydrochloride
Tabs USP 25 mg*

Received: _____

24	16	8
31	23	15
30	22	14
29	21	13
28	20	12
27	19	11
26	18	10
25	17	9
Start Date	Start Time	

STORE AT 20° - 25°C (68° - 77° F)
(SEE USP CONTROLLED ROOM TEMPERATURE)

Dispense in a tight, light-resistant container.

*Each tab contains: Meclizine dihydrochloride, USP 25 mg (Eq. to 21.07 mg of meclizine free base.)

See package insert or label for dosage information
FOR INSTITUTIONAL USE ONLY

16	8
30	23
29	22
28	21
27	20
26	19
25	18
24	17
23	16
22	15
21	14
20	13
19	12
18	11
17	10
16	9
15	8
14	7
13	6
12	5
11	4
10	3
9	2
8	1

Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	8462-AA-B-v00 Vanguard Labs Glasgow, KY 42141 LOT 8462 - EXP 5374644210
Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210
Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210
Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210
Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210
Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210
Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210	Pkg by Vanguard, Glasgow, KY 42141 Meclizine Hydrochloride Tab USP 25 mg* LOT 8462 - EXP 5374644210

Omnicare 24 The overall configuration of this package is a trademark of Omnicare, Inc. 17 9 1

MECLIZINE HYDROCHLORIDE

meclizine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0615-8462(NDC:53746-442)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TALC (UNII: 7SEV7J4R1U)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	

Product Characteristics

Color	yellow (light)	Score	no score
Shape	OVAL	Size	13mm
Flavor		Imprint Code	AN;442
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0615-8462-39	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	04/26/2023	01/31/2025

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201451	04/22/2022	01/31/2025

Labeler - NCS HealthCare of KY, LLC dba Vanguard Labs (050052943)

Establishment

Name	Address	ID/FEI	Business Operations
NCS HealthCare of KY, LLC dba Vanguard Labs		050052943	repack(0615-8462)

Revised: 5/2023

NCS HealthCare of KY, LLC dba Vanguard Labs