HEMATRON-AF- ascorbic acid, dl-alpha tocopheryl acetate, l-5methyltetrahydrofolate calcium, cyanocobalamin, biotin, carbonyl iron, cupric sulfate tablet Sever Pharmatec, Inc.

\_\_\_\_\_

#### **Hematron-AF**<sup>™</sup>

Supplement Facts Serving Size: 1 Tablet / Servings Per Container: 90				
	Amount	% Daily		
	Per Serving	Value		
Vitamin C (as Ascorbic Acid)	500 mg	556%		
Vitamin E (as dl-Alpha Tocopheryl Acetate)	13.5 mg	90%		
Folate (as L-5-Methyltetrahydrofolate Calcium)	1,700 mcg DFE	425%		
	(1,000 mcg Folic Acid)			
Vitamin B12 (as Cyanocobalamin)	60 mcg	2,500%		
Biotin	150 mcg	500%		
Iron (as Carbonyl Iron)	150 mg	833%		
Copper (as Cupric Sulfate)	3 mg	333%		
*Daily Value not established				

#### Other Ingredients:

Microcrystalline Cellulose, Citric Acid, Stearic Acid, Dicalcium Phosphate, Croscarmellose Sodium, Polyvinylpyrrolidone, Magnesium Stearate, Silicon Dioxide, Starch, Glucose Syrup, Crospovidone, Polyethylene Glycol, Talc, Sodium Lauryl Sulfate, FD&C Yellow #6 Lake, Sucrose, Titanium Dioxide, Soy Polysaccharide, FD&C Red #40 Lake, Artificial French Vanilla Flavor, Propylene Glycol and Tricalcium Phosphate. **Contains:** Soy, a bioengineered food ingredient.

**HEMATRON AF** is a multi-vitamin / multi-mineral dietary supplement indicated for use in improving the nutritional status of patients with iron deficiency<sup>†</sup>

<sup>†</sup>This statement has not been evaluated by the FDA. This product is not intended to diagnose, treat, cure, or prevent any disease.

**WARNING** Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6 years of age. **KEEP THIS PRODUCT OUT OF REACH OF CHILDREN**. In case of accidental overdose, call a doctor or poison control center immediately.

**DO NOT USE** with individuals with a known hypersensitivity to any of the ingredients. Patients with hemochromatosis and hemosiderosis are contra-indications to iron therapy. This product is not for use in children under 12 years of age.

## CAUTION

This product contains 1 mg folic acid, the maximum safe Reference Daily Intake (RDI) amount for this nutrient. Folic Acid alone should not be used for individuals with pernicious anemia and other megaloblastic anemias where Vitamin B-12 is deficient. Folic acid in amounts above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations progress. Pregnant Women, nursing mothers or women prior to conception who take **HEMATRON AF** should take this supplement only after their medical condition and other drugs, herbs and / or supplements consumption are considered. Use for elderly individuals should be cautious due to the greater frequency of decreased hepatic, renal, or cardiac function.

### **ADVERSE REACTIONS:**

Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

### DIRECTIONS FOR USE

One caplet daily with or without food or as prescribed by your healthcare provider.

#### STORAGE

Store at room temperature, USP.

To report a serious adverse event or to obtain product information, contact (888) 444-5072.

Manufactured in the USA for: Seyer Pharmatec, Inc. Caguas, Puerto Rico 00726

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

**Dietary Supplement** 

EZY-DOSE TABLETS

#### Packaging



Vitamin E (as dl-Alp Folate (as L-5-Mett drofolate Calcium)

Vitamin B12 (as (

Vitamin C (as Aso

From (as Carbonyl Ir Copper (as Cupric ) \*Daily Value not es

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Supplement Facts Serving Size: 1 Tablet / Servings Per Container: 90

	F	Amount Per Serving	% Daily Value
Vitamin C (as Ascorbic Acid)		500 mg	556%
Vitamin E (as dl-Alpha Tocopher	Acetate)	13.5 mg	90%
Folate (as L-5-Methyltetrahy	1,70	0 mcg DFE	425%
drofolate Calcium)	1,000 mcg	Folic Acid)	
Vitamin B12 (as Cyanocobalamir	1)	60 mcg	2,500%
Biotin		150 mcg	500%
ron (as Carbony Iron)		150 mg	833%
Copper (as Cupric Sulfate)		3 mg	333%
*Daily Value not established			

Other Ingredients: Microcrystalline Cellulose, Citric Acid, Stearic Acid, Dicalcium Phosphate, Croscarmellose Sodium, Polyvinylpyrrolidone, Magnesium Stearate, Silicon Dioxide, Starch, Glucose Syrup, Crospovidone, Polyethylene Glycol, Talc, Sodium Lauryl Sulfate, FD&C Yellow #6 Lake, Sucrose, Titanium Dioxide, Soy Polysaccharide, FD&C Red #40 Lake, Artificial French Vanilla Havor, Propylene Glycol and Tricalcium Phosphate. Contains: Soy, a bioengineered food ingredient.



**HEMATRON AF** is a multi-vitamin / muliti-mineral dietary supplement indicated for use in improving the nutritional status of patients with iron deficiency<sup>†</sup>

<sup>1</sup>This statement has not been evaluated by the FDA. Thhis product in not intended to diagnose, treat, cure, or prevent any disease.

WARNING Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6 years of age. KEEP THIS PRODUCT OUT OF REACH OF CHILDREN. In case of accidental overdose, call a doctor or poison control center immediately.

**DO NOT USE** with individuals with a known hypersensitivity to any of the ingredients. Patients with hemochromatosis and hemosiderosis are contra-indications to iron therapy. This product is not for use in children under 12 years of age.

**CAUTION** This product contains 1 mg folic acid, the maximum safe Reference Daily Intake (RDI) amount for this nutrient. Folic Acid alone should not be used for individuals with pernicious anemia and other megaloblastic anemias where Vitamin B-12 is deficient. Folic acid in amounts above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations progress. Pregnant women, nursing mothers or women prior to conception who take **HEMATRON AF** should take this supplement only after their medical condition and other drugs, herbs and /or supplements consumption are considered. Use for elderly individuals should be cautious due to the greater frequency of decreased hepatic, renal, or cardiac function.

ADVERSE REACTIONS Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

**DIRECTIONS FOR USE** One caplet daily with or without food or as prescribed by your healthcare provider.

STORAGE Store at room temperature, USP

To report a serious adverse event or to obtain product information, contact (888) 444-5072.

Code# SEY2755-9 L Rev 12/21

Manufactured in the USA for:



Seyer Pharmatec, Inc. Caguas, Puerto Rico 00726

# **HEMATRON-AF**

ascorbic acid, dl-alpha tocopheryl acetate, l-5-methyltetrahydrofolate calcium, cyanocobalamin, biotin, carbonyl iron, cupric sulfate tablet

Product Information				
Product Type	DIETARY SUPPLEMENT	Item Code (Source)	NHRIC:11026-2760	
Route of Administration	ORAL			

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASCORBIC ACID (UNII: PQ6CK8PD0R) (ASCORBIC ACID - UNII:PQ6CK8PD0R)	ASCORBIC ACID	500 mg
<b>.ALPHATOCOPHEROL ACETATE, DL-</b> (UNII: WR1WPI7EW8) (.ALPHATOCOPHEROL, DL UNII:7QWA1RIO01)	.ALPHA TOCOPHEROL, DL-	13.5 mg
<b>LEVOMEFOLATE CALCIUM</b> (UNII: A9R10K3F2F) (LEVOMEFOLIC ACID - UNII:8S95DH25XC)	LEVOMEFOLATE CALCIUM	1700 ug
CYANOCOBALAMIN (UNII: P6YC3EG204) (CYANOCOBALAMIN - UNII:P6YC3EG204)	CYANOCOBALAMIN	60 ug
BIOTIN (UNII: 6SO6U10H04) (BIOTIN - UNII:6SO6U10H04)	BIOTIN	150 ug
IRON (UNII: E1UOL152H7) (IRON - UNII:E1UOL152H7)	IRON	150 mg
CUPRIC SULFATE (UNII: LRX7AJ16DT) (CUPRIC CATION - UNII:8CBV67279L)	CUPRIC CATION	3 mg

### **Inactive Ingredients**

Ingredier	nt Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D6	61U)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UN	III: L11K75P92J)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)		
POVIDONE K60 (UNII: SZR7Z3Q2YH)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
STARCH, CORN (UNII: 08232NY3SJ)		
CORN SYRUP (UNII: 9G5L16BK6N)		
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E56	1)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3W	JQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)		
SODIUM LAURYL SULFATE (UNII: 368GB5141J)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
SUCROSE (UNII: C151H8M554)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		
SOYBEAN COTYLEDON CELL WALL POLYSACCH	ARIDES (UNII: 4UL6DF56YQ)	
FD&C RED NO. 40 (UNII: WZ B9127XOA)		

	PYLENE GLYCOL (U	JNII: 6DC9Q167V3)				
TRIC	ALCIUM PHOSPHA	<b>TE</b> (UNII: K4C08XP666)				
Pac	ckaging					
#	ltem Code	<b>Package Description</b>	Marketin	g Start Date	Marl	keting End Date
<b>1</b> N	HRIC:11026-2760-9	90 in 1 BOTTLE				
Ma	rketing Info	ormation				
	rketing Info					
	<b>rketing Info</b> Marketing Category	<b>Ormation</b> Application Number or Mo Citation	onograph	Marketing S Date	tart	Marketing End Date

Supplement Facts				
Serving Size :		Serving per Container :		
	Amount Per Serving	% Daily Value		
color				
shape				
size (solid drugs)	21 mm			
scoring	1			
imprint				
flavor				

Labeler - Seyer Pharmatec, Inc. (832947126)

Revised: 2/2023

Seyer Pharmatec, Inc.