SUNMARK ASPIRIN- as pirin tablet, chewable Strategic Sourcing Services LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

McKesson Aspirin Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

• store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, D&C red #27 aluminum lake, dextrose excipient, FD&C red #40 aluminum lake, flavors, saccharin sodium

Questions or comments?

1-800-719-9260

Principal Display Panel

COMPARE TO BAYER $^{\circledR}$ CHEWABLE LOW DOSE ASPIRIN ACTIVE INGREDIENT aspirin

Pain Reliever (NSAID)

Low Dose Aspirin Regimen

Talk to your doctor or other healthcare provider before using this product for your heart.

CHEWABLE CHERRY FLAVORED

Actual Size

GLUTEN FREE

36 TABLETS – 81 mg



Drug Facts

Active ingredient Purpose (in each tablet)

Aspirin 81 mg (NSAID)*. ..Pain reliever *nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur. consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare butserious illness.

Allergy alert: Aspirin may cause a severe aller gic reaction which may include:

■ hives ■ facial swelling
■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product
contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding
- take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more orfora longer time than directed

Do notuse if you have ever had an allergic reaction to any other pain reliever/fever reducer

■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

- Stop use and ask a doctor if
- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint vomit blood
- have bloody or black stools
- have stomach pain that does not get better ▶

Drug Facts (continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present new symptoms occur
- ringing in the ears or a loss of hearing occurs These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center rightaway (1-800-222-1222).

- drink a full glass of water with each dose
- adults and children 12 years and over:take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

■ store at 20-25°C (68-77°F)

MCKESSON

Questions or comments?

Drug Inactive in gred ients om starch, D& C red #27 aluminum lale, dextros e excipient FD&C red #4 Oaluminum lake flavors, sæcharin sodium

Facts (continued)

SUNMARK ASPIRIN

aspirin tablet, chewable

Prod	luct	Info	rma	tion
Proc	luct	11110	miliai	uon

Product Type HUMAN OTC DRUG Item Code (Source) NDC:49348-191

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients Ingredient Name Strength STARCH, CORN (UNII: O8232NY3SJ) D&C RED NO. 27 (UNII: 2LRS185U6K) FD&C RED NO. 40 (UNII: WZB9127XOA) SACCHARIN SODIUM (UNII: SB8ZUX40TY)

Product Characteristics				
Color	PINK	Score	no score	
Shape	ROUND	Size	8 mm	
Flavor	CHERRY	Imprint Code	L274	
Contains				

Packaging				
# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1 NDC:49348-191-07	1 in 1 CARTON	09/06/2003		
1	36 in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part343	09/06/2003		

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 2/2020 Strategic Sourcing Services LLC