

**DERACOXIB CHEWABLE- deracoxib tablet**  
**BUTLER ANIMAL HEALTH SUPPLY, LLC DBA COVETRUS NORTH AMERICA**

-----  
**Deracoxib**

**Chewable Tablets**

**For Oral Use in Dogs Only**


**Do Not Use in Cats**

**Caution:**

Federal Law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

**Description:**

Deracoxib Chewable Tablets is a non-narcotic, non-steroidal anti-inflammatory drug (NSAID) of the coxib class. Deracoxib Chewable Tablets are round, light brown colored, chewable tablets that contain deracoxib formulated with beefy flavoring. The molecular weight of deracoxib is 397.38. The empirical formula is  $C_{17}H_{14}F_3N_3O_3S$ . Deracoxib is 4-[3-(difluoromethyl)-5-(3-fluoro-4-methoxyphenyl)-1H-pyrazole-1-yl] benzenesulfonamide, and can be termed a diaryl substituted pyrazole. The structural formula is:



**Clinical Pharmacology**

**Mode of Action:**

Deracoxib Chewable Tablets are a member of the coxib class of non-narcotic, nonsteroidal, cyclooxygenase-inhibiting anti-inflammatory drugs for the control of postoperative pain and inflammation associated with orthopedic and dental surgery and for the control of pain and inflammation associated with osteoarthritis in dogs.

Data indicate that deracoxib inhibits the production of PGE<sub>1</sub> and 6-keto PGF<sub>1</sub> by its

inhibitory effects on prostaglandin biosynthesis.<sup>1</sup> Deracoxib inhibited COX-2 mediated PGE2 production in LPS-stimulated human whole blood.<sup>2</sup>

Cyclooxygenase-1 (COX-1) is the enzyme responsible for facilitating constitutive physiological processes (e.g., platelet aggregation, gastric mucosal protection, renal perfusion).<sup>3</sup> Cyclooxygenase-2 (COX-2) is responsible for the synthesis of inflammatory mediators.<sup>4</sup> Both COX isoforms are constitutively expressed in the canine kidney.<sup>5</sup> At doses of 2-4 mg/kg/day, deracoxib tablets do not inhibit COX-1 based on *in vitro* studies using cloned canine cyclooxygenase.<sup>6</sup> The clinical relevance of this *in vitro* data has not been shown.

Although the plasma terminal elimination half-life for deracoxib tablets is approximately 3 hours, a longer duration of clinical effectiveness is observed.

Summary pharmacokinetics of deracoxib tablets are listed in Table 1.

**Table 1: Pharmacokinetics of Deracoxib**

<b>Parameter</b>	<b>Value</b>
Tmax <sup>a</sup>	2 hours
Oral Bioavailability (F) <sup>a</sup>	> 90% at 2 mg/kg
Terminal elimination half-life <sup>b</sup>	3 hours at 2-3 mg/kg 19 hours at 20 mg/kg
Systemic Clearance <sup>b</sup>	~ 5 ml/kg/min at 2 mg/kg ~1.7 ml/kg/min at 20 mg/kg
Volume of Distribution <sup>c</sup>	~ 1.5 L/kg
Protein binding <sup>d</sup>	> 90%

<sup>a</sup>Values obtained following a single 2.35 mg/kg dose

<sup>b</sup>Estimates following IV administration of deracoxib as an aqueous solution

<sup>c</sup>Based upon a dose of 2 mg/kg of deracoxib

<sup>d</sup>Based upon *in vitro* plasma concentrations of 0.1, 0.3, 1.0, 3.0, 10.0 µg/ml

Non-linear elimination kinetics are exhibited at doses above 8 mg/kg/day, at which competitive inhibition of constitutive COX-1 may occur.

Deracoxib is not excreted as parent drug in the urine. The major route of elimination of deracoxib is by hepatic biotransformation producing four major metabolites, two of which are characterized as products of oxidation and o-demethylation. The majority of deracoxib is excreted in feces as parent drug or metabolite.

Large intersubject variability was observed in drug metabolite profiles of urine and feces. No statistically significant differences between genders were observed.

### **Indications and Usage:**

**Always provide “Information for Dog Owners” Sheet with prescription. Carefully consider the potential benefits and risk of Deracoxib Chewable**

**Tablets and other treatment options before deciding to use Deracoxib Chewable Tablets. Use the lowest effective dose for the shortest duration consistent with individual response.**

Osteoarthritis Pain and Inflammation:

Deracoxib Chewable Tablets are indicated for the control of pain and inflammation associated with osteoarthritis in dogs.

**Dosage and Administration:**

**Osteoarthritis Pain and Inflammation: 0.45 - 0.91 mg/lb/day (1 to 2 mg/kg/day) as a single daily dose, as needed.**

Dogs needing a dose of less than 12.5 mg can only be accurately dosed through use of the 12 mg tablet, which can be broken in half to provide 6 mg. Do not attempt to accurately dose smaller dogs through the use of breaking larger tablets. **Inaccurate dosing may result in adverse drug events (see Adverse Reactions, Animal Safety, and Post-Approval Experience).**

Postoperative Orthopedic Pain and Inflammation:

Deracoxib Chewable Tablets are indicated for the control of postoperative pain and inflammation associated with orthopedic surgery in dogs.

**Dosage and Administration:**

**Postoperative Orthopedic Pain and Inflammation: 1.4 - 1.8 mg/lb/day (3 to 4 mg/kg/day) as a single daily dose, as needed, not to exceed 7 days of administration.**

Dogs needing a dose of less than 12.5 mg can only be accurately dosed through use of the 12 mg tablet, which can be broken in half to provide 6 mg. Do not attempt to accurately dose smaller dogs through the use of breaking larger tablets. **Inaccurate dosing may result in adverse drug events (see Adverse Reactions, Animal Safety, and Post-Approval Experience).**

Postoperative Dental Pain and Inflammation:

Deracoxib Chewable Tablets are indicated for the control of postoperative pain and inflammation associated with dental surgery in dogs.

**Dosage and Administration:**

**Postoperative Dental Pain and Inflammation: 0.45 - 0.91 mg/lb/day (1 to 2 mg/kg/day) as a single daily dose, for 3 days.**

The first dose should be given approximately 1 hour prior to dental surgery and subsequent doses should be given daily for up to two additional treatments.

Dogs needing a dose of less than 12.5 mg can only be accurately dosed through use of the 12 mg tablet, which can be broken in half to provide 6 mg. Do not attempt to accurately dose smaller dogs through the use of breaking larger tablets. **Inaccurate dosing may result in adverse drug events (see Adverse Reactions, Animal Safety, and Post-Approval Experience).**

Since deracoxib tablet bioavailability is greatest when taken with food, postprandial administration is preferable. However, deracoxib tablets have been shown to be effective under both fed and fasted conditions; therefore, they may be administered in the fasted state if necessary. For postoperative orthopedic and dental pain, administer Deracoxib Chewable Tablets prior to the procedure. Tablets are scored and dosage should be calculated in half-tablet increments. In clinical practice it is recommended to

adjust the individual patient dose while continuing to monitor the dog's status until a minimum effective dose has been reached.

### **Contraindications:**

Dogs with known hypersensitivity to deracoxib should not receive Deracoxib Chewable Tablets.

### **Warnings:**

Not for use in humans. Keep this and all medications out of reach of children. Consult a physician in case of accidental ingestion by humans. **For use in dogs only. Do not use in cats.**

Dogs needing a dose of less than 12.5 mg can only be accurately dosed through use of the 12 mg tablet, which can be broken in half to provide 6 mg. Do not attempt to accurately dose smaller dogs through the use of breaking larger tablets. **Inaccurate dosing may result in adverse drug events (see Adverse Reactions, Animal Safety, and Post-Approval Experience).**

All dogs should undergo a thorough history and physical examination before the initiation of NSAID therapy. Appropriate laboratory tests to establish hematological and serum biochemical baseline data prior to, and periodically during, administration of any NSAID is recommended. **Owners should be advised to observe for signs of potential drug toxicity (see Adverse Reactions, Animal Safety and Post-Approval Experience) and be given an "Information for Dog Owners" Sheet.**

Keep Deracoxib in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

### **Precautions:**

Dogs needing a dose of less than 12.5 mg can only be accurately dosed through use of the 12 mg tablet, which can be broken in half to provide 6 mg. Do not attempt to accurately dose smaller dogs through the use of breaking larger tablets. **Inaccurate dosing may result in adverse drug events (see Adverse Reactions, Animal Safety, and Post-Approval Experience).**

Since NSAIDs possess the potential to produce gastrointestinal ulceration and/or perforation, concomitant use of Deracoxib Chewable Tablets with other anti-inflammatory drugs, such as NSAIDs or corticosteroids, should be avoided. As a class, NSAIDs may be associated with gastrointestinal, renal and hepatic toxicity. The following collective group of clinical signs has been reported with some serious gastrointestinal events, in decreasing order of reported frequency: anorexia, tachycardia, tachypnea, pyrexia, ascites, pale mucous membranes, dyspnea. In some cases, circulatory shock, collapse and cardiac arrest have also been reported. Sensitivity to drug-associated adverse events varies with the individual patient. Dogs that have experienced adverse reactions from one NSAID may experience adverse reactions from another NSAID. Patients at greatest risk for adverse events are those that are dehydrated, on concomitant diuretic therapy, or those with existing renal, cardiovascular, and/or hepatic dysfunction. Plasma levels of deracoxib may increase in a greater than dose-

proportional fashion above 8 mg/kg/day. Deracoxib tablets have been safely used during field studies in conjunction with other common medications, including heartworm preventatives, anthelmintics, anesthetics, pre-anesthetic medications, and antibiotics. If additional pain medication is needed after a daily dose of Deracoxib Chewable Tablets, a non-NSAID/non-corticosteroid class of analgesic may be necessary. It is not known whether dogs with a history of hypersensitivity to sulfonamide drugs will exhibit hypersensitivity to Deracoxib Chewable Tablets. The safe use of deracoxib tablets in dogs younger than 4 months of age, dogs used for breeding, or in pregnant or lactating dogs has not been evaluated.

NSAIDs may inhibit the prostaglandins which maintain normal homeostatic function. Such anti-prostaglandin effects may result in clinically significant disease in patients with underlying or pre-existing disease that has not been previously diagnosed. Appropriate monitoring procedures should be employed during all surgical procedures. The use of parenteral fluids during surgery should be considered to decrease potential renal complications when using NSAIDs perioperatively. Concurrent administration of potentially nephrotoxic drugs should be carefully approached.

The use of concomitantly protein-bound drugs with deracoxib tablets has not been studied in dogs. Commonly used protein-bound drugs include cardiac, anticonvulsant and behavioral medications. The influence of concomitant drugs that may inhibit metabolism of deracoxib tablets has not been evaluated. Drug compatibility should be monitored in patients requiring adjunctive therapy. Consider appropriate washout times when switching from one NSAID to another or when switching from corticosteroid use to NSAID use.

### **Animal Safety:**

In a 6-month study, dogs were dosed with deracoxib at 0, 2, 4, 6, 8 and 10 mg/kg with food once daily for 6 consecutive months. There were no abnormal feces, and no abnormal findings on clinical observations, food and water consumption, body weights, physical examinations, ophthalmoscopic evaluations, macroscopic pathological examinations, hematology, or buccal bleeding time. Urinalysis results showed hyposthenuria (specific gravity <1.005) and polyuria in one male and one female in the 6 mg/kg group after 6 months of treatment. After 6 months of treatment, the mean BUN values for dogs treated with 6, 8, or 10 mg/kg/day were 30.0, 35.3, and 48.2 mg/dL respectively. No effects were seen on any other clinical chemistry parameters, including other variables associated with renal physiology (serum creatinine, serum electrolytes, and urine sediment evaluation). Dose dependent focal renal tubular degeneration/regeneration was seen in some dogs treated at 6, 8, and 10 mg/kg/day. Focal renal papillary necrosis was seen in 3 dogs dosed at 10 mg/kg/day and in one dog dosed at 8 mg/kg/day. No renal lesions were seen at the label doses of 2 and 4 mg/kg/day. There was no evidence of gastrointestinal, hepatic, or hematopoietic pathology at any of the doses tested.

In a laboratory study, healthy young dogs were dosed with deracoxib tablets once daily, within 30 minutes of feeding, at doses of 0, 4, 6, 8, and 10 mg/kg body weight for 21 consecutive days. No adverse drug events were reported. There were no abnormal findings reported for clinical observations, food and water consumption, body weights, physical examinations, ophthalmic evaluations, organ weights, macroscopic pathologic evaluation, hematology, urinalyses, or buccal mucosal bleeding time. In the clinical

chemistry results there was a statistically significant ( $p < 0.0009$ ) dose-dependent trend toward increased levels of blood urea nitrogen (BUN). Mean BUN values remained within historical normal limits at the label dose. No effects on other clinical chemistry values associated with renal function were reported. There was no evidence of renal, gastrointestinal, hepatic or biliary lesions noted during gross necropsy. Renal histopathology revealed trace amounts of tubular degeneration/regeneration in all dose groups including placebo, but no clear dose relationship could be determined. There was no histopathologic evidence of gastrointestinal, hepatic or biliary lesions.

In another study, micronized deracoxib in gelatin capsules was administered once daily to healthy young dogs at doses of 10, 25, 50, and 100 mg/kg body weight for periods up to 14 consecutive days. Food was withheld prior to dosing. Non-linear elimination kinetics occurred at all doses. At doses of 25, 50, and 100 mg/kg, reduced body weight, vomiting, and melena occurred. Necropsy revealed gross gastrointestinal lesions in dogs from all dose groups. The frequency and severity of the lesions increased with escalating doses. At 10 mg/kg, moderate diffuse congestion of gut associated lymphoid tissues (GALT) and erosions/ulcers in the jejunum occurred. At 100 mg/kg, all dogs exhibited gastric ulcers and erosions/ulcerations of the small intestines. There were no hepatic or renal lesions reported at any dose in this study.

In a 13-week study, deracoxib in gelatin capsules was administered to healthy dogs at doses of 0, 2, 4, and 8 mg/kg/day. No test-article related changes were identified in clinical observations, physical exams, or any of the other parameters measured. One dog in the 8 mg/kg dose group died from bacterial septicemia secondary to a renal abscess. The relationship between deracoxib administration and the renal abscess is not clear.

### **Effectiveness:**

Deracoxib tablets were evaluated in masked, placebo-controlled multi-site field studies involving client-owned animals to determine effectiveness.

### **Osteoarthritis Pain and Inflammation Field Study:**

Two hundred and nine (209) client-owned dogs with clinical and radiographic signs of osteoarthritis of at least one appendicular joint were enrolled in this study. A total of 194 dogs were included in the safety evaluation and a total of 181 dogs were included in the effectiveness evaluation. The effectiveness of deracoxib tablets in the control of pain and inflammation associated with osteoarthritis was demonstrated in a placebo-controlled, masked study evaluating the anti-inflammatory and analgesic effects of deracoxib tablets. Tablets were administered by the owner at approximately 1-2 mg/kg/day for forty-three (43) consecutive days. In general, statistically significant ( $p \leq 0.05$ ) differences in favor of deracoxib were seen for force plate parameters (vertical impulse area, peak vertical force) and owner evaluations (quality of life, lameness and overall level of activity).

The results of this field study demonstrate that deracoxib tablets, when administered at 1-2 mg/kg/day for 43 days, are effective for the control of pain and inflammation associated with osteoarthritis.

### **Adverse Reactions:**

Deracoxib was well tolerated and the incidence of clinical adverse reactions was

comparable in deracoxib and placebo-treated animals. A total of 209 dogs of 41 breeds, 1-14 years old, weighing 17-177 lbs were included in the field safety analysis. The following table shows the number of dogs displaying each adverse reaction.

### **Abnormal Health Findings in the Osteoarthritis Field Study<sup>1</sup>**

<b>Clinical Observation</b>	<b>Deracoxib tablets N = 105</b>	<b>Placebo N = 104</b>
Vomiting	3	4
Diarrhea/soft stool	3	2
Weight loss	1	0
Abdominal pain (splinting)	0	1
Seizure	1	0
Lethargy	0	1
Pyoderma/Dermatitis	2	0
Unilateral conjunctivitis	1	0
Scleral injection	0	1
Hematuria/UTI	1	0
Splenomegaly*	1	0
Grade II murmur systolic	1	0

<sup>1</sup>Dogs may have experienced more than one adverse reaction during the study.

\* This dog was less active and eating less on enrollment, with elevated WBC, amylase, and AST and died 1 month after exiting the study. The dog was withdrawn from the study on Day 17 with anorexia, lethargy and a suspicion of diarrhea. Follow-up laboratory analyses revealed hypoalbuminemia, hyperphosphatemia, elevated AST and decreased BUN. Follow-up treatment included other anti- inflammatories and antibiotics.

Complete blood count, serum chemistry, and buccal bleeding time analysis were conducted at the beginning and end of the trial. Mean values of all CBC and chemistry results for both deracoxib and placebo-treated dogs were within normal limits. There was no statistically significant difference in the buccal bleeding time between deracoxib and placebo-treated dogs before or after the study, and all results remained within normal limits (less than 5 minutes). The results of this field study demonstrate that deracoxib is safe and effective for the control of pain and inflammation associated with osteoarthritis in dogs. During this trial, dogs were safely treated with a variety of commonly used medications, including antibiotics, anti-parasiticides, topical flea adulticides and thyroid supplements.

The results of this field study demonstrate that deracoxib tablets are well tolerated when administered at 1-2 mg/kg/day for up to 43 days for the control of pain and inflammation associated with osteoarthritis.

### **Postoperative Orthopedic Pain and Inflammation Field Study:**

In this study, 207 dogs admitted to veterinary hospitals for repair of a cranial cruciate injury were randomly administered deracoxib tablets or a placebo. Drug administration started the evening before surgery and continued once daily for 6 days postoperatively. Effectiveness was evaluated in 119 dogs and safety was evaluated in 207 dogs. Statistically significant differences in favor of deracoxib tablets were found for lameness

at walk and trot, and pain on palpation values at all post-surgical time points. The results of this field study demonstrate that deracoxib tablets, when administered daily for 7 days are effective for the control of postoperative pain and inflammation associated with orthopedic surgery.

### **Adverse Reactions:**

A total of 207 dogs of forty-three (43) different breeds, 1-15 years old, weighing 7-141 lbs were included in the field safety analysis. The following table shows the number of dogs displaying each adverse reaction.

### **Abnormal Health Findings in The Postoperative Orthopedic Pain Field Study<sup>1</sup>**

<b>Clinical Observation</b>	<b>Deracoxib tablets N = 105</b>	<b>Placebo N = 102</b>
Vomiting	11	6
Diarrhea	6	7
Hematochezia	4	0
Melena	0	1
Anorexia	0	4
Incision site lesion (drainage, oozing)	11	6
Non-incision skin lesions (moist dermatitis, pyoderma)	2	0
Otitis externa	2	0
Positive joint culture	1	0
Phlebitis	1	0
Hematuria	2	0
Conjunctivitis	1	2
Splenomegaly	1	0
Hepatomegaly	1	0
Death	0	1

<sup>1</sup>Dogs may have experienced more than one adverse reaction during the study.

This table does not include one dog that was dosed at 16.92 mg/kg/day for the study duration. Beginning on the last day of treatment, this dog experienced vomiting, diarrhea, increased water intake and decreased appetite. Hematology and clinical chemistry values were unremarkable. The dog recovered uneventfully within 3 days of cessation of dosing.

Incisional drainage was most prevalent in dogs enrolled at a single study site. There were no statistically significant changes in the mean values for hepatic or renal clinical pathology indices between deracoxib tablet and placebo treated dogs. Four deracoxib tablet-treated dogs and two placebo-treated dogs exhibited elevated bilirubin during the dosing phase. One deracoxib tablet-treated dog exhibited elevated ALT, BUN and total bilirubin and a single vomiting event. None of the changes in clinical pathology values were considered clinically significant.

The results of this clinical study demonstrate that deracoxib tablets, when administered daily for 7 days to control postoperative orthopedic pain and inflammation in dogs, are well tolerated.



## Postoperative Dental Pain and Inflammation Field Study:

In this study, 62 dogs admitted to veterinary hospitals for dental extractions were randomly administered deracoxib tablets or a placebo. Drug administration started approximately 1 hour before surgery and continued once daily for 2 days postoperatively. Effectiveness was evaluated in 57 dogs and safety was evaluated in 62 dogs. There was a statistically significant reduction ( $p=0.0338$ ) in the proportion of dogs that required rescue therapy to control post-surgical pain in the deracoxib treated group compared to the placebo control group. Pain assessors used a modification of the Glasgow Composite Pain Scale (mGCPS) to assess pain.<sup>7</sup> A dog was rescued if it scored  $\geq 4$  on the combined mGCPS variables of Posture/Activity, Demeanor, Response to Touch, and Vocalization, or if the investigator determined at any time that pain intervention was needed. The results of this field study demonstrate that deracoxib, when administered once daily for 3 days, is effective for the control of postoperative pain and inflammation associated with dental surgery.

### Adverse Reactions:

A total of 62 male and female dogs of various breeds, 1.5-16 years old, were included in the field safety analysis. The following table shows the number of dogs displaying each adverse reaction. Digestive tract disorders (diarrhea and vomiting) and systemic disorders (abnormal clinical chemistry results) were the most frequently reported findings. There were no distinct breed, age, or sex predilections for adverse reactions that were reported. No dogs were withdrawn from the study due to the occurrence of an adverse reaction.

### Abnormal Health Findings in the Dental Pain Field Study<sup>1</sup>

Clinical Observation	Deracoxib tablets N = 31	Placebo N = 31
Vomiting	4	1
Diarrhea/soft stool	3	1
Regurgitation	0	2
Increased AST <sup>2</sup>	3	0
Increased ALT <sup>2</sup>	1	0
Hematuria	1	0
Leukocytosis	1	1
Neutrophilia	1	1
Lameness	1	0
Facial swelling	0	1
Tachycardia	0	1

<sup>1</sup>Dogs may have experienced more than one adverse reaction during the study.

<sup>2</sup>Included animals with results over 2x the high normal.

### Post Approval Experience (Rev. 2010):

The following adverse events are based on post-approval adverse drug experience reporting. Not all adverse reactions are reported to FDA CVM. It is not always possible to

reliably estimate the adverse event frequency or establish a causal relationship to product exposure using this data. The following adverse events are grouped by body system and are presented in decreasing order of reporting frequency.

Gastrointestinal: vomiting, diarrhea, hypoalbuminemia, melena, hematochezia, elevated amylase/lipase, hematemesis, abdominal pain, peritonitis, decreased or increased total protein and globulin, gastrointestinal perforation, gastrointestinal ulceration, hypersalivation.

General: anorexia, depression/lethargy, weight loss, weakness, fever, dehydration

Hepatic: elevated liver enzymes, hyperbilirubinemia, icterus, ascites, decreased BUN

Hematologic: anemia, leukocytosis, leukocytopenia, thrombocytopenia

Neurologic: seizures, ataxia, recumbency, trembling, confusion, collapse, hind limb paresis, nystagmus, proprioceptive disorder, vestibular signs

Behavioral: nervousness, hyperactivity, aggression, apprehension

Urologic: elevated BUN/creatinine, polydipsia, polyuria, hyper-phosphatemia, hematuria, low urine specific gravity, urinary incontinence, renal failure, urinary tract infection

Dermatologic: pruritus, erythema, urticaria, moist dermatitis, facial/muzzle edema, dermal ulceration/necrosis

Respiratory: panting, dyspnea, epistaxis, coughing

Cardiovascular: tachycardia, heart murmur, bradycardia, arrest

Sensory: Vestibular signs, glazed eyes, uveitis.

Ophthalmic: blindness, mydriasis, conjunctivitis, keratoconjunctivitis sicca, uveitis.

In some cases, death has been reported as an outcome of the adverse events listed above.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet, contact Covetrus North America at (855) 724-3461. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

## **Chewable Tablets**

### **Information for Dog Owners:**

Deracoxib Chewable Tablets, like other drugs of its class, is not free from adverse reactions. Owners should be advised of the potential for adverse reactions and be informed of the clinical signs associated with drug intolerance. Adverse reactions may include vomiting, diarrhea, decreased appetite, dark or tarry stools, increased water consumption, increased urination, anemia, yellowing of gums, skin or white of the eye due to jaundice, lethargy, incoordination, seizure, or behavioral changes. **Serious adverse reactions associated with this drug class can occur without warning and in some cases result in death (see Warnings, Post-Approval Experience and Adverse Reactions). Owners should be advised to discontinue Deracoxib Chewable Tablets therapy and contact their veterinarian immediately if signs of intolerance are observed.** The vast majority of patients with drug related adverse

reactions have recovered when the signs are recognized, the drug is withdrawn, and veterinary care, if appropriate, is initiated. Owners should be advised of the importance of periodic follow up for all dogs during administration of any NSAID.

### **Storage Conditions:**

Deracoxib Chewable Tablets should be Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° and 30°C (between 59° and 86°F) [see USP Controlled Room Temperature]

**Keep this and all medications out of reach of children.**

**Use within 90 days of splitting.**

### **How Supplied:**

Deracoxib Chewable Tablets are available as 12 mg, 25 mg, 75 mg and 100 mg round, brownish, half-scored tablets in 30 and 90 count bottles.

<b>NDC Number</b>	<b>Tablet Size</b>	<b>Tablets/Bottle</b>
11695-6995-1	12 mg	30
11695-6995-2	12 mg	90
11695-6996-1	25 mg	30
11695-6996-2	25 mg	90
11695-6997-1	75 mg	30
11695-6997-2	75 mg	90
11695-6998-1	100 mg	30
11695-6998-2	100 mg	90

### **References:**

<sup>1</sup> Data on file under NADA 141-203

<sup>2</sup> Data on file under NADA 141-203

<sup>3</sup> Smith, et al.: "Pharmacological Analysis of Cyclo-oxygenase-1 in Inflammation," Proc. Natl. Acad. Sci. USA (October 1998) 95: 13313-13318, Pharmacology.

<sup>4</sup> Zhang, et al.: "Inhibition of Cyclo-oxygenase-2 Rapidly Reverses Inflammatory Hyperalgesia and Prostaglandin E2 Production," *JPET*, (1997) 283: 1069-1075.

<sup>5</sup> Verburg, KM et al. "Cox-2 Specific Inhibitors: Definition of a New Therapeutic Concept." Amer J of Therapeutics 8, 49-64, 2001.

<sup>6</sup> Data on file under NADA 141-203

<sup>7</sup> Holton, L., Reid, J., Scott, E.M., Pawson, P. and Nolan, A. (2001). Development of a behaviour-based scale to measure acute pain in dogs. Veterinary Record, 148, 525-53.

Approved by FDA under ANADA # 200-704

**Questions?** (855) 724-3461

Distributed by: Covetrus North America,  
400 Metro Place North, Dublin, OH 43017

**covetrus.com**

Made in India

## **INFORMATION FOR DOG OWNERS:**

### **Deracoxib**

#### **Chewable Tablets**

#### **For Oral Use in Dogs Only**

#### **Do Not Use in Cats**

#### **Information for Dog Owners**

Deracoxib Chewable Tablets are for the control of pain and inflammation due to osteoarthritis or following orthopedic and dental surgery.

This summary contains important information about Deracoxib Chewable Tablets. You should read this information before starting your dog on Deracoxib Chewable Tablets. This sheet is provided only as a summary and does not take the place of instructions from your veterinarian. Talk to your veterinarian if you do not understand any of this information or you want to know more about Deracoxib Chewable Tablets.

#### **What is Deracoxib Chewable Tablets?**

Deracoxib Chewable Tablets are a prescription non-steroidal anti-inflammatory drug (NSAID) of the coxib class. They are indicated for the control of postoperative pain and inflammation associated with orthopedic (bone) and dental surgery in dogs and for the control of pain and inflammation (soreness) associated with osteoarthritis in dogs. The tablets are flavored to make administration more convenient.

#### **What kind of results can I expect when my dog takes Deracoxib Chewable Tablets for postoperative orthopedic and dental pain and inflammation?**

Deracoxib Chewable Tablets allow your dog to recover more comfortably by controlling pain and inflammation that follow orthopedic and dental surgery.

- The postoperative orthopedic pain dose is a higher dose; administration at this dose **should not exceed 7 days total, including the days the patient is administered Deracoxib Chewable Tablets while in the hospital.**
- Control of pain and inflammation may vary from dog to dog.
- If Deracoxib Chewable Tablets are not given according to your veterinarian's directions, your dog's pain may return.
- Consult your veterinarian if your dog appears to be uncomfortable.

#### **What kind of results can I expect when my dog takes Deracoxib Chewable Tablets for pain and inflammation due to osteoarthritis?**

Osteoarthritis is a painful condition caused by damage to cartilage and other parts of the joint that may result in the following changes or signs in your dog:

- Limping or lameness
- Decreased activity or exercise (reluctance to stand, climb stairs, jump or run, or difficulty in performing these activities)
- Stiffness or decreased movement of joints

While Deracoxib Chewable Tablets is not a cure for osteoarthritis, it can control the pain and inflammation of osteoarthritis and improve your dog's mobility. Response may vary from dog to dog but can be quite dramatic. **Deracoxib Chewable Tablets may need to be given on a periodic basis for the animal's lifetime. Use the lowest dose to provide adequate relief. Always consult with your veterinarian before altering the dose.**

### **What dogs should not take Deracoxib Chewable Tablets?**

Your dog should not be given Deracoxib Chewable Tablets if s/he:

- Has had an allergic reaction to deracoxib, the active ingredient in Deracoxib Chewable Tablets
- Has had an allergic reaction (such as hives, facial swelling, or red or itchy skin) to aspirin or other NSAIDs
- Is presently taking aspirin, other NSAIDs, or corticosteroids (unless directed by your veterinarian)
- Has bloody stool or vomit
- Has a pre-existing kidney or liver condition
- Has any condition predisposing to dehydration
- Is anorexic (loss of appetite)

### **Deracoxib Chewable Tablets should only be given to dogs. Do not use in cats.**

People should not take Deracoxib Chewable Tablets. Keep Deracoxib Chewable Tablets and all medication out of reach of children. Call your physician immediately if you accidentally take Deracoxib Chewable Tablets.

### **What to discuss with your veterinarian before giving Deracoxib Chewable Tablets?**

Tell your veterinarian about:

- Any side effects your dog has experienced from Deracoxib Chewable Tablets or other NSAIDs
- Any digestive upset (vomiting or diarrhea) your dog has had
- Any kidney disease your dog has had
- Any other medical problems or allergies that your dog has now or has had in the past
- All medications that you are giving your dog or plan to give your dog, including those you can get without prescription and any dietary supplements
- If you plan to breed your dog, or if your dog is pregnant or nursing

Talk to your veterinarian about:

- The orthopedic or dental surgery your dog will undergo
- What tests might be done before surgery is performed or Deracoxib Chewable Tablets are prescribed
- The signs of pain or inflammation that may occur after surgery
- Normal events that can be expected after your dog undergoes surgery
- The proper amount of exercise after surgery to aid recovery
- The signs of osteoarthritis you have observed (for example, limping or stiffness)
- The importance of weight control, physical therapy and exercise in the management of osteoarthritis
- How often your dog may need to be examined by your veterinarian

- The risks and benefits of using Deracoxib Chewable Tablets

## **How to give Deracoxib Chewable Tablets to your dog.**

Deracoxib Chewable Tablets should be given according to your veterinarian's instructions. Your veterinarian will tell you what amount of Deracoxib Chewable Tablets is right for your dog and for how long they should be given. Do not change the way you give Deracoxib Chewable Tablets to your dog without first speaking with your veterinarian. Deracoxib Chewable Tablets should be given by mouth and may be given with or without food, although with food is preferable.

## **What are the possible side effects that may occur in my dog during therapy with Deracoxib Chewable Tablets?**

Deracoxib Chewable Tablets may cause some side effects in individual dogs. Serious side effects associated with this drug can occur with or without warning and, in some cases, result in death. The most common side effects associated with Deracoxib Chewable Tablets therapy involve the digestive tract (vomiting, decreased appetite and diarrhea). Liver and kidney problems have also been reported. It is important to stop the medication and contact your veterinarian immediately if you think your dog may have a medical problem or side effect while on Deracoxib Chewable Tablets. If you have additional questions about possible side effects, talk with your veterinarian or call Covetrus North America at (855) 724-3461.

Look for the following side effects that may indicate that your dog is having a problem with Deracoxib Chewable Tablets or may have another medical problem:

- Vomiting
- Decrease in appetite
- Change in behavior, such as depression, restlessness, aggression or apprehension
- Change in bowel movements such as diarrhea or change in stool color (black, tarry or bloody stool)
- Change in drinking or urination
- Yellowing of gums, skin or whites of the eyes (jaundice)

## **Can Deracoxib Chewable Tablets be given with other medications?**

Deracoxib Chewable Tablets should not be given with other non-steroidal anti-inflammatory drugs (NSAIDs) or corticosteroids (for example, aspirin, carprofen, etodolac, prednisone).

Tell your veterinarian about all medications that you have given your dog in the past, and any medications that you are planning to give with Deracoxib Chewable Tablets. This should include any medications that you can get without a prescription and any dietary supplements. Your veterinarian may want to evaluate the potential for any drug interactions and to assure drug compatibility.

## **What can I do in case my dog eats more than the prescribed amount of Deracoxib Chewable Tablets?**

Contact your veterinarian immediately if your dog eats more than the prescribed amount of Deracoxib Chewable Tablets.

## **What else should I know about Deracoxib Chewable Tablets?**

This sheet provides a summary of information about Deracoxib Chewable Tablets. If you

have any questions or concerns about Deracoxib Chewable Tablets, postoperative orthopedic and dental pain and inflammation, or osteoarthritis, talk to your veterinarian.

As with all prescribed medications, Deracoxib Chewable Tablets should only be given to the dog for which they are prescribed. They should be given to your dog only for the condition for which they were prescribed, at the prescribed dose, as directed by your veterinarian. It is important to periodically discuss your dog's response to Deracoxib Chewable Tablets at regular checkups. Your veterinarian will best determine if your dog is responding as expected and if your dog should continue receiving Deracoxib Chewable Tablets.

Keep Deracoxib in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

**Use within 90 days of splitting.**

Approved by FDA under ANADA # 200-704

**Questions?** (855) 724-3461

Distributed by: Covetrus North America,  
400 Metro Place North, Dublin, OH 43017

**covetrus.com**

Made in India

**PRINCIPAL DISPLAY PANEL - 12mg Tablet x 30 Count Bottle**

NDC: 11695-6995-1

**Deracoxib**

**Chewable Tablets**

**12 mg**

**For Oral Use in Dogs Only**

**Do Not Use in Cats**

**A coxib-class NSAID**

Caution: Federal law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-704

30 Tablets



## PRINCIPAL DISPLAY PANEL - 12mg Tablet x 90 Count Bottle

NDC: 11695-6995-2

**Deracoxib**

**Chewable Tablets**

**12 mg**

**For Oral Use in Dogs Only**

**Do Not Use in Cats**

**A coxib-class NSAID**

Caution: Federal law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-704

90 Tablets



## PRINCIPAL DISPLAY PANEL - 25mg Tablet x 30 Count Bottle

NDC: 11695-6996-1

**Deracoxib**

**Chewable Tablets**



**25 mg**

**For Oral Use in Dogs Only**


**Do Not Use in Cats**

**A coxib-class NSAID**

Caution: Federal law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-704

30 Tablets



**PRINCIPAL DISPLAY PANEL - 25mg Tablet x 90 Count Bottle**

NDC: 11695-6996-2

**Deracoxib**

**Chewable Tablets**

**25 mg**

**For Oral Use in Dogs Only**

**Do Not Use in Cats**

**A coxib-class NSAID**

Caution: Federal law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-704

90 Tablets

**covetrus**

NDC: 11695-6996-2

# Deracoxib

**Chewable Tablets**

**For Oral Use in Dogs Only**  
**Do Not Use in Cats**  
**A coxib-class NSAID**

**25 mg**

8 40066 51220 4

11 001 482

**Reorder #074738**  
AH-074738-L-01 REV: 0621  
Questions? (855) 724-3461  
Distributed by: Covetrus North America, 400 Metro Place North, Dublin, OH 43017, covetrus.com

**Made in India**  
Neutral Code No. DRUGS/TS/32/2008.

**90 Tablets**

**Desage & Administration.**  
For complete product information, see package insert.

**Storage Conditions:**  
Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° and 30°C (between 59° and 86°F) [see USP Controlled Room Temperature]

**Use within 90 days of splitting.**

**WARNINGS:**  
**Keep this and all medications out of the reach of children.**  
Keep Deracoxib in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

Caution: Federal law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-704

## PRINCIPAL DISPLAY PANEL - 75mg Tablet x 30 Count Bottle

NDC: 11695-6997-1

**Deracoxib**

**Chewable Tablets**

**75 mg**

**For Oral Use in Dogs Only**

**Do Not Use in Cats**

**A coxib-class NSAID**

Caution: Federal law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-704

30 Tablets

**covetrus**

NDC: 11695-6997-1

# Deracoxib

**Chewable Tablets**

**For Oral Use in Dogs Only**  
**Do Not Use in Cats**  
**A coxib-class NSAID**

**75 mg**

8 40066 51221 1

11 001 481

**Reorder #074739**  
AH-074739-L-01 REV: 0621  
Questions? (855) 724-3461  
Distributed by: Covetrus North America, 400 Metro Place North, Dublin, OH 43017, covetrus.com

**Made in India**  
Neutral Code No. DRUGS/TS/32/2008.

**30 Tablets**

**Desage & Administration.**  
For complete product information, see package insert.

**Storage Conditions:**  
Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° and 30°C (between 59° and 86°F) [see USP Controlled Room Temperature]

**Use within 90 days of splitting.**

**WARNINGS:**  
**Keep this and all medications out of the reach of children.**  
Keep Deracoxib in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

Caution: Federal law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-704

## PRINCIPAL DISPLAY PANEL - 75mg Tablet x 90 Count Bottle

NDC: 11695-6997-2

**Deracoxib**

**Chewable Tablets**

**75 mg**

**For Oral Use in Dogs Only**

**Do Not Use in Cats**

**A coxib-class NSAID**

Caution: Federal law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-704

90 Tablets

**covetrus**

NDC: 11695-6997-2

# Deracoxib

## Chewable Tablets

**For Oral Use in Dogs Only**  
**Do Not Use in Cats**  
**A coxib-class NSAID**

**75 mg**

Caution: Federal law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-704

**90 Tablets**

**Dosage & Administration.**  
For complete product information, see package insert.

**Storage Conditions:**  
Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° and 30°C (between 59° and 86°F) [see USP Controlled Room Temperature]

**Use within 90 days of splitting.**

**WARNINGS:**  
**Keep this and all medications out of the reach of children.**

Keep Deracoxib in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

Reorder #074740  
AH-074740-L-01 REV: 0621  
Questions? (855) 724-3461  
Distributed by: Covetrus North America, 400 Metro Place North, Dublin, OH 43017, covetrus.com  
Made in India  
Neutral Code No. DRUGS/TS/32/2008.

8 40066 51222 8 11001480

**PRINCIPAL DISPLAY PANEL - 100 mg Tablet x 30 Count Bottle**

NDC: 11695-6998-1

**Deracoxib**

**Chewable Tablets**

**100 mg**

**For Oral Use in Dogs Only**

**Do Not Use in Cats**

**A coxib-class NSAID**

Caution: Federal law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-704

30 Tablets

**Dosage & Administration.**  
For complete product information, see package outsert.

**Storage Conditions:**  
Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° and 30°C (between 59° and 86°F) [see USP Controlled Room Temperature]

**Use within 90 days of splitting.**

**WARNINGS:**  
**Keep this and all medications out of the reach of children.**  
Keep Deracoxib in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

**covetrus**  
NDC: 11695-6998-1  
**Deracoxib**  
**Chewable Tablets**  
**For Oral Use in Dogs Only**  
**Do Not Use in Cats**  
**A coxib-class NSAID**  
**100 mg**  
8 40066 51223 5  
11001478  
Reorder #074741  
AH-074741-L-01 REV: 0621  
Questions? (855) 724-3461  
Distributed by: Covetrus North America, 400 Metro Place North, Dublin, OH 43017, covetrus.com  
Made in India  
Neutral Code No. DRUGS/TS/3/2/2008.

30 Tablets

**PRINCIPAL DISPLAY PANEL - 100 mg Tablet x 90 Count Bottle**

NDC: 11695-6998-2

**Deracoxib**

**Chewable Tablets**

**100 mg**

**For Oral Use in Dogs Only**

**Do Not Use in Cats**

**A coxib-class NSAID**

Caution: Federal law (U.S.) restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-704

90 Tablets

**Dosage & Administration.**  
For complete product information, see package outsert.

**Storage Conditions:**  
Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° and 30°C (between 59° and 86°F) [see USP Controlled Room Temperature]

**Use within 90 days of splitting.**

**WARNINGS:**  
**Keep this and all medications out of the reach of children.**  
Keep Deracoxib in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

**covetrus**  
NDC: 11695-6998-2  
**Deracoxib**  
**Chewable Tablets**  
**For Oral Use in Dogs Only**  
**Do Not Use in Cats**  
**A coxib-class NSAID**  
**100 mg**  
8 40066 51224 2  
11001479  
Reorder #074742  
AH-074742-L-01 REV: 0621  
Questions? (855) 724-3461  
Distributed by: Covetrus North America, 400 Metro Place North, Dublin, OH 43017, covetrus.com  
Made in India  
Neutral Code No. DRUGS/TS/3/2/2008.

90 Tablets

**DERACOXIB CHEWABLE**

deracoxib tablet

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:11695-6995
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>DERACOXIB</b> (UNII: VX29JB5XWW) (DERACOXIB - UNII:VX29JB5XWW)	DERACOXIB	12 mg

### Product Characteristics

<b>Color</b>	brown	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>	MEAT (beef)	<b>Imprint Code</b>	12;F;1
<b>Contains</b>			

### Packaging

#	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11695-6995-1	30 in 1 BOTTLE		
2	NDC:11695-6995-2	90 in 1 BOTTLE		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANADA	ANADA200704	12/15/2021	

## DERACOXIB CHEWABLE

deracoxib tablet

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:11695-6996
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>DERACOXIB</b> (UNII: VX29JB5XWW) (DERACOXIB - UNII:VX29JB5XWW)	DERACOXIB	25 mg

### Product Characteristics

<b>Color</b>	brown	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	11mm
<b>Flavor</b>	MEAT (beef)	<b>Imprint Code</b>	25;F;1
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11695-6996-1	30 in 1 BOTTLE		
2	NDC:11695-6996-2	90 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200704	12/15/2021	

## DERACOXIB CHEWABLE

deracoxib tablet

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:11695-6997
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DERACOXIB (UNII: VX29JB5XWW) (DERACOXIB - UNII: VX29JB5XWW)	DERACOXIB	75 mg

### Product Characteristics

<b>Color</b>	brown	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	16mm
<b>Flavor</b>	MEAT (beef)	<b>Imprint Code</b>	75;F;1
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11695-6997-1	30 in 1 BOTTLE		
2	NDC:11695-6997-2	90 in 1 BOTTLE		

### Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
ANADA	ANADA200704	12/15/2021	

## DERACOXIB CHEWABLE

deracoxib tablet

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:11695-6998
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DERACOXIB (UNII: VX29JB5XWW) (DERACOXIB - UNII: VX29JB5XWW)	DERACOXIB	100 mg

### Product Characteristics

<b>Color</b>	brown	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	19mm
<b>Flavor</b>	MEAT (beef)	<b>Imprint Code</b>	100;F;1
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11695-6998-1	30 in 1 BOTTLE		
2	NDC:11695-6998-2	90 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200704	12/15/2021	

**Labeler** - BUTLER ANIMAL HEALTH SUPPLY, LLC DBA COVETRUS NORTH AMERICA (603750329)

**Registrant** - Felix Pharmaceuticals Private Limited (985612369)

Revised: 11/2021

BUTLER ANIMAL HEALTH SUPPLY, LLC DBA COVETRUS NORTH AMERICA