

**NOREL AD- acetaminophen, chlorpheniramine maleate, and phenylephrine hcl tablet, multilayer**  
**U.S. PHARMACEUTICAL CORPORATION**

-----  
**Norel® AD**

***Drug Facts***

***Active Ingredients (per tablet)***

Acetaminophen 325 mg

Chlorpheniramine Maleate 4 mg

Phenylephrine HCl 10 mg

***Purpose***

Pain reliever

Antihistamine

Nasal decongestant

***Uses***

Temporarily relieves these symptoms due to the common cold or flu

- nasal congestion
- runny nose
- minor aches and pains
- headache
- sore throat
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- sinus congestion and pressure
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 6 tablets in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reaction.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, speak with a healthcare provider.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking any other nasal decongestant or stimulant
- taking sedatives or tranquilizers

### **When using this product**

- do not exceed recommended dosage
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

### **Stop use and ask a doctor if**

- pain or nasal congestion gets worse or lasts more than 7 days

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- you get nervous, dizzy, or sleepless
- new symptoms occur

### **If pregnant or breast feeding**

- ask a health professional before use

### **Keep out of the reach of children.**

Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- do not take more than directed (see overdose warning)
- adults and children 12 years of age and over: take 1 tablet every 4 hours, while symptoms persist
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years of age: ask a doctor

### **Other Information**

store at 20°-25°C (68°-77°F)

**Overdose Warning:** Keep this and all medication out of the reach of children. In case of accidental overdose, seek medical help or contact a Poison Control Center immediately.

### **Inactive Ingredients**

Colloidal Silicon Dioxide, Crospovidone Polyplasdone, Magnesium Stearate (vegetable base), Microcrystalline Cellulose, Povidone K-30, Pregelatinized Starch, Stearic Acid, D&C Yellow #10, FD&C Yellow #6.

### **Questions or Comments**

Please visit [www.uspco.com](http://www.uspco.com) or contact us at US Pharmaceutical Corporation, P.O. Box 360465, Decatur, GA 30036

### **Principal Display Panel**

**Figure 1:** Norel<sup>®</sup> AD container label

Lot No.:  
Exp. Date:



NDC 52747-475-70

**Norel<sup>®</sup> AD**

**Acetaminophen,**

**Chlorpheniramine Maleate,  
Phenylephrine HCl**

Acetaminophen 325 mg ..... Pain reliever  
Chlorpheniramine Maleate 4 mg ... Antihistamine  
Phenylephrine HCl 10 mg .... Nasal Decongestant  
**20 Tablets**

Do not use if foil seal under the cap is broken or missing. You should contact your healthcare provider for medical advice about adverse events. To report a serious adverse event, contact US Pharmaceutical Corporation, P.O. Box 360465, Decatur, GA 30036. Marketed by US Pharmaceutical Corporation, P.O. Box 360465, Decatur, GA 30036. The white and yellow triangular shape and name of NOREL<sup>®</sup> AD tablets are registered trademarks of US Pharmaceutical Corporation, Decatur, GA 30036.

Lift Here  
for  
Drug Facts

400776 Rev. 08/2022

## NOREL AD

acetaminophen, chlorpheniramine maleate, and phenylephrine hcl tablet, multilayer

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:52747-475
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

### Product Characteristics

<b>Color</b>	yellow, white	<b>Score</b>	no score
<b>Shape</b>	TRIANGLE (Arc)	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	0425;US

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52747-475-70	20 in 1 BOTTLE; Type 0: Not a Combination Product	04/27/2012	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/27/2012	

**Labeler** - U.S. PHARMACEUTICAL CORPORATION (079467662)

Revised: 10/2023

U.S. PHARMACEUTICAL CORPORATION