MINOXIDIL SOLUTION- minoxidil solution solution Nutrastim Hair LLC

NUTRASTIM HAIR REGROWTH TREATMENT FOR WOMEN

To regrow hair on the scalp.

- For external use only.
- For use by women only.
- Flammable. Keep away from fire or flame.
- Your degree of hair loss is different than that shown on the side of this carton, because this product may not work for you.
- You have no family history of hair loss.
- Your hair loss is sudden and/or patchy.
- Your hair loss is associated with child birth.
- You do not know the reason for your hair loss.
- You are under 18 years of age.
- Do not use on babies and children.
- Your scalp is red, inflamed, infected, irritated, or painful.
- You use other medicines on the scalp.

Ask a doctor before use if you have heart disease

- Do not apply on other parts of the body.
- Avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
- Some people have experienced changes in hair color and/or texture.
- It takes time to regrow hair. Results may occur at 2 months with twice a day usage. For some women, you may need to use this product for at least 4 months before you see results.
- The amount of hair regrowth is different with each person. This product will not work for all women.
- Chest pain, rapid heartbeat, faintness, or dizziness occurs.
- Sudden, unexplained weight gain occurs.
- Your hands or feet swell.
- Scalp irritation or redness occurs.
- Unwanted facial hair growth occurs.
- You do not see hair regrowth in 4 months.

May be harmful if used when pregnant or breast-feeding.

□Keep out of reach of children. □If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

Directions

- Apply 1 ml, 2 times daily directly with the dropper onto the scalp in the hair loss area.
- Using more or more often will not improve results.
- Continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again.

Other information

• Before use see hair loss pictures on side of this box.

- Keep the box, it contains important information.
- In clinical studies of mostly white women aged 18-45 years with mild to moderate degrees of hair loss, the following response to minoxidil topical solution was 2% was reported: 19% of women reported moderate hair regrowth after using minoxidil topical solution 2% for 8 months (19% had moderate regrowth: 40% had minimal regrowth). This compares with 7% of women reporting moderate hair regrowth after using the placebo, the liquid without minoxidil in it, for 8 months (7% had moderate regrowth; 33% had minimal regrowth).
- Store at 20° to 25° C (68° to 77° F).

Keep tightly closed.

Alcohol, propylene glycol, purified water

Call us at 1-844-892-8578

Visit nutrastimhair.com

for more information about hair thinning, visit

nutrastimhair.com/women-education/



MINOXIDIL SOLUTION

minoxidil solution solution

Prod	luct	Info	rma	tion
FIUU	luct	HILL	I IIIa	uvu

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71517-001

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength	
MINO XIDIL (UNII: 5965120 SH1) (MINO XIDIL - UNII:5965120 SH1)	MINOXIDIL	20 g in 1 mL	

Inactive Ingredients

mactive ingredients			
Ingredient Name	Strength		
ALCOHOL (UNII: 3K9958V90M)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059 OF0 KOOR)			

Packaging

# Item Code	Package Description	Marketing Start Date	Marketing End Date
NDC:71517-001- 02	2 in 1 CARTON	06/29/2017	
NDC:71517-001- 01	60 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		

Marketing Information

muinemg mormuon			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075598	06/29/2017	

Labeler - Nutrastim Hair LLC (080274282)

Revised: 11/2019 Nutrastim Hair LLC