

CARE ONE ALL DAY PAIN RELIEF- naproxen sodium tablet, film coated
American Sales Company

American Sales Company All Day Pain Relief Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - you have difficulty swallowing
 - it feels like the pill is stuck in your throat
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older:

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years:

- ask a doctor

Other information

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the Active Ingredient in Aleve®

ALL DAY PAIN RELIEF

Naproxen Sodium Tablets, 220 mg


ACTUAL SIZE

PAIN RELIEVER/FEVER REDUCER (NSAID)

Strength to Last 12 Hours

50 CAPLETS**

(**Capsule-Shaped Tablets)



DO NOT USE IF PRINTED FOIL UNDER
CAP IS BROKEN OR MISSING

ACTUAL SIZE



Compare to the
Active Ingredient
in Aleve®***

NDC 41520-368-71

CAREONE®

ALL DAY PAIN RELIEF

Naproxen Sodium Tablets, 220 mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

Strength to Last 12 Hours



ACTUAL
SIZE

50
CAPLETS**

**Capsule-Shaped Tablets






EXP. DATE :

EXP.

LOT NO.



Drug Facts

Active ingredient (in each caplet) Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*
Purposes Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ minor pain of arthritis ■ muscular aches ■ backache
■ menstrual cramps ■ headache ■ toothache
■ the common cold ■ temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you: ■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have asthma

Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition
■ taking any other drug

Drug Facts (continued)

When using this product

■ take with food or milk if stomach upset occurs
■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint

■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ you have difficulty swallowing
■ it feels like the pill is stuck in your throat
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

■ do not take more than directed
■ the smallest effective dose should be used
■ drink a full glass of water with each dose

Adults and children 12 years and older:

■ take 1 caplet every 8 to 12 hours while symptoms last
■ for the first dose you may take 2 caplets within the first hour
■ do not exceed 2 caplets in any 8- to 12-hour period
■ do not exceed 3 caplets in a 24-hour period

Children under 12 years: ■ ask a doctor

Other information

■ each caplet contains: sodium 21 mg
■ store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Drug Facts (continued)

Inactive ingredients: FD&C Blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-800-719-9280

KEEP CARTON FOR REFERENCE

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve®.

GLUTEN FREE

**DISTRIBUTED BY FOODHOLD U.S.A., LLC
ANDOVER, MD 20785 • 1-877-846-9949
©2013 S&S Brands, LLC**

Quality guaranteed or your money back.

CARE ONE ALL DAY PAIN RELIEF

naproxen sodium tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-368
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONES (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BLUE (Light Blue)	Score	no score
Shape	CAPSULE (oval)	Size	12mm
Flavor		Imprint Code	L368
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-368-71	1 in 1 CARTON		
1		50 in 1 BOTTLE		
2	NDC:41520-368-78	1 in 1 CARTON		
2		100 in 1 BOTTLE		
3	NDC:41520-368-72	1 in 1 CARTON		
3		60 in 1 BOTTLE		
4	NDC:41520-368-82	2 in 1 CARTON		
4		100 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	04/05/2004	

Labeler - American Sales Company (809183973)

Revised: 12/2013

American Sales Company