

**ANTACID MINT- aluminum hydroxide, magnesium hydroxide,
dimethicone suspension**
Chain Drug Marketing Association

qc ant mnt

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 200 mg (equivalent to dried gel, USP)

Magnesium hydroxide 200 mg

Simethicone 20mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before each use

- adults and children 12 years and older: take 2 to 4 teaspoonfuls, as needed, up to four times a day, or as directed by a doctor
- do not take more than 16 teaspoonfuls in 24 hours unless directed by a doctor or use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

Other information

- **each 5 mL teaspoonful contains:** magnesium 85 mg, sodium 3 mg
- store at room temperature
- protect from freezing
- keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

package Label

NDC 63868-694-57



*Compare to the
Active Ingredients
in MAALOX® Regular
Strength

Regular Strength Antacid Suspension

Antacid & Antigas

Alumina, Magnesia, and
Simethicone
Oral Suspension USP

Fast Relief of:
Acid Indigestion
Heartburn
Sour Stomach
Pressure & Bloating



Cool Mint
Flavored Liquid

12 FL OZ (355 mL)

REV GC-0119
231-11212-3

Drug Facts

TAMPER EVIDENT: DO NOT USE IF BREAKAWAY
BAND ON BOTTLE CAP IS MISSING OR BROKEN.

Active ingredients

(in each 5 mL teaspoonful)

Aluminum hydroxide 200 mg (equivalent to dried gel USP)	Antacid
Magnesium hydroxide 200 mg	Antacid
Simethicone 20 mg	Antigas

Purposes

Uses relieves • heartburn • sour stomach • acid indigestion
• the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescriptions drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

if pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Directions

- shake well before each use
- adults and children 12 years and older: take 2 to 4 teaspoonfuls four times a day or as directed by a doctor
- do not exceed 16 teaspoonfuls in a 24 hour period or use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

Other information

- each 5 mL teaspoonful contains: magnesium 85 mg, sodium 3 mg
- store at room temperature
- protect from freezing
- keep tightly closed

Inactive ingredients benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

*This product is not manufactured or distributed by the owner of the registered trademark MAALOX®

REV GC-0119
931-11212-3



Distributed by C.D.M.A., Inc.®
43157 W. Nine Mile
Novi, MI 48376-0995
www.qualitychoice.com
Questions: 248-449-9300



ANTACID MINT

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-694
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE -	ALUMINUM	200 mg

UNII:5QB0T2IUN0)	HYDROXIDE	in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	200 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	20 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-694-57	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	03/01/2014	

Labeler - Chain Drug Marketing Association (011920774)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(63868-694)

