


**HYDROXYZINE HYDROCHLORIDE- hydroxyzine hydrochloride tablet, film coated**  
**Dispensing Solutions, Inc.**

-----  
**HYDROXYZINE HYDROCHLORIDE TABLETS, USP**

**Rx only**

**DESCRIPTION**

Hydroxyzine hydrochloride has the chemical name of 2-[2-[4-(*p*-Chloro- $\alpha$ -phenylbenzyl)-1-piperazinyl]ethoxy]ethanol dihydrochloride.



$C_{21}H_{27}ClN_2O_2 \cdot 2HCl$

M.W. = 447.83

Hydroxyzine hydrochloride occurs as a white, odorless powder which is very soluble in water.

Each tablet for oral administration contains 10 mg, 25 mg, or 50 mg hydroxyzine HCl. Inactive ingredients include colloidal silicon dioxide, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, sodium starch glycolate, and titanium dioxide.

The 10 mg tablet also contains FD&C Red #40 Aluminum Lake and FD&C Yellow #6 Aluminum Lake. The 25 mg tablet also contains D&C Yellow #10 Aluminum Lake and FD&C Blue #1 Aluminum Lake. The 50 mg tablet also contains FD&C Red #40 Aluminum Lake and FD&C Yellow #5 Aluminum Lake.

**CLINICAL PHARMACOLOGY**

Hydroxyzine hydrochloride is unrelated chemically to the phenothiazines, reserpine, meprobamate or the benzodiazepines. Hydroxyzine is not a cortical depressant, but its action may be due to a suppression of activity in certain key regions of the subcortical area of the central nervous system.

Primary skeletal muscle relaxation has been demonstrated experimentally. Bronchodilator activity, and antihistaminic and analgesic effects have been demonstrated experimentally and confirmed clinically. An antiemetic effect, both by the apomorphine test and the veriloid test, has been demonstrated.

Pharmacological and clinical studies indicate that hydroxyzine in therapeutic dosage does not increase gastric secretion or acidity and in most cases has mild antisecretory activity.

Hydroxyzine is rapidly absorbed from the gastrointestinal tract and hydroxyzine's clinical effects are usually noted within 15 to 30 minutes after oral administration.

## INDICATIONS AND USAGE

For symptomatic relief of anxiety and tension associated with psychoneurosis and as an adjunct in organic disease states in which anxiety is manifested.

Useful in the management of pruritus due to allergic conditions such as chronic urticaria and atopic and contact dermatoses and in histamine-mediated pruritus.

As a sedative when used as a premedication and following general anesthesia, **hydroxyzine may potentiate meperidine and barbiturates**, so their use in pre-anesthetic adjunctive therapy should be modified on an individual basis. Atropine and other belladonna alkaloids are not affected by the drug. Hydroxyzine is not known to interfere with the action of digitalis in any way and it may be used concurrently with this agent.

The effectiveness of hydroxyzine as an antianxiety agent for long term use, that is more than 4 months, has not been assessed by systematic clinical studies. The physician should reassess periodically the usefulness of the drug for the individual patient.

## CONTRAINDICATIONS

Hydroxyzine, when administered to the pregnant mouse, rat, and rabbit induced fetal abnormalities in the rat and mouse at doses substantially above the human therapeutic range. Clinical data in human beings are inadequate to establish safety in early pregnancy. Until such data are available, hydroxyzine is contraindicated in early pregnancy.

Hydroxyzine is contraindicated for patients who have shown a previous hypersensitivity to any component of this medication.

## WARNINGS

### **Nursing Mothers:**

It is not known whether this drug is excreted in human milk. Since many drugs are so excreted, hydroxyzine should not be given to nursing mothers.

## PRECAUTIONS

### **General:**

THE POTENTIATING ACTION OF HYDROXYZINE MUST BE CONSIDERED WHEN THE DRUG IS USED IN CONJUNCTION WITH CENTRAL NERVOUS SYSTEM DEPRESSANTS SUCH AS NARCOTICS, NON-NARCOTIC ANALGESICS AND BARBITURATES. Therefore, when central nervous system depressants are administered concomitantly with hydroxyzine their dosage should be reduced.

Since drowsiness may occur with use of this drug, patients should be warned of this possibility and cautioned against driving a car or operating dangerous machinery while taking hydroxyzine. Patients should also be advised against the simultaneous use of other CNS depressant drugs, and cautioned that the effects of alcohol may be increased.

Hydroxyzine Hydrochloride Tablets 50 mg contain FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

### **Geriatric Use:**

A determination has not been made whether controlled clinical studies of hydroxyzine included sufficient numbers of subjects aged 65 and over to define a difference in response from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

The extent of renal excretion of hydroxyzine has not been determined. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selections.

Sedating drugs may cause confusion and over sedation in the elderly; elderly patients generally should be started on low doses of hydroxyzine and observed closely.

## **ADVERSE REACTIONS**

Side effects reported with the administration of hydroxyzine hydrochloride are usually mild and transitory in nature.

**Anticholinergic:** Dry mouth.

**Central Nervous System:** Drowsiness is usually transitory and may disappear in a few days of continued therapy or upon reduction of dose. Involuntary motor activity including rare instances of tremor and convulsions have been reported, usually with doses considerably higher than those recommended. Clinically significant respiratory depression has not been reported at recommended doses.

In post-marketing experience, the following additional undesirable effects have been reported:

**Body as a Whole:** allergic reaction.

**Nervous System:** headache.

**Psychiatric:** hallucination.

**Skin and Appendages:** pruritus, rash, urticaria.

## **OVERDOSAGE**

The most common manifestation of hydroxyzine overdosage is hypersedation. Other reported signs and symptoms were convulsions, stupor, nausea and vomiting. As in the management of overdosage with any drug, it should be borne in mind that multiple agents may have been taken.

If vomiting has not occurred spontaneously, it should be induced. Immediate gastric lavage is also recommended. General supportive care, including frequent monitoring of the vital signs and close observation of the patient, is indicated. Hypotension, though unlikely, may be controlled with intravenous fluids and levarterenol, or metaraminol. Do not use epinephrine as hydroxyzine counteracts its pressor action.

There is no specific antidote. It is doubtful that hemodialysis would be of any value in the treatment of overdosage with hydroxyzine. However, if other agents such as barbiturates have been ingested concomitantly, hemodialysis may be indicated. There is no practical method to quantitate hydroxyzine in body fluids or tissue after its ingestion or administration.

## **DOSAGE AND ADMINISTRATION**

For symptomatic relief of anxiety and tension associated with psychoneurosis and as an adjunct in organic disease states in which anxiety is manifested: Adults, 50–100 mg q.i.d.; children under 6 years, 50 mg daily in divided doses; children over 6 years, 50–100 mg daily in divided doses.

For use in the management of pruritus due to allergic conditions such as chronic urticaria and atopic and contact dermatoses and in histamine-mediated pruritus: adults, 25 mg t.i.d. or q.i.d.; children under 6

years, 50 mg daily in divided doses; children over 6 years, 50–100 mg daily in divided doses.

As a sedative when used as a premedication and following general anesthesia: 50–100 mg for adults and 0.6 mg/kg of body weight in children.

When treatment is initiated by the intramuscular route of administration, subsequent doses may be administered orally.

As with all potent medication, the dosage should be adjusted according to the patient's response to therapy.

## **HOW SUPPLIED**

Hydroxyzine Hydrochloride Tablets, USP are available as:

10 mg – orange, round tablets debossed "3615" on one side and debossed "V" on the reverse side.

Supplied in bottles of:

- Bottles of 10: NDC 0603-3967-10
- Bottles of 100: NDC 0603-3967-21
- Bottles of 500: NDC 0603-3967-28
- Bottles of 1000: NDC 0603-3967-32

25 mg – green, round tablets debossed "3616" on one side and debossed "V" on the reverse side. Supplied in bottles of:

- Bottles of 10: NDC 0603-3968-10
- Bottles of 100: NDC 0603-3968-21
- Bottles of 500: NDC 0603-3968-28
- Bottles of 1000: NDC 0603-3968-32

50 mg – light orange, round tablets debossed "3617" on one side and debossed "V" on the reverse side.

Supplied in bottles of:

- Bottles of 10: NDC 0603-3969-10
- Bottles of 100: NDC 0603-3969-21
- Bottles of 500: NDC 0603-3969-28
- Bottles of 1000: NDC 0603-3969-32

**Dispense in a tight container as defined in the USP.**

**Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].**

Manufactured for:

**QUALITEST PHARMACEUTICALS**

Huntsville, AL 35811

8181154

R2/13-R2

And Relabeled By:

**Dispensing Solutions Inc.**

3000 West Warner Ave

Santa Ana, CA 92704

United States

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

NDC 66336-0086-XX

**BULK SOURCE DATA**  
MFD. FOR: QUALITEST PHARMACEUTICALS  
HUNTSVILLE, AL 35811

**PRODUCT ID:**  
GREEN ROUND TABLET  
DEBOSSED 3616 / V

**BULK SOURCE NDC: 00603-3968-32**  
MFR. LOT: XXXXXX  
PEDIGREE #: 0409042  
DISPENSE IN THIS  
TIGHT/LIGHT RESISTANT CONTAINER





**HYDROXYZINE HCl 25 mg**  
**XX TABLETS**  
**NDC 66336-0086-XX**  
**PRODUCT # 824-XX**

Rev. Date: 03/13

**EACH TABLET CONTAINS:**  
HYDROXYZINE  
HYDROCHLORIDE USP . . . . 25 mg

LOT# SAMPLE EXP: 00-00 Rx # 31758078  
**RX ONLY**

**WARNING: KEEP OUT OF CHILDREN'S REACH  
STORE AT 68° - 77° F. SEE USP.**

824- XX NDC 66336-0086- XX  
HYDROXYZINE HCl 25 mg  
XX TABLETS  
LOT # SAMPLE EXP: 00-00  
MN 00603-3968-32 RX# 31758078

824- XX NDC 66336-0086- XX  
HYDROXYZINE HCl 25 mg  
XX TABLETS  
LOT # SAMPLE EXP: 00-00  
MN 00603-3968-32 RX# 31758078

824- XX NDC 66336-0086- XX  
HYDROXYZINE HCl 25 mg  
XX TABLETS  
LOT # SAMPLE EXP: 00-00  
MN 00603-3968-32 RX# 31758078



Packaged Exclusively By:  
DISPENSING SOLUTIONS<sup>Phc</sup>  
Santa Ana, CA 92704

TAKE \_\_\_\_\_ ORALLY EVERY  
\_\_\_\_\_ HOURS \_\_\_\_\_ TIMES A DAY.  
\_\_\_\_\_ AS NEEDED  
AVOID ALCOHOL.  
MAY CAUSE DROWSINESS.

NDC 66336-0086-60

<b>HYDROXYZINE HYDROCHLORIDE</b>			
hydroxyzine hydrochloride tablet, film coated			
Product Information			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:66336-086(NDC:0603-3968)
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
HYDROXYZINE HYDROCHLORIDE (UNII: 76755771U3) (HYDROXYZINE - UNII:30S50YM8OG)	HYDROXYZINE HYDROCHLORIDE	25 mg	
Inactive Ingredients			
Ingredient Name	Strength		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
CALCIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: L11K75P92J)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)			
POVIDONES (UNII: FZ989GH94E)			
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			

## Product Characteristics

<b>Color</b>	GREEN	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	3616;V
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66336-086-60	60 in 1 BOTTLE, PLASTIC		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040574	05/27/2005	

**Labeler** - Dispensing Solutions, Inc. (066070785)

**Registrant** - PSS World Medical, Inc. (101822682)

## Establishment

Name	Address	ID/FEI	Business Operations
Dispensing Solutions, Inc.		066070785	relabel(66336-086) , repack(66336-086)

Revised: 8/2013

Dispensing Solutions, Inc.