

RECOTHROM- thrombin topical recombinant

Baxter Healthcare Corporation

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use RECOTHROM safely and effectively. See full prescribing information for RECOTHROM.

RECOTHROM Thrombin topical (Recombinant)

Lyophilized Powder for Solution - For Topical Use Only

Initial U.S. Approval: 2008

RECENT MAJOR CHANGES

Warnings and Precautions (5.1)

01/2024

INDICATIONS AND USAGE

RECOTHROM Thrombin topical (Recombinant), is a topical thrombin indicated to aid hemostasis whenever oozing blood and minor bleeding from capillaries and small venules is accessible and control of bleeding by standard surgical techniques (such as suture, ligature, or cautery) is ineffective or impractical in adults and pediatric populations greater than or equal to one month of age. (1)

RECOTHROM may be used in conjunction with an absorbable gelatin sponge, USP. (1)

DOSAGE AND ADMINISTRATION

For topical use only. DO NOT INJECT. (2)

Reconstitute RECOTHROM Thrombin topical (Recombinant) powder with sterile 0.9% sodium chloride, USP, yielding a solution containing 1000 units (international units of potency) per mL. (2.1)

Apply RECOTHROM solution directly to bleeding site surface or in conjunction with absorbable gelatin sponge. The amount required depends upon the area of tissue to be treated. (2.2)

5000-unit RECOTHROM may be used in conjunction with FLOSEAL NT Hemostatic Matrix, 5mL (2.2)

DOSAGE FORMS AND STRENGTHS

RECOTHROM Thrombin topical (Recombinant) is available as 5000-unit and 20,000-unit vials of sterile recombinant topical thrombin lyophilized powder for solution. When reconstituted as directed, the final solution contains 1000 units/mL of RECOTHROM. (3)

CONTRAINDICATIONS

- Do not inject directly into the circulatory system. (4)
- Do not use for the treatment of massive or brisk arterial bleeding. (4)
- Do not administer to patients with a history of hypersensitivity to RECOTHROM Thrombin topical (Recombinant), any components of RECOTHROM or hamster proteins. (4)

WARNINGS AND PRECAUTIONS

- RECOTHROM Thrombin topical (Recombinant) causes thrombosis if it enters the circulatory system. (5.1)
- Hypersensitivity reactions, including anaphylaxis, may occur. (5.2)

ADVERSE REACTIONS

- The most common adverse reaction (incidence 6%) was thromboembolic events. (5.1, 6)
- Antibody formation to RECOTHROM Thrombin topical (Recombinant) occurred in <1% of patients. None of the antibodies detected neutralized native human thrombin. (6.2)

To report SUSPECTED ADVERSE REACTIONS, contact Baxter Healthcare at 1-866-888-2472 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 1/2024

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE**2 DOSAGE AND ADMINISTRATION**

2.1 Reconstitution of RECOTHROM Thrombin topical (Recombinant)

2.2 Application Techniques

3 DOSAGE FORMS AND STRENGTHS**4 CONTRAINDICATIONS****5 WARNINGS AND PRECAUTIONS**

5.1 Thrombosis

5.2 Hypersensitivity Reactions

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Immunogenicity

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

11 DESCRIPTION**12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES**15 REFERENCES****16 HOW SUPPLIED/STORAGE AND HANDLING****17 PATIENT COUNSELING INFORMATION**

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION**1 INDICATIONS AND USAGE**

RECOTHROM Thrombin topical (Recombinant), is a topical thrombin indicated to aid hemostasis whenever oozing blood and minor bleeding from capillaries and small venules is accessible and control of bleeding by standard surgical techniques (such as suture, ligature, or cautery) is ineffective or impractical in adults and pediatric populations greater than or equal to one month of age.

RECOTHROM may be used in conjunction with an absorbable gelatin sponge, USP.

2 DOSAGE AND ADMINISTRATION

For topical use only. DO NOT INJECT.

2.1 Reconstitution of RECOTHROM Thrombin topical (Recombinant)

The volume of reconstituted RECOTHROM Thrombin topical (Recombinant) required will vary depending on the size and number of bleeding sites to be treated and the method of application.

Inspect the integrity of the RECOTHROM package and contents. Do not use if the packaging or contents have been damaged or opened.

Reconstitute the lyophilized powder using the supplied diluent. Use aseptic technique when handling vials and syringes.

5000-unit RECOTHROM Thrombin topical (Recombinant) Reconstitution

Units used herein represent international units of potency determined using a reference standard that has been calibrated against the World Health Organization Second International Standard for Thrombin.

1. Remove flip-off cap from the top of the RECOTHROM vial.
2. Attach the needle-free transfer device and snap it into place on the vial by placing the vial on a flat surface and attaching the transfer device straight into the center of the vial stopper.
3. Attach the prefilled diluent syringe to the needle-free transfer device.
4. Inject the 5 mL of diluent from the syringe into the product vial. Keep the syringe plunger depressed.
5. DO NOT reuse the diluent syringe for transfer of the reconstituted product. Remove and discard the diluent syringe.
6. Gently swirl and invert the product vial until the powder is completely dissolved. Avoid excessive agitation. The powder should dissolve in less than one minute at room temperature.
7. Apply the pre-printed "DO NOT INJECT" label to the sterile, empty transfer syringe provided, then draw up the RECOTHROM solution.

20,000-unit RECOTHROM Thrombin topical (Recombinant) Reconstitution

1. Remove the flip-off cap from the top of the RECOTHROM vial and the diluent vial.
2. Attach a needle-free transfer device (one each) to the RECOTHROM and diluent vials and snap them into place by placing the vial on a flat surface and attaching the transfer device straight into the center of the vial stopper.
3. Open the sterile, empty 20-mL syringe package.
4. Attach the 20-mL syringe to the needle-free transfer device on the diluent vial (injection of air into the diluent vial may facilitate withdrawal of the diluent).
5. Draw up 20 mL of diluent from the vial into the syringe.
6. Remove the diluent-filled syringe from the diluent vial and attach it to the transfer device on the RECOTHROM vial.
7. Transfer the 20 mL of diluent from the syringe into the RECOTHROM vial; the vacuum in the vial facilitates transfer.
8. Leave the syringe attached and gently swirl and invert the RECOTHROM vial until the powder is completely dissolved. Avoid excessive agitation. The powder should dissolve in less than one minute at room temperature.
9. Apply the pre-printed 'DO NOT INJECT' label to the syringe. Draw up the RECOTHROM solution.

2.2 Application Techniques

Topically apply RECOTHROM Thrombin topical (Recombinant) solution directly or in conjunction with absorbable gelatin sponge onto the bleeding site. DO NOT INJECT.

The amount required depends upon the area of tissue to be treated and the method of application.

Vials are for single use only. Discard unused contents.

Use with Absorbable Gelatin Sponge

Refer to the absorbable gelatin sponge labeling for safety information and instructions on appropriate use.

1. Transfer solution from syringe to a sterile bowl or basin.
2. Place the desired size pieces of the absorbable gelatin sponge into the bowl containing reconstituted RECOTHROM to completely saturate the sponge(s).
3. Remove the saturated sponge(s) and squeeze gently to remove excess RECOTHROM.
4. Apply the sponge to the bleeding site in a single layer.

Use with RECOTHROM Thrombin Spray Applicator Kit

1. Hold the outer sealed tray, peel back the lid, and aseptically transfer the inner sealed sterile tray to the sterile field.
2. Open the inner tray seal and use the sterile bowl as the receptacle for reconstituted RECOTHROM solution.
3. Refer to Spray Applicator Kit instructions for spray pump and syringe spray assembly and use.

Use in conjunction with FLOSEAL NT Hemostatic Matrix, 5mL

1. 5000-unit RECOTHROM may be used with FLOSEAL NT Hemostatic Matrix, 5mL according to the FLOSEAL NT Hemostatic Matrix, 5mL instructions for use (IFU).

3 DOSAGE FORMS AND STRENGTHS

RECOTHROM Thrombin topical (Recombinant) is available as a sterile lyophilized powder in 5000- and 20,000-unit single-use vials. When reconstituted with the sterile 0.9% sodium chloride, USP provided, the powder yields a solution containing 1000 units/mL of Thrombin topical (Recombinant).

4 CONTRAINDICATIONS

- Do not inject directly into the circulatory system.
- Do not use for the treatment of massive or brisk arterial bleeding.
- Do not administer to patients with a history of hypersensitivity to RECOTHROM Thrombin topical (Recombinant) or any components of RECOTHROM.
- Do not use in patients with known hypersensitivity to hamster proteins.

5 WARNINGS AND PRECAUTIONS

5.1 Thrombosis

RECOTHROM Thrombin topical (Recombinant) causes thrombosis if it enters the circulatory system. Apply topically. DO NOT INJECT.

5.2 Hypersensitivity Reactions

Hypersensitivity reactions, including anaphylaxis, may occur.

RECOTHROM Thrombin topical (Recombinant) is produced in a genetically modified Chinese Hamster Ovary (CHO) cell line and may contain hamster or snake proteins [see *Contraindications (4) and Description (11)*].

6 ADVERSE REACTIONS

Thromboembolic adverse reactions were reported in 6% of surgical patients treated with RECOTHROM Thrombin topical (Recombinant) in all completed clinical trials (N=644) [see *Warnings and Precautions (5.1)*].

Antibody formation to RECOTHROM occurred in <1% of patients. None of the antibodies detected neutralized native human thrombin [see *Adverse Reactions (6.2)*].

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug product cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Clinical trials have been performed with RECOTHROM Thrombin topical (Recombinant) applied with absorbable gelatin sponge and applied with a spray applicator. A total of 644 patients were exposed to RECOTHROM in these studies.

RECOTHROM Thrombin topical (Recombinant) Used in Conjunction with Absorbable Gelatin Sponge

Four hundred eleven (411) patients were treated in a randomized, double-blind, controlled trial that compared RECOTHROM to bovine thrombin. Both thrombins were applied with a gelatin sponge in patients undergoing spinal surgery, hepatic resection, peripheral arterial bypass surgery, or arteriovenous graft formation for hemodialysis access.¹ The incidence of thromboembolic adverse reactions was similar between the RECOTHROM and bovine thrombin treatment groups (see Table 1).

Table 1: Incidence of Adverse Reactions with RECOTHROM Thrombin topical (Recombinant) and Bovine Thrombin

Adverse Reaction Category	RECOTHROM (N=205) n (%)	Bovine Thrombin* (N=206) n (%)
Thromboembolic events	11 (5%)	12 (6%)

* THROMBIN-JMI Thrombin, Topical (Bovine)

In an open-label, single-group trial (N=209), patients with documented or highly likely

prior exposure to bovine thrombin within the previous three years were treated with RECOTHROM when undergoing surgeries (spinal, peripheral arterial bypass, or arteriovenous graft formation for hemodialysis access).² The incidence of thromboembolic adverse reactions in this study was 9%.

In an open-label, single-group trial of re-exposure to RECOTHROM (N=31), patients with documented prior exposure to RECOTHROM were treated with RECOTHROM during surgery (spinal, peripheral arterial bypass, arteriovenous graft formation, or other procedures).³ The incidence of thromboembolic adverse reactions in this study was 3%.

In other randomized, double-blind trials across a range of surgical settings (N=130; spinal surgery, hepatic resection, peripheral arterial bypass surgery, or arteriovenous graft formation for hemodialysis access), the safety of RECOTHROM (n=88 patients) was compared to placebo (RECOTHROM excipients reconstituted with sterile 0.9% sodium chloride, USP) (n=42 patients). The incidence of thromboembolic adverse reactions in this study was 5% for RECOTHROM and 12% for placebo.

RECOTHROM Thrombin topical (Recombinant) Applied with Spray Applicator

RECOTHROM was applied with a spray applicator in two open-label clinical trials: a single-group trial in adult and pediatric burn patients (N=72; ≤ 16 years of age, (n=4) and ≥ 17 years of age, (n=68)) treated with RECOTHROM applied to the wound excision site prior to autologous skin grafting⁴; and in a single-group trial in pediatric patients (one month to 17 years of age) undergoing synchronous burn wound excision and autologous skin grafting (N=30; ≤ 16 years of age, (n=26); ≥ 17 years of age, (n=4)).⁵ In the first study, the incidence of thromboembolic adverse reactions was 1%. In the second study, there were no reported thromboembolic adverse reactions [*see Use in Specific Populations (8.4)*].

6.2 Immunogenicity

The detection of antibody formation is highly dependent upon the sensitivity and specificity of the assay. The absolute immunogenicity rates reported here are difficult to compare with the results from other products due to differences in assay methodology, patient populations, and other underlying factors.

The potential for development of antibodies to RECOTHROM Thrombin topical (Recombinant) was evaluated in multiple clinical trials and included patients with a single exposure to RECOTHROM as well as patients who were re-exposed to RECOTHROM during a subsequent surgical procedure. Only patients with both baseline and post-treatment antibody specimens available were evaluated for the development of specific anti-RECOTHROM product antibodies, which was defined as seroconversion or a ≥ 1.0 titer unit (≥ 10 -fold) increase in antibody levels after study treatment. Five of 609 (0.8%; 95% CI, 0.4%-2.8%) evaluable patients developed specific anti-RECOTHROM product antibodies. None of these antibodies were found to neutralize native human thrombin. There was no difference in anti-RECOTHROM product antibody formation incidence among patients exposed to RECOTHROM applied with absorbable gelatin sponge, USP or with spray applicator.

In a clinical trial comparing RECOTHROM to bovine thrombin (N=411; n=398 antibody evaluable) for the development of specific anti-product antibodies, blood samples were collected at baseline and at Day 29 in both treatment groups and were analyzed by ELISA.¹ At baseline, 1.5% of RECOTHROM patients (n=3/198) had positive anti-product

antibody titers compared with 5% of bovine thrombin patients (n=10/200). Of these patients, none of the RECOTHROM group and eight in the bovine thrombin group exhibited ≥ 1.0 titer unit (≥ 10 -fold) increases in anti-product antibody levels after study treatment.

At Day 29, three of 198 (1.5%; 95% CI, 0%-4%) patients in the RECOTHROM group developed specific anti-product antibodies (one patient also developed anti-CHO host cell protein antibodies); 43 of 200 patients in the bovine thrombin group (22%; 95% CI, 16%-28%) developed specific antibodies to bovine thrombin product. Treatment with RECOTHROM resulted in a statistically significant lower incidence of specific anti-product antibody development. Because the study was not powered to detect a difference in clinical outcomes attributable to antibody formation, no conclusions can be drawn regarding the clinical significance of the difference in antibody formation based on the results of this study. None of the antibodies in the RECOTHROM group neutralized native human thrombin. Antibodies against bovine thrombin product were not tested for neutralization of native human thrombin.

In a trial of patients with a high likelihood of prior exposure to bovine thrombin, 15.6% of patients (n=32/205) had anti-bovine thrombin product antibodies and 2% of patients (n=4/200) had anti-RECOTHROM product antibodies at baseline.² Following treatment, none of the 200 evaluable patients (patients for whom post-treatment specimens were available) developed antibodies to RECOTHROM.

In a trial of patients previously exposed to RECOTHROM, 31 patients were re-exposed to RECOTHROM during a subsequent surgery.³ None of the evaluable patients (n=30) had anti-RECOTHROM product antibodies at baseline and none developed antibodies at Day 29.

In a trial of RECOTHROM, including 26 pediatric patients (aged one month to 16 years) and four patients 17 years of age, one patient without prior thrombin exposure had pre-existing anti-RECOTHROM product antibodies at baseline.⁵ None of the 27 evaluable patients developed anti-RECOTHROM product antibodies at Day 29.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no available data regarding RECOTHROM Thrombin topical (Recombinant) in pregnant women. No animal reproductive and developmental toxicity studies have been conducted with RECOTHROM.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

8.2 Lactation

Risk Summary

There is no information regarding the presence of RECOTHROM Thrombin topical (Recombinant) in human milk, the effects on the breastfed infant, and the effects on milk production.

The developmental and health benefits of breastfeeding should be considered along with

the mother's clinical need for RECOTHROM and any potential adverse effects on the breastfed child from RECOTHROM or from underlying maternal condition.

8.4 Pediatric Use

A total of 30 pediatric patients, ages 0 to 16 years (one month to 2 years, n=10; 2 to 12 years, n=12; 12 to 16 years, n=8), were treated in clinical trials with RECOTHROM Thrombin topical (Recombinant) using a spray applicator to burn wound excision sites prior to autologous skin grafting. No patient experienced a thromboembolic adverse reaction. The safety of RECOTHROM in pediatric patients greater than or equal to one month of age is supported by these data and by extrapolation of efficacy from adequate and well-controlled studies of RECOTHROM in adults. Safety and efficacy have not been established in neonates [see *Adverse Reactions (6)*].

8.5 Geriatric Use

Of 644 patients in clinical studies of RECOTHROM Thrombin topical (Recombinant), 36% (n=232/644) were ≥ 65 years old and 15% (n=95/644) were ≥ 75 years old.

No differences in safety or effectiveness were observed between these patients and younger patients, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

11 DESCRIPTION

RECOTHROM Thrombin topical (Recombinant), is a human coagulation protein produced via recombinant DNA technology from a genetically modified CHO cell line. RECOTHROM is identical in amino acid sequence and structurally similar to naturally occurring human thrombin. RECOTHROM precursor is secreted to culture medium as single chain form that is proteolytically converted to a two-chain active form (using a protein derived from snakes) and is purified by a chromatographic process that yields a product having hemostatic activities similar to native human thrombin. The cell line used to manufacture RECOTHROM has been tested and shown to be free of known infectious agents. The cell culture process used in the manufacture of RECOTHROM employs no additives of human or animal origin. The purification process includes solvent-detergent treatment and nano-filtration steps dedicated to viral clearance.

RECOTHROM is provided as a sterile, white to off-white, preservative-free, lyophilized powder in vials for reconstitution with diluent (sterile 0.9% sodium chloride, USP). Reconstitution with the provided diluent, as described [see *Dosage and Administration (2.1)*], yields a solution with a pH of 6.0 containing 1000 units/mL of recombinant thrombin for topical use. The formulated product is a clear, colorless solution upon reconstitution and contains the following excipients: histidine, mannitol, sucrose, polyethylene glycol 3350, sodium chloride, and calcium chloride dihydrate, USP.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

RECOTHROM Thrombin topical (Recombinant), is a specific human serine protease that

promotes hemostasis and acts locally when applied topically to a site of bleeding.

RECOTHROM activates platelets and catalyzes the conversion of fibrinogen to fibrin, which are steps that are essential for blood clot formation.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In vitro cytotoxicity studies have been performed in mouse L929 fibroblast cell cultures and

demonstrate a concentration-dependent effect on cell morphology. The thrombin-induced

morphological changes were similar to those seen with bovine thrombin.

13.2 Animal Toxicology and/or Pharmacology

In a study in nonhuman primates, RECOTHROM Thrombin topical (Recombinant) was applied directly to a liver wound with an absorbable gelatin sponge, USP. In a second study, RECOTHROM was administered subcutaneously once weekly for four weeks to nonhuman primates following repeat doses of 5405 units/m². In both studies, RECOTHROM had no effect on clinical signs, serum chemistry, coagulation parameters, or histopathology; only normal postsurgical findings were observed. No animals developed anti-RECOTHROM product antibodies in either study.

RECOTHROM was found to be non-irritating when instilled in the eyes (200 units) or applied to normal or abraded skin of rabbits (up to 1000 units/site).

To evaluate RECOTHROM inhibition and clearance from the bloodstream, radiolabeled RECOTHROM was administered intravenously or subcutaneously to nonhuman primates and applied with an absorbable gelatin sponge, USP, in a rabbit hepatic wound model. RECOTHROM did not circulate in the blood as free, active molecule, but was rapidly inactivated (<5 minutes) after formation of complexes with endogenous inhibitors (e.g., antithrombin III); these complexes were cleared by the liver. RECOTHROM applied with an absorbable gelatin sponge, USP, was shown to decrease time to hemostasis (TTH) when compared to saline in a rabbit hepatic wound model and rat heminephrectomy model. RECOTHROM also reduced TTH when directly applied in a porcine partial-thickness excisional skin-wound model as compared to saline control (or no treatment).

RECOTHROM applied with a gauze sponge decreased TTH in a concentration-dependent manner in both the rabbit and rat models. Concentrations of RECOTHROM >1000 units/mL were no different than 1000 units/mL while the effect of RECOTHROM diluted to a concentration of 100 units/mL on TTH was indistinguishable from placebo.

The performance of RECOTHROM mixed with gelatin matrix (FLOSEAL NT Hemostatic Matrix, 5mL) was compared to FLOSEAL Hemostatic Matrix, 5mL (human plasma-derived thrombin mixed with gelatin matrix) in a blinded nonclinical study using a heparinized porcine liver punch model of moderate to severe bleeding. The analysis demonstrated that the performance of RECOTHROM mixed with gelatin matrix was non-inferior to human plasma-derived thrombin mixed with gelatin matrix.

14 CLINICAL STUDIES

RECOTHROM Thrombin topical (Recombinant) was evaluated in a randomized, double-blind comparative clinical trial to bovine thrombin. Each thrombin was topically applied to bleeding sites with an absorbable gelatin sponge at a nominal concentration of 1000 units/mL.¹ Patients (N=411) were a heterogeneous surgical population undergoing surgery in one of four surgical settings: spinal surgery (n=122, 30%), hepatic resection (n=125, 30%), peripheral arterial bypass surgery (n=88, 21%), and arteriovenous graft formation for hemodialysis access (n=76, 18%). Patients with prior heparin-induced thrombocytopenia were excluded. Patient ages ranged from 21 to 89 years, gender was 53% male and 47% female, and the distribution by race was 68% white, 18% black or African American, and 14% other. The distribution of these characteristics was similar in both the RECOTHROM and bovine thrombin treatment groups.

The objectives of the study were to evaluate the comparative efficacy, safety, and immunogenicity of RECOTHROM and bovine thrombin in combination with an absorbable gelatin sponge as adjuncts to hemostasis in surgery. Efficacy was evaluated by the incidence of hemostasis within 10 minutes. Bleeding appropriate for evaluation was defined as mild to moderate bleeding, either on its own or remaining after brisk bleeding was controlled by standard surgical modalities. Although multiple bleeding sites could be treated, only one bleeding site per patient was selected to determine effectiveness.

Table 2 summarizes the incidence of hemostasis within 10 minutes for each treatment for the 401

efficacy evaluable patients. The incidence of hemostasis within 10 minutes was comparable for the RECOTHROM and bovine thrombin groups.

Table 2: Hemostasis Within 10 Minutes*

	RECOTHROM (N=198) (%)	Bovine Thrombin[†] (N=203) (%)
Overall	95%	95%
Spinal surgery	98%	98%
Hepatic resection	98%	97%
Peripheral arterial bypass	85%	86%
Arteriovenous graft formation	97%	97%

* Evaluation of hemostasis at ≤ 10 minutes for patients treated at 1 of 4 primary TTH bleeding site types: epidural venous plexus, hepatic resection site, peripheral arterial bypass proximal anastomosis, and arteriovenous graft arterial anastomosis.

† THROMBIN-JMI Thrombin, Topical (Bovine)

The percentage of patients achieving hemostasis at 1.5, 3, 6, and 10 minutes is listed in Table 3.

Table 3: Cumulative Incidence of Hemostasis Over Time

Time (Minutes)	RECOTHROM (N=198) (%)	Bovine Thrombin* (N=203) (%)
1.5	48%	46%

3	81%	72%
6	92%	88%
10	95%	95%

* THROMBIN-JMI Thrombin, Topical (Bovine)

15 REFERENCES

1. Chapman WC, Singla N, Genyk Y, et al. A phase 3, randomized, double-blind comparative study of the efficacy and safety of topical recombinant human thrombin and bovine thrombin in surgical hemostasis. *J Am Coll Surg.* 2007;205(2):256-265.
2. Singla NK, Ballard JL, Moneta G, et al. A phase 3b open-label, single-group immunogenicity and safety study of topical recombinant thrombin in surgical hemostasis. *J Am Coll Surg.* 2009;209(1):68-74.
3. Singla NK, Gasparis AP, Ballard JL, et al. Immunogenicity and safety of re-exposure to recombinant human thrombin in surgical hemostasis. *J Am Coll Surg.* 2011;213(6):722-727.
4. Greenhalgh DG, Gamelli RL, Collins J, et al. Recombinant thrombin: safety and immunogenicity in burn wound excision and grafting. *J Burn Care Res.* 2009;30(3):371-379.
5. Foster KN, Mullins RF, Greenhalgh DG, et al. Recombinant human thrombin: safety and immunogenicity in pediatric burn wound excision. *J Ped Surg.* 2011;46(10):1992-1999.

16 HOW SUPPLIED/STORAGE AND HANDLING

RECOTHROM Thrombin topical (Recombinant), is supplied in single-use, preservative-free vials in the following packages:

NDC 0338-0322-01

A 5000-unit vial of RECOTHROM with a 5-mL prefilled diluent syringe (containing sterile 0.9% sodium chloride, USP), a sterile needle-free transfer device, a 5-mL sterile empty syringe, and a pre-printed label.

NDC 0338-0326-01

A 20,000-unit vial of RECOTHROM with a 20-mL vial of diluent (containing sterile 0.9% sodium chloride, USP), 2 sterile needle-free transfer devices, a 20-mL sterile empty syringe, and a pre-printed label.

NDC 0338-0330-01

The 20,000-unit RECOTHROM kit co-packaged with RECOTHROM Spray Applicator Kit containing a spray pump, a spray bottle, a syringe spray tip, a syringe, a bowl, and 2 blank labels.

No RECOTHROM kit components contain latex.

Store RECOTHROM sterile powder vials at 2°C to 25°C (36°F to 77°F). Do not freeze.

Reconstituted solutions of RECOTHROM prepared with sterile 0.9% sodium chloride, USP, may be stored for up to 24 hours at 2°C to 25°C (36°F to 77°F). Discard reconstituted solution after 24 hours.

17 PATIENT COUNSELING INFORMATION

Because topical thrombin may cause the formation of clots in blood vessels if absorbed systemically, advise patients to consult their physician if they experience leg tenderness or swelling, chest pain, shortness of breath, or difficulty speaking or swallowing [see *Warnings and Precautions (5.1)*].

Baxter and Recothrom are registered trademarks of Baxter International Inc. Thrombin-JMI is a registered trademark of King Pharmaceuticals Research and Development, LLC.

Manufactured for:


Baxter Healthcare Corporation

Deerfield, IL 60015 USA

US License No. 0140

07-19-00-7463

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL



Baxter
Recothrom
THROMBIN TOPICAL
[RECOMBINANT]

FOR TOPICAL USE ONLY - DO NOT INJECT

Manufactured for:
Baxter Healthcare Corporation
Deerfield, IL 60015 USA
US License No. 0140
Made in Italy
Rx only
Store at 2-25°C (36-77°F).
Do not freeze.
See package insert
Full Prescribing Information.
No components
contain latex.
No preservative.

NDC: 0338-0322-01

5,000 units

5,000 units

NDC: 0338-0322-01

5,000 units

FOR TOPICAL USE ONLY - DO NOT INJECT

Baxter
Recothrom
THROMBIN TOPICAL
[RECOMBINANT]

Manufactured for:
Baxter Healthcare Corporation
Deerfield, IL 60015 USA
US License No. 0140
Made in Italy

FPO


FPO

Rx only

CONTENTS:

- One single-use vial containing 5,000 International Units of sterile Thrombin topical (Recombinant)
- One 5-mL prefilled diluent syringe (containing sterile 0.9% sodium chloride injection, USP)
- One sterile needle-free transfer device
- One sterile 5-mL empty syringe for transfer
- One syringe label

PS



Carton Label

PS

5,000 units

NDC: 0338-0322-01

FOR TOPICAL USE ONLY - DO NOT INJECT

Baxter Logo

RECOTHROM
 THROMBIN TOPICAL
 (RECOMBINANT)

Manufactured for:
Baxter Healthcare Corporation
 Deerfield, IL 60015 USA
 US License No. 0140
 Made in Italy

FPO

FPO

5,000 units

Rx only

CONTENTS:

- One single-use vial containing 5,000 International Units of sterile Thrombin topical (Recombinant)
- One 5-mL prefilled diluent syringe (containing sterile 0.9% sodium chloride injection, USP)
- One sterile needle-free transfer device
- One sterile 5-mL empty syringe for transfer
- One syringe label

PS

5,000 units

NDC: 0338-0322-01

FOR TOPICAL USE ONLY - DO NOT INJECT

Baxter Logo

RECOTHROM

THROMBIN TOPICAL

(RECOMBINANT)

Manufactured for: Baxter Healthcare Corporation Deerfield, IL 60015 USA US License No. 0140 Baxter and Recothrom are registered trademarks of Baxter International Inc. Made in Italy

Rx only

CONTENTS:

- One single-use vial containing 5,000 International Units of sterile Thrombin topical (Recombinant)
- One 5-mL prefilled diluent syringe (containing sterile 0.9% sodium chloride injection, USP)
- One sterile needle-free transfer device
- One sterile 5-mL empty syringe for transfer
- One syringe label

FPO

5,000 UNITS

Rx only

Store at 2-25°C (36-77°F).

Do not freeze.

See package insert

Full Prescribing Information.

Baxter Logo

07-01-00-0673

5,000 units

NDC: 0338-0322-01

FOR TOPICAL USE ONLY - DO NOT INJECT

Baxter Logo

RECOTHROM

THROMBIN TOPICAL
(RECOMBINANT)

Manufactured for:

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

US License No. 0140

Made in Italy

Rx only

Store at 2-25°C (36-77°F).

Do not freeze.

See package insert

Full Prescribing Information.

No components

contain latex.

No preservative.

Baxter and Recothrom are registered trademarks of Baxter International Inc.

Manufactured for:
Baxter Healthcare Corporation
Deerfield, IL 60015 USA
US License No. 0140
Made in Italy

Rx only
Store at 2-25°C (36-77°F).
Do not freeze.
See package insert
Full Prescribing Information.
No components
contain latex.
No preservative.

Recothrom

THROMBIN TOPICAL

[RECOMBINANT]

FOR TOPICAL USE ONLY - DO NOT INJECT

NDC: 0338-0326-01

20,000 units

20,000 units

FOR TOPICAL USE ONLY - DO NOT INJECT

Baxter

Recothrom

THROMBIN TOPICAL

[RECOMBINANT]

Rx only
Store at 2-25°C (36-77°F).
Do not freeze.
See package insert
Full Prescribing Information.
Manufactured for:
Baxter Healthcare Corporation
Deerfield, IL 60015 USA
US License No. 0140
Made in Italy

FPO

FPO



NDC: 0338-0326-01



20,000 units

NDC: 0338-0326-01

CONTENTS:

- One single-use vial containing 20,000 International Units of sterile Thrombin topical (Recombinant)
- One vial diluent (containing 20-mL sterile 0.9% sodium chloride injection, USP)
- Two sterile needle-free transfer devices
- One sterile 20-mL empty syringe for transfer
- One syringe label

Rx only
Store at 2-25°C (36-77°F).
Do not freeze.
See package insert
Full Prescribing Information.





20,000 units

NDC: 0338-0326-01

20,000 units

NDC: 0338-0326-01

FOR TOPICAL USE ONLY - DO NOT INJECT

Baxter
Recothrom
THROMBIN TOPICAL
[RECOMBINANT]

Rx only
Store at 2-25°C (36-77°F).
Do not freeze.
See package insert
Full Prescribing Information.

CONTENTS:

- One single-use vial containing 20,000 International Units of sterile Thrombin topical (Recombinant)
- One vial diluent (containing 20-mL sterile 0.9% sodium chloride injection, USP)
- Two sterile needle-free transfer devices
- One sterile 20-mL empty syringe for transfer
- One syringe label


FOR TOPICAL USE ONLY - DO NOT INJECT


Baxter
Recothrom
THROMBIN TOPICAL
[RECOMBINANT]

Rx only
Store at 2-25°C (36-77°F).
Do not freeze.
See package insert
Full Prescribing Information.

07-01-00-0675

FPO





Carton Label

20,000 units

NDC: 0338-0326-01

FOR TOPICAL USE ONLY - DO NOT INJECT

Baxter Logo

RECOTHROM

THROMBIN TOPICAL
(RECOMBINANT)

Rx only

Store at 2-25°C (36-77°F).

Do not freeze.

See package insert

Full Prescribing Information.

Manufactured for:

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

US License No. 0140

Made in Italy

FPO

FPO

20,000 units
NDC: 0338-0326-01

Rx only
Store at 2-25°C (36-77°F).
Do not freeze.

See package insert
Full Prescribing Information.

CONTENTS:

- One single-use vial containing 20,000 International Units of sterile Thrombin topical (Recombinant)
- One vial diluent (containing 20-mL sterile 0.9% sodium chloride injection, USP)
- Two sterile needle-free transfer devices
- One sterile 20-mL empty syringe for transfer
- One syringe label

20,000 units
NDC: 0338-0326-01

FOR TOPICAL USE ONLY - DO NOT INJECT

Baxter Logo

RECOTHROM
THROMBIN TOPICAL
(RECOMBINANT)

Rx only
Store at 2-25°C (36-77°F).
Do not freeze.

See package insert
Full Prescribing Information.

CONTENTS:

- One single-use vial containing 20,000 International Units of sterile Thrombin topical (Recombinant)
- One vial diluent (containing 20-mL sterile 0.9% sodium chloride injection, USP)
- Two sterile needle-free transfer devices
- One sterile 20-mL empty syringe for transfer
- One syringe label

FPO

20,000 units
NDC: 0338-0326-01

FOR TOPICAL USE ONLY - DO NOT INJECT

Baxter Logo

RECOTHROM
THROMBIN TOPICAL
(RECOMBINANT)

Rx only
Store at 2-25°C (36-77°F).
Do not freeze.
See package insert
Full Prescribing Information.

07-01-00-0675

20,000 units
NDC: 0338-0326-01

FOR TOPICAL USE ONLY - DO NOT INJECT

Baxter Logo


RECOTHROM
THROMBIN TOPICAL
(RECOMBINANT)

Manufactured for:
Baxter Healthcare Corporation
Deerfield, IL 60015 USA
US License No. 0140
Made in Italy

Rx only
Store at 2-25°C (36-77°F).
Do not freeze.
See package insert
Full Prescribing Information.

No Components
contain latex.
No preservative.

Baxter and Recothrom are registered trademarks of Baxter International Inc.



Carton Label

20,000 units
NDC: 0338-0330-01

FOR TOPICAL USE ONLY - DO NOT INJECT
RECOTHROM

THROMBIN TOPICAL
(RECOMBINANT)

20,000 units

WITH SPRAY APPLICATOR KIT

Rx only

Store at 2-25°C (36-77°F).

Do not freeze.

See package insert

Full Prescribing Information.

No components contain latex.

No preservative.

RECOTHROM CONTENTS:

- One single-use vial containing 20,000 International Units of sterile Thrombin topical (Recombinant)
- One vial diluent (containing 20-mL sterile 0.9% sodium chloride injection, USP)
- Two sterile needle-free transfer devices
- One sterile 20-mL empty syringe for transfer
- One syringe label

SPRAY APPLICATOR CONTENTS:

- One spray pump
- One spray bottle
- One syringe spray tip
- One 20-mL syringe
- One 30-mL bowl
- Two blank labels
- Instructions for use

Manufactured for:

Baxter Healthcare Corporation

Deerfield, IL 60015 uSA

US License No. 0140

Made in Italy

Baxter Logo

Baxter and Recothrom are registered trademarks of Baxter International Inc.

20,000 units

NDC: 0338-0330-01

FOR TOPICAL USE ONLY - DO NOT INJECT

RECOTHROM

THROMBIN TOPICAL
(RECOMBINANT)

20,000 units

WITH SPRAY APPLICATOR KIT

Manufactured for:

Baxter Healthcare Corporation

Deerfield, IL 60015 uSA

US License No. 0140

Made in Italy

Baxter Logo

07-03-00-0614

FPO

20,000 units

NDC: 0338-0330-01

FOR TOPICAL USE ONLY - DO NOT INJECT

RECOTHROM

THROMBIN TOPICAL
(RECOMBINANT)

20,000 units

WITH SPRAY APPLICATOR KIT

Rx only

Store at 2-25°C (36-77°F).

Do not freeze.

See package insert

Full Prescribing Information.

No components contain latex.

No preservative.

07-03-00-0614

20,000 units

NDC: 0338-0330-01

FOR TOPICAL USE ONLY - DO NOT INJECT

RECOTHROM

THROMBIN TOPICAL
(RECOMBINANT)

20,000 units

WITH SPRAY APPLICATOR KIT

Rx only

Store at 2-25°C (36-77°F).

Do not freeze.

See package insert

Full Prescribing Information.

No components contain latex.

No preservative.

RECOTHROM CONTENTS:

- One single-use vial containing 20,000 International Units of sterile Thrombin topical (Recombinant)
- One vial diluent (containing 20-mL sterile 0.9% sodium chloride injection, USP)
- Two sterile needle-free transfer devices
- One sterile 20-mL empty syringe for transfer
- One syringe label

SPRAY APPLICATOR CONTENTS:

- One spray pump
- One spray bottle
- One syringe spray tip
- One 20-mL syringe
- One 30-mL bowl
- Two blank labels
- Instructions for use

FPO

FPO

20,000 units

NDC: 0338-0330-01

***Baxter* Logo**

RECOTHROM

THROMBIN TOPICAL

(RECOMBINANT)

20,000 units

WITH SPRAY APPLICATOR KIT

FOR TOPICAL USE ONLY - DO NOT INJECT

Rx only

Store at 2-25°C (36-77°F).

Do not freeze.

See package insert

Full Prescribing Information.

No components contain latex.

No preservative.

RECOTHROM CONTENTS:

- One single-use vial containing 20,000 International Units of sterile Thrombin topical (Recombinant)
- One vial diluent (containing 20-mL sterile 0.9% sodium chloride injection, USP)
- Two sterile needle-free transfer devices
- One sterile 20-mL empty syringe for transfer
- One syringe label

SPRAY APPLICATOR CONTENTS:

- One spray pump
- One spray bottle
- One syringe spray tip
- One 20-mL syringe
- One 30-mL bowl
- Two blank labels
- Instructions for use

RECOTHROM				
thrombin topical recombinant kit				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-0322	
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0322-01	1 in 1 KIT; Type 1: Convenience Kit of Co-Package	01/29/2008	
Quantity of Parts				
Part #	Package Quantity		Total Product Quantity	
Part 1	1 VIAL		5 mL	
Part 2	1 SYRINGE		5 mL	
Part 1 of 2				
RECOTHROM				
thrombin topical recombinant powder, for solution				

Product Information

Item Code (Source) NDC:0338-0324

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
THROMBIN ALFA (UNII: SCK81AMR7R) (THROMBIN ALFA - UNII:SCK81AMR7R)	THROMBIN ALFA	1000 [iU] in 1 mL

Inactive Ingredients

Ingredient Name	Strength
MANNITOL (UNII: 3OWL53L36A)	
SUCROSE (UNII: C151H8M554)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
HISTIDINE (UNII: 4QD397987E)	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0324-01	5 mL in 1 VIAL; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125248	01/29/2008	

Part 2 of 2

SODIUM CHLORIDE

normal saline solution

Product Information

Route of Administration TOPICAL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		5 mL in 1 SYRINGE; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125248	01/29/2008	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125248	01/29/2008	

RECOTHROM

thrombin topical recombinant kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-0326
---------------------	-------------------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0326-01	1 in 1 KIT; Type 1: Convenience Kit of Co-Package	06/09/2008	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 VIAL	20 mL
Part 2	1 SYRINGE	20 mL

Part 1 of 2

RECOTHROM

thrombin topical recombinant powder, for solution

Product Information**Item Code (Source)** NDC:0338-0328**Route of Administration** TOPICAL**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
THROMBIN ALFA (UNII: SCK81AMR7R) (THROMBIN ALFA - UNII:SCK81AMR7R)	THROMBIN ALFA	1000 [iU] in 1 mL

Inactive Ingredients

Ingredient Name	Strength
MANNITOL (UNII: 3OWL53L36A)	
SUCROSE (UNII: C151H8M554)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
HISTIDINE (UNII: 4QD397987E)	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0328-01	20 mL in 1 VIAL; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125248	06/09/2008	

Part 2 of 2**SODIUM CHLORIDE**

sodium chloride solution

Product Information**Route of Administration** TOPICAL**Inactive Ingredients**

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		20 mL in 1 SYRINGE; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125248	06/09/2008	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125248	06/09/2008	

RECOTHROM

thrombin topical recombinant kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-0330
---------------------	-------------------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0330-01	1 in 1 KIT; Type 1: Convenience Kit of Co-Package	06/09/2008	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 VIAL	20 mL
Part 2	1 VIAL	20 mL

Part 1 of 2

RECOTHROM

thrombin topical recombinant powder, for solution

Product Information

Item Code (Source) NDC:0338-0328

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
THROMBIN ALFA (UNII: SCK81AMR7R) (THROMBIN ALFA - UNII:SCK81AMR7R)	THROMBIN ALFA	1000 [iU] in 1 mL

Inactive Ingredients

Ingredient Name	Strength
MANNITOL (UNII: 3OWL53L36A)	
SUCROSE (UNII: C151H8M554)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
HISTIDINE (UNII: 4QD397987E)	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0328-01	20 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125248	06/09/2008	

Part 2 of 2

SODIUM CHLORIDE

sodium chloride solution

Product Information

Route of Administration TOPICAL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		20 mL in 1 VIAL; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125248	06/09/2008	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125248	06/09/2008	

Labeler - Baxter Healthcare Corporation (005083209)

Revised: 1/2024

Baxter Healthcare Corporation