

IBUPROFEN PM- ibuprofen, diphenhydramine hcl capsule, liquid filled Walgreens

Drug Facts

Active ingredient (in each capsule)

Diphenhydramine hydrochloride USP 25 mg

**Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)**

***non-steroidal anti-inflammatory drug**

Purpose

Nighttime sleep-aid

Pain Reliever

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other antihistamines
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - leg swelling

- slurred speech
- weakness in one part or side of body
- pain gets worse or last more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- adults and children 12 years and over: take 2 capsules at bedtime
- do not take more than 2 capsules in 24 hours

Other information

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F)
- protect from light
- Organic Impurities Test is pending
- USP Dissolution Test is pending
- swallow whole; do not crush, chew, or dissolve

Inactive ingredients

D&C red #33, FD&C blue #1, gelatin, hypromellose, isopropyl alcohol, polyethylene glycol, propylene glycol, potassium hydroxide, povidone, purified water, sorbitan, sorbitol, titanium dioxide

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal display panel

Compare to the active ingredients in Advil® PM††

Ibuprofen **PM**

IBUPROFEN, 200 mg / PAIN RELIEVER (NSAID)

DIPHENHYDRAMINE HCl, 25 mg / NIGHTTIME SLEEP AID

NIGHTTIME SOFTGELS

SOFTGELS**

(**LIQUID-FILLED CAPSULES)

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor Advil® PM.

TAMPER EVIDENT: DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

Product Packaging

Lot No.:
Exp. Date:

PLD-A690A FC007036



DISTRIBUTED BY: WALGREEN CO.
200 WILLMOT RD., DERRFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2020 Walgreen Co.
ITEM 621604 W10024-0720-L

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey
†† This product is not manufactured or distributed by
Pfizer Consumer Healthcare, distributor of Advil® PM.

Drug Facts
Active ingredients
(in each capsule)

Diphenhydramine hydrochloride USP 25 mg.....Nighttime sleep-aid
Sulbuzolid ibuprofen equal to
200 mg ibuprofen USP (NSAID)*.....Pain reliever
(present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Uses
■ for relief of occasional sleeplessness when associated with minor aches and pains
■ helps you fall asleep and stay asleep

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ unless you have time for a full night's sleep
■ in children under 12 years of age
■ right before or after heart surgery

Drug Facts (continued)
■ with any other product containing diphenhydramine, even one used on skin
■ if you have sleeplessness without pain

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
■ you are taking a diuretic
■ you have a breathing problem such as emphysema or chronic bronchitis

■ you have glaucoma
■ you have trouble urinating due to an enlarged prostate gland
Ask a doctor or pharmacist before use if you are
■ taking sedatives or tranquilizers, or any other sleep-aid
■ under a doctor's care for any continuing medical illness
■ taking any other antihistamines
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
■ taking any other drug

When using this product
■ drowsiness will occur
■ avoid alcoholic drinks
■ do not drive a motor vehicle or operate machinery
■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding
■ feel faint ■ vomit blood
■ have bloody or black stools
■ you have stomach pain that does not get better
■ you have symptoms of heart problems or stroke
■ chest pain ■ trouble breathing
■ weakness in one part or side of body
■ leg swelling ■ slurred speech
■ pain gets worse or lasts more than 10 days

Drug Facts (continued)
■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ do not take more than directed
■ adults and children 12 years and over: take 2 capsules at bedtime
■ do not take more than 2 capsules in 24 hours

Other information
■ each capsule contains: potassium 20 mg
■ read all warnings and directions before use. Keep carton.
■ store at 20° to 25°C (68° to 77°F)
■ avoid excessive heat above 40°C (104°F)
■ protect from light
■ Organic Impurities Test is pending
■ USP Dissolution Test is pending
■ swallow whole; do not crush, chew, or dissolve

Inactive ingredients D&C Red No. 33, gelatin, hypromellose, isopropyl alcohol, Neelcent PB&C Blue No. 1, polyethylene glycol, propylene glycol, potassium hydroxide, polydioxane, purified water, sorbitol sorbitan solution, titanium dioxide

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST



how2recycle.info
Empire & Republic Cards
TAMPER EVIDENT: DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION



Compare to the active ingredients in Advil® PM††

Ibuprofen PM

IBUPROFEN CAPSULES 200 mg / PAIN RELIEVER (NSAID)
DIPHENHYDRAMINE HCl 25 mg / NIGHTTIME SLEEP AID

NDC 0363-0250-40

NIGHTTIME SOFTGELS

40 SOFTGELS**
(**LIQUID-FILLED CAPSULES)



WALGREENS Ibuprofen PM Softgels

IBUPROFEN PM

ibuprofen, diphenhydramine hcl capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0250
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ALCOHOL (UNII: 3K9958V90M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POVIDONE (UNII: FZ989GH94E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	DHI25
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0250-40	1 in 1 BOX	01/01/2021	
1		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210676	01/01/2021	

Labeler - Walgreens (008965063)

Revised: 11/2022

Walgreens