

**RITEAID NIGHTTIME SLEEP AID- diphenhydramine hydrochloride and acetaminophen liquid**  
**RITE AID CORPORATION**

-----  
**RITEAID Nighttime Sleep-aid Pain Reliever**

***Drug Facts***

**Active ingredients (in each 30 mL)**

Acetaminophen 1000 mg

Diphenhydramine HCl 50 mg

**Purpose**

Pain reliever

Nighttime sleep-aid

**Uses**

- for the temporary relief of occasional minor aches and pains with accompanying sleeplessness.

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs., which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age

- with any other product containing diphenhydramine, even one used on skin.

### **Ask a doctor before use if you have**

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- avoid alcoholic beverages
- drowsiness will occur
- do not drive a motor vehicle or operate machinery

### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- Sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- take only one dose (30 mL) per day (24 hours)
- Only use dose cup provided

adults & children 12 yrs & over	30 mL at bedtime
children under 12 yrs	do not use

### **Other information**

- **each 30 mL contains:** sodium 10 mg
- store at room temperature
- very low sodium

## **Inactive ingredients**

citric acid, flavors, FD&C Blue No. 1, FD&C Red No. 40, glycerin, potassium citrate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose.

## **Questions or comments?**

**1-866-467-2748**

## **PRINCIPAL DISPLAY PANEL - 354 ml bottle**

Compare to the active ingredient in Vicks® ZzzQuil™ Night Pain Sleep-Aid\*

NDC# 11822-8791-1

## **Nighttime Sleep-Aid**

### **Pain Reliever**

Diphenhydramine HCl

### **Acetaminophen**

- Fall Asleep Fast
- Max Strength Pain Reliever
- Non-Habit Forming

### **Berry Flavor**

Naturally and Artificially Flavored

No Added Alcohol

### **12 FL. OZ. (354 ml)**

\*This product is not manufactured or distributed by Procter & Gamble, the distributor of Vicks® ZzzQuil™ Night Pain Sleep-Aid.

### **Distributed by:**

7.875" Width

3.5" Height

**Drug Facts** (continued)

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only one dose (30 mL) per day (24 hours)
- only use the dose cup provided

adults & children 12 yrs & over	30 mL at bedtime
children under 12 yrs	do not use

**Other information**

- each 30 mL contains: sodium 10 mg
- store at room temperature
- very low sodium

**Inactive ingredients** citric acid, flavors, FD&C Blue No. 1, FD&C Red No. 40, glycerin, potassium citrate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

**Questions or comments?** 1-866-467-2748

\*This product is not manufactured or distributed by Procter & Gamble, the distributor of Vicks® ZzzQuil™ Night Pain Sleep-Aid.

**DISTRIBUTED BY:** RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011 [www.riteaid.com](http://www.riteaid.com)  
60012RITLR

Compare to the active ingredients in  
**Vicks® ZzzQuil™ Night Pain Sleep-Aid\***

NDC 11822-8791-1

**NIGHTTIME SLEEP-AID PAIN RELIEVER**

**DIPHENHYDRAMINE HCl**  
50 mg per 30 mL

**ACETAMINOPHEN 1000 mg per 30 mL**

max strength pain reliever  
fall asleep fast  
non-habit forming  
no added alcohol

**BERRY FLAVOR**  
NATURALLY AND ARTIFICIALLY FLAVORED

12 FL OZ (354 mL)

**Drug Facts** **TAMPER EVIDENT:** Do not use if printed shrink band is missing or broken.

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 1000 mg	Pain reliever
Diphenhydramine HCl 50 mg	Nighttime sleep-aid

**Uses** ■ for the temporary relief of occasional minor aches and pains with accompanying sleeplessness.

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product ■ taken with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- for children under 12 years of age
- with any other product containing diphenhydramine, even one used on skin

**Ask a doctor before use if you have** ■ liver disease ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are** ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

**When using this product** ■ avoid alcoholic beverages ■ drowsiness will occur ■ do not drive a motor vehicle or operate machinery

**Stop use and ask a doctor if** ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ Sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness. These could be signs of a serious condition. ▶

LOT: \_\_\_\_\_

EXP: \_\_\_\_\_

0 11822 12989 3

## RITEAID NIGHTTIME SLEEP AID

diphenhydramine hydrochloride and acetaminophen liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-8791
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	1000 mg in 30 mL
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	50 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POTASSIUM CITRATE</b> (UNII: EE90ONI6FF)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

## Product Characteristics

<b>Color</b>	RED	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-8791-1	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2021	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M010	06/01/2021	

**Labeler** - RITE AID CORPORATION (014578892)

Revised: 11/2023

RITE AID CORPORATION