

X RAY PAIN RELIEVER- acetaminophen, aspirin, and caffeine tablet

Genomma Lab USA

X Ray® Pain Reliever Tablets

Drug Facts

Active ingredients (in each tablet)	Purposes
Acetaminophen 250mg	Pain reliever
Aspirin 250mg (NSAID*)	Pain reliever
Caffeine 65mg	Pain reliever aid

* nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to

- headache
- a cold
- arthritis
- muscular aches
- toothache
- premenstrual and menstrual cramps

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)

- shock

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning

This product contains an NSAID which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non prescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning

The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug, or under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice the signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose.
- **adults and children 12 years and over:** take 2 tablets every 6 hours; not more than 8 tablets in 24 hours
- **children under 12 years:** ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- keep carton for full product information
- do not use if imprinted safety seal under cap is broken or missing
- you may report serious side effects to 1-888-952-0050

Inactive ingredients

corn starch, croscarmellose sodium, crospovidone, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, silicon dioxide, stearic acid, titanium dioxide

Distributed by:
Genomma Lab USA Inc., Houston, TX 77098

PRINCIPAL DISPLAY PANEL - 24 Tablet Bottle Carton

Acetaminophen / Aspirin (NSAID) and Caffeine
Pain Reliever / Pain reliever aid

X RAY®
PAIN RELIEVER

✓ EXTRA STRENGTH

RELIEVES:

- ARTHRITIS AND JOINT PAIN
- HEADACHE
- ACHES AND PAIN DUE TO A COLD
- MUSCULAR ACHES
- MENSTRUAL CRAMP PAIN

24 | TABLETS

Acetaminophen / Aspirin (NSAID) and Caffeine
Pain Reliever / Pain reliever aid
Acetaminofén / Aspirina (ANE) y Cafeína
Analgésico / Alivia el dolor

X RAY
PAIN RELIEVER

Acetaminophen / Aspirin (NSAID) and Caffeine
 Pain Reliever / Pain reliever aid
 Acetaminofén / Aspirina (ANE) y Cafeína
 Analgésico / Alivia el dolor

X RAY
PAIN RELIEVER

✓ **EXTRA STRENGTH**
 ✓ **EXTRA FUERTE**

RELIEVES / ALIVIA:

- ARTHRITIS AND JOINT PAIN / DOLOR ARTICULAR Y DE ARTRITIS
- HEADACHE / CABEZA
- ACHE AND PAIN DUE TO A COLD / MOLESTIA Y DOLOR POR RESFRIADO COMÚN
- MUSCULAR ACHE / DOLORES MUSCULARES
- MENSTRUAL CRAMP PAIN / DOLORES MENSTRUALES

24 TABLETS / TABLETAS

Drug Facts

Active Ingredients (in each tablet)
 Acetaminophen 250mg.....Pain reliever
 Aspirin 250mg (NSAID).....Pain reliever
 Caffeine 65mg.....Nonsteroidal anti-inflammatory drug

Uses Temporarily relieves minor aches and pains due to:
 ● muscle aches ● cold ● headache ● cold ● influenza ● muscle aches ● headaches ● menstrual and menstrual cramps

Warnings
 Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. Worsening of liver disease or stomach problems may occur. Consult a doctor because these conditions may be worsened by this product. Aspirin should not be given to children or teenagers because it may cause severe skin reactions. Symptoms may include:
 ● skin redness ● blisters ● rash ● skin reaction occurs, stop use and seek medical help right away.
 ● Allergy alert: Aspirin may cause a severe allergic reaction which may include:
 ● hives ● facial swelling ● asthma (wheezing) ● shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 8 tablets in 24 hours or the maximum daily amount with other drugs containing acetaminophen, or if you drink alcohol every day while using this product. This product contains an NSAID which may cause severe stomach bleeding.

Drug Facts (continued)

The chance is higher if you:
 ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinning (anticoagulant) or steroid drug ● take other drugs containing aspirin (Aspirin, Ibuprofen, naproxen) ● take other drugs containing acetaminophen ● take other drugs containing caffeine ● take other drugs containing caffeine containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat

Do not use if you have ever had an allergic reaction to acetaminophen, aspirin or other pain reliever/fever reducer ● with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if ● you have liver disease ● stomach bleeding warning applies to you ● you have a history of stomach problems, such as heartburn ● you have high blood pressure, heart disease, liver arthritis, or kidney disease ● you are taking a diuretic ● you have asthma

Ask a doctor or pharmacist before use if you are ● taking a prescription drug for diabetes, gout, or arthritis ● taking any other drug, or under a doctor's care for any serious condition

Stop use and ask a doctor if ● an allergic reaction occurs. Seek medical help right away ● you experience any of the following signs of stomach bleeding ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ● ringing in the ears or loss of hearing occurs ● palms/soles feel red or swollen ● pain gets worse or lasts for more than 10 days ● fever gets worse or lasts for more than 3 days ● any new symptoms occur. These could be signs of a serious condition

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during or after birth. Avoid use of aspirin unless directed by a doctor. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice the signs or symptoms.

Directions
 ● do not use more than directed ● drink a full glass of water with each dose.
 ● adults and children 12 years and over: take 2 tablets every 6 hours, not more than 8 tablets in 24 hours ● children under 12 years: ask a doctor

Other information
 ● store at 20°-25°C (68°-77°F) ● keep carton for full product information ● do not use if imprinted safety seal under cap is broken or missing ● you may report serious side effects to 1-888-952-0150

Inactive ingredients
 hydroxypropylcellulose, croscarmellose sodium, croscopolone, hydroxypropylcellulose, polyethylene glycol, povidone, pregelatinized starch, silicon dioxide, stearic acid, titanium dioxide

Genomma Lab

Lot/Lote:
Exp:

Made in USA with foreign and domestic ingredients.
 Distribuido por: / Distributed by:
 Genomma Lab USA Inc., Houston, TX 77098

Información del Medicamento

Ingrediente Activo (en cada tableta)	Propósito
Acetaminofén 250 mg.....	Analgésico
Aspirina 250 mg (ANE).....	Analgésico
Cafeína 65 mg.....	Auxiliar del Analgésico

* medicamento antiinflamatorio no esteroideo

Usos alivia temporalmente dolencias y dolores menores debidos a:
 ● cefalea ● un resfriado ● artritis ● dolores musculares ● dolor de muelas ● cólicos premenstruales y menstruales

Advertencias
 Síndrome de Reye: No deberá administrar este producto a niños y adolescentes que tienen o se están recuperando de varicela o sifomas similares a la gripa. Si al administrar este producto ocurren cambios en el comportamiento con náusea o vómito, consulte a un médico. Debido a que este síndrome se reduce por un alto consumo del Sistema de Dosis...

Incluye información en español

¡Atención! Consulte a su médico antes de que cause síntomas que le sea difícil respirar o que se hinche la cara, una enfermedad rara pero grave.
Alerta de alergia: El Paracetamol puede causar reacciones graves en la piel incluyendo: ■ enrojecimiento ■ ampollas ■ salpullido. Si se producen reacciones de la piel, deje de usar y consulte a un médico inmediatamente. La aspirina puede ocasionar una reacción alérgica severa, la cual incluye: urticaria, hinchazón facial, asma (resaca), o choque.

Información del Medicamento (continuación)

Advertencia hepática: Este producto contiene paracetamol. Puede presentarse daño hepático severo si usted toma:
 ■ más de 8 tabletas en 24 horas, la cual es la cantidad diaria máxima ■ junto con otros medicamentos que contengan paracetamol ■ 3 o más bebidas alcohólicas cada día al mismo tiempo que usa este producto ■ tomar más o por un tiempo más prolongado de lo indicado.
Advertencia de hemorragia estomacal: Este producto contiene un medicamento antiinflamatorio no esteroideo (AINE), el cual puede ocasionar hemorragia estomacal severa. Las probabilidades son mayores si usted: ■ tiene 60 años o más ■ ha tenido úlceras estomacales o problemas de hemorragia ■ toma un adelgazante sanguíneo (anticoagulante) o un medicamento esteroideo ■ toma otros medicamentos que contengan AINEs de prescripción o no de prescripción (aspirina, ibuprofeno, naproxeno, u otros) ■ toma 3 o más bebidas alcohólicas cada día al mismo tiempo que utiliza este producto ■ toma más o por un tiempo más prolongado de lo indicado.
Advertencia de la Cafeína: La dosis recomendada de este producto contiene aproximadamente tanta cafeína como un taza de café. Limite el uso de medicamentos, alimentos, o bebidas que contengan cafeína mientras está tomando este producto debido a que un exceso de cafeína puede ocasionar nerviosismo, irritabilidad, insomnio, y ocasionalmente taquicardia.
No utilizar ■ Si presenta reacciones alérgicas a paracetamol, aspirina, o cualquier analgésico / Antipirético.
 ■ Con otro medicamento que contenga paracetamol, (de prescripción o no de prescripción). Si no está seguro sobre si el medicamento contiene paracetamol, consulte a su médico o farmacéutico.
 Consulte a su médico antes de usarlo si ■ padece enfermedad hepática ■ la advertencia sobre hemorragia estomacal

Información del Medicamento (continuación)

se aplica a usted ■ tiene una historia de problemas estomacales, tales como acidez ■ Padece de hipertensión, enfermedad cardíaca, cirrosis hepática, o enfermedad renal ■ asma ■ está tomando un diurético
 Consulte a su médico o farmacéutico antes de usarlo si usted está tomando ■ cualquier otro medicamento que contenga un AINE (de prescripción o no de prescripción) ■ un adelgazante sanguíneo (anticoagulante) o medicamento esteroideo ■ un medicamento de prescripción para diabetes, gota o artritis ■ cualquier otro medicamento, o está bajo el cuidado del médico por alguna condición grave.
 Suspenda su uso y consulte a su médico si se presenta una reacción alérgica. Busque ayuda médica de inmediato.
 ■ experimenta cualquiera de los siguientes signos de hemorragia estomacal: se siente mareado, vomita sangre, tiene heces con sangre, tiene dolor estomacal que no mejora. ■ zumbidos en los oídos o presenta pérdida auditiva ■ el área del dolor está enrojecida o hinchada ■ el dolor empeora o persiste más de 10 días ■ la fiebre empeora o persiste más de 3 días ■ aparece cualquier síntoma nuevo.
 En caso de embarazo o lactancia, consulte a un profesional médico antes de usarlo. Es especialmente importante no usar la aspirina durante los últimos 3 meses del embarazo a menos que sea definitivamente indicada de tomarla por parte de su médico debido a que puede ocasionar problemas en el feto o complicaciones durante el parto.
 Manténgase fuera del alcance de los niños.
 En caso de sobredosis, contacte a su médico o al Centro de Control de Envenenamiento (1-800-222-1222) inmediatamente. La atención médica pronta es crítica para adultos al igual que para niños incluso si usted no nota ningún signo o síntoma.

Información del Medicamento (continuación)

Indicaciones ■ no usar más de lo indicado ■ con cada dosis tome un vaso lleno de agua
 ■ adultos y niños de 12 años y mayores: tomar 2 tabletas cada 6 horas; no tomar más de 8 tabletas en 24 horas
 ■ niños menores de 12 años consulte a su médico
Información adicional
 ■ almacenar a temperatura ambiente entre 20-25 °C (68-77 °F)
 ■ cierre tapa seguramente
 ■ lea toda la información de producto antes de usar Mantenga cajilla para obtener información del medicamento.
Ingredientes inactivos
 almidón de maíz, croscarmelosa sódica, crospovidón a, hipromelosa, microcristalina celulosa, polietilenglicol, povidona, amilón pregelatinizado, dióxido de silicio, estearato ácido, dióxido de titanio
¿Preguntas o comentarios? Número gratuito 1-877-994-3666

X RAY PAIN RELIEVER

acetaminophen, aspirin, and caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50066-455
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
acetaminophen (UNII: 362O9ITL9D) (acetaminophen - UNII:362O9ITL9D)	acetaminophen	250 mg
aspirin (UNII: R16CO5Y76E) (aspirin - UNII:R16CO5Y76E)	aspirin	250 mg
caffeine (UNII: 3G6A5W338E) (caffeine - UNII:3G6A5W338E)	caffeine	65 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE 112 (UNII: X7XJ6RM9Q2)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
silicon dioxide (UNII: ETJ7Z6XBU4)	
stearic acid (UNII: 4ELV7Z65AP)	
talc (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	AZ;285
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50066-455-24	1 in 1 CARTON	12/22/2023	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M013	12/22/2023	

Labeler - Genomma Lab USA (832323534)

Revised: 12/2023

Genomma Lab USA