

**MINOXIDIL (FOR MEN)- minoxidil aerosol, foam**  
**TARGET Corporation**

-----  
**Minoxidil**  
**5% (For Men)**

***Drug Facts***

**Active ingredient**

Minoxidil 5% w/w (without propellant)

**Purpose**

Hair regrowth treatment for men

**Use**

to regrow hair on the top of the scalp (vertex only, see pictures on side of this carton)

**Warnings**

**For external use only.**

**For use by men only.**

**Extremely Flammable:**Avoid fire, flame, or smoking during and immediately following application.

**Do not use if**

- you are a woman
- your amount of hair loss is different than that shown on side of this carton or your hair loss is on the front of the scalp. 5% minoxidil topical foam is not intended for frontal baldness or receding hairline.
- you have no family history of hair loss
- your hair loss is sudden and/or patchy
- you do not know the reason for your hair loss
- you are under 18 years of age. Do not use on babies and children.
- your scalp is red, inflamed, infected, irritated, or painful
- you use other medicines on the scalp

**Ask a doctor before use if you have**heart disease

**When using this product**

- do not apply on other parts of the body
- avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
- some people have experienced changes in hair color and/or texture

- it takes time to regrow hair. Results may occur at 2 months with twice a day usage. For some men, you may need to use this product for at least 4 months before you see results.
- the amount of hair regrowth is different for each person. This product will not work for all men.

**Stop use and ask a doctor if**

- chest pain, rapid heartbeat, faintness, or dizziness occurs
- sudden, unexplained weight gain occurs
- your hands or feet swell
- scalp irritation or redness occurs
- unwanted facial hair growth occurs
- you do not see hair regrowth in 4 months

**May be harmful if used when pregnant or breast-feeding.**

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- apply half a capful 2 times a day to the scalp in the hair loss area
- massage into scalp with fingers, then wash hands well
- see enclosed leaflet for complete directions on how to use
- using more or more often will not improve results
- continued use is necessary to increase and keep your hair regrowth or hair loss will begin again

**Other information**

- hair growth has been shown in a clinical study of men (mostly white) aged 18-49 years who used it for 4 months
- see hair loss pictures on side of this carton
- before use, read all information on carton and enclosed leaflet
- keep the carton. It contains important information.
- store at controlled room temperature 20° to 25°C (68° to 77°F)
- contents under pressure. Do not puncture or incinerate container.
- do not expose to heat or store at temperatures above 120°F (49°C).

**Inactive ingredients**

butylated hydroxytoluene, cetyl alcohol, citric acid anhydrous, dehydrated alcohol, isobutane, lactic acid, n-butane, polysorbate 60, propane, purified water, stearyl alcohol

**Questions?**

- Call **1-866-923-4914**

Distributed by Target Corporation  
Minneapolis, MN 55403

**PRINCIPAL DISPLAY PANEL - 60 g Can Carton**

hair regrowth  
treatment  
for men

minoxidil topical aerosol, 5% (for men)

reactivates hair follicles to  
stimulate regrowth

easy-to-use foam  
unscented

not for use by women

up&up <sup>TM</sup>

FOAM

3  
MONTH  
SUPPLY

3 MONTH SUPPLY  
THREE - 60 g (2.11 OZ) CANS



## MINOXIDIL (FOR MEN)

minoxidil aerosol, foam

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-157
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MINOXIDIL (UNII: 5965120SH1) (MINOXIDIL - UNII:5965120SH1)	MINOXIDIL	50 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BUTANE (UNII: 6LV4FOR43R)	

<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>CETYL ALCOHOL</b> (UNII: 936JST6JCN)	
<b>ISOBUTANE</b> (UNII: BXR49TP611)	
<b>LACTIC ACID, UNSPECIFIED FORM</b> (UNII: 33X04XA5AT)	
<b>POLYSORBATE 60</b> (UNII: CAL22UVI4M)	
<b>PROPANE</b> (UNII: T75W9911L6)	
<b>STEARYL ALCOHOL</b> (UNII: 2KR89I4H1Y)	
<b>WATER</b> (UNII: 059QF0KO0R)	

### Product Characteristics

<b>Color</b>	white (white to off-white)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-157-03	3 in 1 CARTON	08/16/2022	
1	NDC:11673-157-01	60 g in 1 CAN; Type 0: Not a Combination Product		
2	NDC:11673-157-09	3 in 1 CARTON	08/16/2022	
2	NDC:11673-157-01	60 g in 1 CAN; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209074	08/16/2022	

**Labeler** - TARGET Corporation (006961700)

### Establishment

Name	Address	ID/FEI	Business Operations
Taro Pharmaceutical Industries Ltd.		600072078	manufacture(11673-157)

Revised: 5/2024

TARGET Corporation