

**MUCUS RELIEF DM MAXIMUM STRENGTH- dextromethorphan hbr,
guaifenesin liquid
Rite Aid Corporation**

Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Purpose

Cough suppressant

Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of cough
 - the impulse to cough to help you get to sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm(mucus)

When using this product,

- **do not use more than directed**

Stop use and ask a doctor if

cough lasts more than 7 days, comes back, or occurs with fever, rash or persistent

headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age : do not use

Other information

- **each 20 mL contains:** sodium 20 mg
- store between 20-25°C (68-77°F). do not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C red 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® DM Max*

MAXIMUM STRENGTH

MUCUS RELIEF

DM MAX

MULTI-SYMPTOM RELIEF

COUGH SUPPRESSANT EXPECTORANT

DEXTROMETHORPHAN HBr 20 mg

GUAIFENESIN 400 mg

Controls cough

Relieves chest congestion

Thins & loosen mucus

For ages 12 & over

4-hour dosing

For ages 12 and over

FL OZ(mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® DM Max.

DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE,

CAMP HILL, PA 17011

www.riteaid.com

Package Label

Compare to the active ingredients in
Maximum Strength Mucinex®
Fast-Max® DM Max*

NDC 11822-7224-6

MAXIMUM STRENGTH

MUCUS RELIEF DM MAX

MULTI-SYMPTOM RELIEF

COUGH SUPPRESSANT
EXPECTORANT

**DEXTROMETHORPHAN HBr 20 mg
GUAIFENESIN 400 mg**

Controls cough
Relieves chest congestion
Thins & loosens mucus
For ages 12 & over
4-hour dosing



PLD-B409A LB006733

6 FL OZ
(177 mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY
SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

*This product is not manufactured or distributed by
Reckitt Benckiser, distributor of Maximum Strength
Mucinex® Fast-Max® DM Max.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

PLD-A409A LB007075



DISTRIBUTED BY:
RITE AID, 30 HUNTER LANE,
CAMP HILL, PA 17011
www.riteaid.com

SATISFACTION GUARANTEE

If you're not satisfied, we'll
happily refund your money.

PEEL CORNER FOR DRUG FACTS ▶

Drug Facts

Active ingredients Purposes (in each 20 mL)

Dextromethorphan HBr 20 mg.....
.....Cough suppressant
Guaifenesin 400 mg.....Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Drug Facts (continued)

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product, do not use more than directed.

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. ▶

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

Other information

- **each 20 mL contains:** sodium 20 mg
- store between 20-25°C (68-77°F). Do not refrigerate. ▶

Drug Facts (continued)

Inactive ingredients

citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday
9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS ▶

RITE AID Maximum Strength Mucus Relief DM

MUCUS RELIEF DM MAXIMUM STRENGTH

dextromethorphan hbr, guaifenesin liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:11822-7224 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------------|--------------------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 20 mg in 20 mL |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 400 mg in 20 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| EDETATE SODIUM (UNII: MP1J8420LU) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| PROPYL GALLATE (UNII: 8D4SNN7V92) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K) | |
| WATER (UNII: 059QF0KO0R) | |
| SORBITOL (UNII: 506T60A25R) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:11822-7224-6 | 117 mL in 1 BOTTLE; Type 0: Not a Combination Product | 11/26/2021 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 11/26/2021 | |

Labeler - Rite Aid Corporation (014578892)

