

**DOLLAR GENERAL PAIN RELIEVING PATCH WITH LIDOCAINE- lidocaine patch
DOLGENCORP, LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Dollar General Pain Relieving Patch with Lidocaine

Active Ingredients

Lidocaine 4% Topical Anesthetic

Purpose

Temporarily relieves minor pain.

Dosage and Administration

Directions Adults and children over 12 years:

- clean and dry affected area
- remove backing from patch by firmly grasping both ends and gently pulling until backing separates in middle
- carefully remove smaller portion of backing from patch starting at corner
- apply sticky side of patch to affected area
- use one patch for up to 12 hours. **Children under 12 years of age:** consult a physician.

Warnings

For external use only.

Indications and Usage

Uses: Temporarily relieves minor pains.

When using this product

- use only as directed
- read and follow all directions and warnings on this carton
- do not allow contact with the eyes
- do not use at the same time as other topical analgesics

- do not bandage tightly or apply local heat (such as heating pads) to the area of use
- do not microwave
- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and consult a doctor

- condition worsens
- redness is present
- irritation develops
- symptoms persist for more than 7 days or clear up and occur again within a few days
- you experience signs of skin injury, such as pain, swelling or blistering where the product was applied.

If pregnant or breastfeeding, ask a health professional before use.

Do not Use

- more than one patch on your body at a time
- on cut, irritated or swollen skin
- on puncture wounds
- for more than one week without consulting a doctor
- if you are allergic to any active or inactive ingredients
- if pouch is damaged or opened.

Keep out of reach of children and pets.

If swallowed, get medical help or contact a Poison Control Center right away.

Other Safety Information

Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

Inactive Ingredients

Aluminum Glycinate, carboxymethylcellulose sodium, Glycerin, iodopropynyl butylcarbamate, Kaolin, petrolatum, Phenoxyethanol, polyacrylic acid, polysorbate 80, povidone, Propylene Glycol, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Water, 3-(2-ethylhexyloxy)propane-1,2-diol

DIST. BY

Dolgenercorp, LLC
 100 Mission Ridge
 Goodlettsville, TN

Made in China



HOW TO APPLY



Drug Facts	Purpose
Active ingredient Lidocaine 4%	Topical anesthetic
Use: Temporarily relieves minor pain.	
Warnings For external use only. Do not use more than one patch on your body at a time. On cut, irritated or swollen skin. On puncture wounds. For more than one week without consulting a doctor if you are allergic to any acids or inactive ingredients. If pouch is damaged or opened. When using the product: Use only as directed. Read and follow all directions and warnings on this carton. Do not allow contact with the eyes. Do not use at the same time as other topical anesthetics. Do not bandage tightly or apply heat (such as heating pads) to the area of use. Do not microwave. Dispose of used patch in manner that avoids release of product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch. Stop use and consult a doctor if: Condition worsens; if redness is present; if irritation develops; if symptoms persist for more than 7 days or clear up and occur again within a few days; if you experience signs of skin injury such as pain, swelling or blistering where the product was applied. If pregnant or breastfeeding, ask a health professional for use. Keep out of reach of children and pets. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
Directions Adults and children 12 years of age and over: Clean and dry affected area. Carefully remove backing from patch starting at a corner. Apply sticky side of patch to affected area. Use one patch for up to 12 hours. Discard patch after single use. Children under 12 years of age: consult a physician.	
Other information Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.	
Inactive ingredients aluminum glycolate, carboxymethylcellulose sodium, glycerol, hydroxypropyl butylcarbamate, isobutylparaffin, dihydroxyethyl polyacrylate acid, polyacrylate 90, polyvinyl, polyethylene glycol, sodium polyacrylate, tartaric acid, titanium dioxide, water, 3-(2-ethylhexyloxy)propane-1,2-diol	

*The product is not bioequivalent to or distributed by Chullin, Inc., owner of the registered trademark Aspercreme®.

DISTRIBUTED BY OLD EAST MAIN CO.
 100 MISSION RIDGE
 GOODLETTSVILLE, TN 37072



Child-Resistant Packaging
 Model 77-4111-1P
 82-27032 01/22
 V220046
 LOT
 EXP

HOW TO APPLY



Drug Facts
Active ingredient.....Purpose
 Lidocaine 4%.....Topical anesthetic
Use Temporarily relieves minor pain.
Warnings
 For external use only
 Do not use ■ More than one patch on your body at a time ■ On cut, irritated or swollen skin ■ On puncture wounds ■ For more than one week without consulting a doctor ■ If you are allergic to any active or inactive ingredients ■ If pouch is damaged or opened.
 When using this product ■ Use only as directed ■ Read and follow all directions and warnings on this carton ■ Do not allow contact with the eyes ■ Do not use at the same time as other topical anesthetics ■ Do not bandage tightly or apply local heat (such as heating pads) to the area of use ■ Do not microwave ■ Dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.
 Stop use and consult a doctor if ■ Condition worsens ■ Redness is present ■ Irritation develops ■ Symptoms persist for more than 7 days or clear up and occur again within a few days ■ You experience signs of skin injury, such as pain, swelling or blistering where the product was applied.
 If pregnant or breastfeeding, ask a health professional before use.
 Keep out of reach of children and pets. If swallowed, get medical help or contact a Poison Control Center 1-800-322-1222 right away.
Directions
 Adults and children 12 years of age and over: Clean and dry affected area. Carefully remove backing from patch, starting at a corner. Apply sticky side of patch to affected area. Use one patch for up to 12 hours. Discard patch after single use. Children under 12 years of age: consult a physician.
Other information
 Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.
Inactive ingredients
 aluminum glycolate, carbonylmethylcellulose sodium, glycerin, lidopropyl butylcarbamate, laolin, petrolatum, phenoxethanol, polyacrylic acid, polyisobutyl methacrylate, polyisobutylene glycol, sodium polyacrylate, tartaric acid, titanium dioxide, water, 3-(2-ethylhexyloxy)propylene-1,2-diol.

*Solonpas® is a registered trademark of Hombach Pharmaceutical Co. Inc.
 Hombach Pharmaceutical Co. Inc. is not affiliated with "Other General" or this product.
 Model 17-5175-29
 82-77035 02/22
 VCS00046
 AT&T
 MADE IN CHINA

LOT
EXP



Maximum Strength Lidocaine Pain Relief Gel-Patch
 Lidocaine 4% Topical Anesthetic
 NDC 55910-912-02

DG™ health

Compare to Solonpas® Lidocaine Patch active ingredient*

Maximum Strength Lidocaine Pain Relief Gel-Patch
 Lidocaine 4% Topical Anesthetic

- Numbing relief
- Odor free
- No-mess
- Stay-put flexible patch

Up to 12 hours

Neck, Shoulder, Back, Knee & Elbow

2 Large Patches
 3.93 in x 5.51 in (10 cm x 14 cm)

Maximum Strength Lidocaine Pain Relief Gel-Patch
 Lidocaine 4% Topical Anesthetic

DOLLAR GENERAL PAIN RELIEVING PATCH WITH LIDOCAINE

lidocaine patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-912
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
DIHYDROXYALUMINUM AMINOACETATE (UNII: DO250MG0W6)	
GLYCERIN (UNII: PDC6A3C0OX)	
KAOLIN (UNII: 24H4NWX5CO)	
POVIDONE (UNII: FZ989GH94E)	
POLYACRYLIC ACID (300000 MW) (UNII: A8371R0U5J)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
SODIUM POLYACRYLATE (8000 MW) (UNII: 285CYO341L)	
TARTARIC ACID (UNII: W4888I119H)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	
PETROLATUM (UNII: 4T6H12BN9U)	
IODOPROPYNYL BUTYLCARBAMATE (UNII: 603P14DHEB)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	

Product Characteristics

Color		Score	
Shape	RECTANGLE	Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-912-01	1 in 1 CARTON	10/01/2017	
1		9 g in 1 PATCH; Type 0: Not a Combination Product		
2	NDC:55910-912-02	2 in 1 CARTON	09/01/2018	
2		9 g in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	10/01/2017	

Labeler - DOLGENCORP, LLC (068331990)

Establishment

Name	Address	ID/FEI	Business Operations
Foshan Aqua Gel Biotech Co.,Ltd.		529128763	manufacture(55910-912)

Revised: 7/2022

DOLGENCORP, LLC