

**NAPROXEN SODIUM- naproxen sodium capsule, liquid filled  
Care One (Retail Business Services, LLC.)**

-----  
**Drug Facts**

**Active ingredient (in each capsule)**

**Naproxen sodium 220 mg**

**(naproxen 200 mg) (NSAID)\***

**\*nonsteroidal anti-inflammatory drug**

**Purposes**

**Pain reliever/fever reducer**

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
  - the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed

**Heart attack and stroke:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

take with food or milk if stomach upset occurs.

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - leg swelling
  - slurred speech
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing

- it feels like the capsule is stuck in your throat

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center(1-800-222-1222) right away.

### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work
- adults and children 12 years and older:
  - take 1 capsule every 8 to 12 hours while symptoms last
  - for the first dose you may take 2 capsules within the first hour
  - do not exceed 2 capsules in any 8- to 12-hour period
  - do not exceed 3 capsules in a 24-hour period
- children under 12 years: ask a doctor

### **Other information**

- **each capsule contains:** sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F). Protect from light.
- read all directions and warnings before use. Keep carton.
- swallow whole; do not crush, chew, or dissolve

### **Inactive ingredients**

FD&C blue #1, gelatin, glycerin, lactic acid, lecithin, light mineral oil, n-butyl alcohol, polyethylene glycol, povidone, propylene glycol, purified water, shellac glaze, sorbitan, sorbitol solution, titanium dioxide, white ink

### **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

Compare to the Active Ingredient in Aleve®†

NAPROXEN SODIUM

Capsules, 220 mg (NSAID)

Pain Reliever; Fever Reducer

Strength to lasts 12 hours

LIQUID GELS\*\*

(\*\*Liquid-Filled Capsules)

†This product is not manufactured or distributed by Bayer HealthCare, LLC distributor of Aleve®.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS  
BROKEN OR MISSING**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT  
INFORMATION.**

DISTRIBUTED BY: ADUSA DISTRIBUTION, LLC

SALISBURY, NC 28147

**Product Label**

Exp. Date:  
Lot No.:

P.L.D.-CS238D  
F00/7851



DISTRIBUTED BY: ADUSA DISTRIBUTION, LLC  
SALISBURY, NC 28147  
For product questions or concerns,  
contact us at 1-833-992-3872  
Quality guaranteed or your money back.

### Drug Facts

#### Active ingredient (in each capsule)

Naproxen sodium 220 mg  
(naproxen 200 mg) (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

#### Purposes

Pain reliever/  
fever reducer

#### Uses

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
- the common cold
- temporarily reduces fever

#### Warnings

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### Drug Facts (continued)

#### Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

#### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product, take with food or milk if stomach upset occurs.

#### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - leg swelling
  - slurred speech
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ do not take more than directed

### Drug Facts (continued)

- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work
- adults and children 12 years and older
  - take 1 capsule every 8 to 12 hours while symptoms last
  - for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- children under 12 years: ask a doctor

### Other information

- each capsule contains: sodium 20 mg
- store between 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F). Protect from light.
- read all directions and warnings before use. Keep carton.
- swallow whole; do not crush, chew, or dissolve

**Inactive ingredients** FD&C blue #1, gelatin, glycerin, lactic acid, lecithin, light mineral oil, n-butyl alcohol, polyethylene glycol, povidone, propylene glycol, purified water, shellac glaze, sorbitol sorbitan solution, titanium dioxide, white ink

### Questions or comments?

Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

This product is not manufactured or distributed by Bayer HealthCare, LLC, distributor of Aleve®.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

CAREone™

NDC 72476-748-80

Compare to the Active Ingredient in Aleve†

NAPROXEN SODIUM

Capsules, 220 mg  
(NSAID)

Actual Size



Pain Reliever/Fever Reducer  
Strength to Last 12 Hours

80 LIQUID GELS\*\*

## NAPROXEN SODIUM

naproxen sodium capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72476-748
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>2-NAPHTHALENEACETIC ACID, 6-METHOXY-.ALPHA.-METHYL-, SODIUM SALT, (S)-</b> (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	2-NAPHTHALENEACETIC ACID, 6-METHOXY-.ALPHA.-METHYL-, SODIUM SALT, (S)-	220 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>LACTIC ACID</b> (UNII: 33X04XA5AT)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	PC19
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC 72476			

1	NDC: 72476-748-80	1 in 1 BOX	04/30/2021	
1		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC: 72476-748-20	1 in 1 BOX	04/30/2021	
2		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA208363	04/30/2021	

**Labeler** - Care One (Retail Business Services, LLC.) (967989935)

Revised: 6/2021

Care One (Retail Business Services, LLC.)