

**HEALTH MART DAYTIME- acetaminophen, dextromethorphan hbr, phenylephrine  
hcl solution**

**Mckesson**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**McKesson Daytime Drug Facts**

**Active ingredients (in each 15 mL)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves common cold/flu symptoms:
- cough due to minor throat and bronchial irritation
- minor aches and pains
- sore throat
- fever
- nasal congestion
- headache

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **When using this product**

do not use more than directed

### **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- each 15 mL contains: sodium 7 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients**

butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

**Questions or comments?**

1-800-719-9260

**Principal Display Panel**

compare to Vicks® DayQuil® Cold & Flu active ingredients

daytime

cold & flu

pain reliever/fever reducer

cough suppressant

nasal decongestant

acetaminophen

phenylephrine HCl

dextromethorphan HBr

alcohol free

multi-symptom relief

non-drowsy - gluten free

8 FL OZ (237 mL)

SEE NEW WARNINGS

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

NDC 62011-0065-2



compare to Vicks® DayQuil®  
Cold & Flu active ingredients\*

# daytime

cold & flu

pain reliever/fever reducer  
cough suppressant  
nasal decongestant

acetaminophen  
phenylephrine HCl  
dextromethorphan HBr  
alcohol free

multi-symptom relief

non-drowsy • gluten free

8 FL OZ (237 mL)

SEE NEW WARNINGS

### Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

### Uses

- temporarily relieves common cold/flu symptoms:
- cough due to minor throat and bronchial irritation
- minor aches and pains    ■ sore throat    ■ fever
- nasal congestion            ■ headache

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin reddening    ■ blisters    ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.



**McKesson**

Another Quality Product  
Distributed by McKesson  
One Post Street, San Francisco, CA 94104  
Money Back Guarantee  
www.healthmart.com/healthmartbrand

**PARENTS:**

Learn about teen medicine abuse.

www.StopMedicineAbuse.org

65634 06 F3

PEEL BACK AT CORNER FOR MORE INFORMATION →

### Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**    ■ liver disease  
 ■ heart disease    ■ thyroid disease    ■ diabetes    ■ high blood pressure  
 ■ trouble urinating due to an enlarged prostate gland  
 ■ cough that occurs with too much phlegm (mucus)  
 ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

#### When using this product do not use more than directed

- Stop use and ask a doctor if**    ■ new symptoms occur
- you get nervous, dizzy or sleepless    ■ redness or swelling is present
  - fever gets worse or lasts more than 3 days
  - pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
  - cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### Drug Facts (continued)

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- take only as directed – see Overdose warning
  - only use the dose cup provided    ■ do not exceed 4 doses per 24 hrs
- |                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs      | 15 mL every 4 hrs |
| children 4 to under 6 yrs       | ask a doctor      |
| children under 4 yrs            | do not use        |

**Other information**    ■ each 15 mL contains: sodium 7 mg  
 ■ store at 20-25°C (68-77°F)

**Inactive ingredients** butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

#### Questions or comments? 1-800-719-9260

\*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Cold & Flu.

## HEALTH MART DAYTIME

acetaminophen, dextromethorphan hbr, phenylephrine hcl solution

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62011-0065
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL

<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BUTYLATED HYDROXYANISOLE</b> (UNII: REK4960K2U)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM</b> (UNII: 3980JIH2SW)	
<b>POLYETHYLENE GLYCOLS</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A)	

### Product Characteristics

<b>Color</b>	ORANGE (clear)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MENTHOL (with fruit)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0065-1	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/11/2012	
2	NDC:62011-0065-2	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/06/2012	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	01/11/2012	

**Labeler** - Mckesson (177667227)