

DURADRY PM SWEAT MINIMIZING ANTIPERSPIRANT GEL- aluminum chloride hexahydrate gel
Novadore USA Inc

DURADRY PM Sweat Minimizing Antiperspirant Gel

Active ingredient

ALUMINUM CHLORIDE (HEXAHYDRATE) 15%

Purpose

Antiperspirant

Uses

Reduces underarm perspiration

Warnings

For external use only
Stop use if rash or irritation occurs.

Do not use

- On broken, irritated or recently shaved skin.
- Immediately after bathing.
- In or near eyes. If contact occurs, rise eyes thoroughly with water.

Ask a doctor before use

if you have kidney disease

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply to underarms only.
- Apply a small amount of product to underarms at bedtime. It may itch during initial application.
- The next morning, rinse underarms with abundant water and apply a regular antiperspirant. For best results, use DURADRY® AM.
- Apply DURADRY® PM for a few nights in a row or until the excessive perspiration is controlled. Then Apply twice a week or as needed for maintenance.

Inactive ingredients

Aqua, Propylene Glycol, Xanthan Gum, Sodium Bicarbonate, Disodium EDTA, Salicylic Acid.

Questions?

Learn more at duradry.com

Dist. by:

NOVADORE USA, LLC
Miami, FL 33179

DURADRY PM (ALUMINUM CHLORIDE HEXAHYDRATE)

Duradry



ALUMINUM CHLORIDE (HEXAHYDRATE) ANTIPERSPIRANT

PM Antiperspirant
0.4 FL. OZ. (10.5 mL)

**A Dry Tomorrow
Starts Tonight.**

USE ONLY AT BEDTIME

swallowed, get medical help or contact a Poison Control Center immediately.

Drug Facts

Active Ingredient:	Purpose:
Aluminum Chloride (Hexahydrate) 15%.....	Antiperspirant

Uses: Reduces underarm perspiration

Warnings:

For external use only

Do not use • On broken, irritated or recently shaved skin • Immediately after bathing • In or near eyes. If contact occurs, rinse eyes thoroughly with water. Stop use if rash or irritation occurs • **Ask a doctor before use if you have kidney disease. Keep out of reach of children.** If

Directions: Apply to underarms only • Apply a small amount of product to underarms at bedtime. It may itch during initial applications • The next morning, rinse underarms with abundant water and soap, for best results use DURADRY® Wash, and apply a regular antiperspirant. For best results, use DURADRY® AM • Apply DURADRY® PM for a few nights in a row or until the excessive perspiration is controlled. Then apply twice a week or as needed for maintenance.

Inactive Ingredients: Aqua, Propylene Glycol, Xanthan Gum, Sodium Bicarbonate, Disodium EDTA, Salicylic Acid.

Questions? Learn more at duradry.com

Dist. by NOVADORE USA, LLC • Miami, FL 33179

**A Dry Tomorrow
Starts Tonight.**

USE ONLY AT BEDTIME

Drug Facts

Active Ingredient: Aluminum Chloride (Hexahydrate)15%..... **Purpose:** Antiperspirant.

Uses: Reduces underarm perspiration.

Warnings:

For external use only.

Do not use broken, irritated or recently shaved skin • Immediately after bathing • In or near eyes. If contact occurs, rinse eyes thoroughly with water.

Stop use and ask a doctor if rash or irritation occurs.

Ask a doctor before use if you have kidney disease.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately.

Directions: Apply to underarms only • Apply a small amount of product to underarms at bedtime. It may itch during initial applications • The next morning, rinse underarms with abundant water and soap (for best results, use DURADRY® Wash) and apply a regular antiperspirant. For best results, use DURADRY® AM • Apply DURADRY® PM for a few nights in a row or until the excessive perspiration is controlled. Then apply twice a week or as needed for maintenance.

Inactive Ingredients: Aqua, Propylene Glycol, Xanthan Gum, Sodium Bicarbonate, Disodium EDTA, Salicylic Acid.

Questions? Learn more at duradry.com

Dist. by
NOVADORE USA, LLC
Miami, FL 33179



Duradry



PM Antiperspirant

DURADRY PM SWEAT MINIMIZING ANTIPERSPIRANT GEL

aluminum chloride hexahydrate gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69990-102
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM CHLORIDE (UNII: 3CYT62D3GA) (ALUMINUM CATION - UNII:3XHB1D032B)	ALUMINUM CHLORIDE	150 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

SALICYLIC ACID (UNII: O414PZ4LPZ)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
WATER (UNII: 059QF0KO0R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69990-102-01	1 in 1 POUCH	02/20/2018	
1		10.5 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M019	01/01/2015	

Labeler - Novadore USA Inc (079777451)

Registrant - Novadore USA Inc (079777451)

Revised: 11/2023

Novadore USA Inc