

**EQUATE MILK OF MAGNESIA- magnesium hydroxide liquid**  
**Wal-Mart Stores Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Wal-Mart Milk of Magnesia Drug Facts**

**Active ingredient (in each 15 mL TBSP)**

Magnesium hydroxide 1200 mg

**Purpose**

Saline laxative

**Uses**

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

**Warnings**

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

**Ask a doctor or pharmacist before use if you are**

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

**Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- shake well before use

- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.
children 6 to 11 years	1 to 2 TBSP.
children under 6 years	ask a doctor

### Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- keep tightly closed

### Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

### Questions or comments?

1-888-287-1915

### Principal Display Panel

Compare to Phillips'® Milk of Magnesia Wild Cherry Active Ingredient

Milk of Magnesia

Magnesium Hydroxide

Saline Laxative

Comfortable, cramp free relief

Stimulant free

Wild Cherry Flavor

ALCOHOL 0.4%

12 FL OZ (355 mL)

NDC 49035-949-40

**equate**<sup>®</sup>

Compare  
to Phillips'<sup>®</sup>  
Milk of Magnesia  
Wild Cherry  
Active  
Ingredient\*

# Milk of Magnesia

## Magnesium Hydroxide

### Saline Laxative

- Comfortable, cramp free relief
- Stimulant free

### Wild Cherry Flavor

ALCOHOL 0.4%

12 FL OZ (355 mL)



: 94940 2E F2

## Drug Facts

**DO NOT USE IF PRINTED NECKBAND IS  
BROKEN OR MISSING.**

**Active ingredient (in each 15 mL TBSP)**      **Purpose**  
Magnesium hydroxide 1200 mg.....Saline laxative

**Uses**    ■ relieves occasional constipation (irregularity)  
■ generally produces bowel movement in ½ to 6 hours

### Warnings

Ask a doctor before use if you have

- kidney disease    ■ a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug.  
Take this product two or more hours before or after other drugs.  
Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided    ■ TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.
children 6 to 11 years	1 to 2 TBSP.
children under 6 years	ask a doctor

### Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)    ■ keep tightly closed

**Inactive ingredients** anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

**Questions or comments?** 1-800-719-9260

EB HD 0h6hb :

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips'® Milk of Magnesia.

GLUTEN FREE

DISTRIBUTED BY FOODHOLD U.S.A., LLC  
LANDOVER, MD 20785 · 1-877-846-9949  
©2013 S&S Brands, LLC

Quality guaranteed or your money back.



**EQUATE MILK OF MAGNESIA**

magnesium hydroxide liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49035-949
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>MAGNESIUM HYDRO XIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDRO XIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	

**Product Characteristics**

<b>Color</b>	PINK (Opaque Light pink)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY (wild cherry)	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:49035-949-35	769 mL in 1 BOTTLE		
2	NDC:49035-949-40	355 mL in 1 BOTTLE		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph not final	part334	04/21/2005	

**Labeler** - Wal-Mart Stores Inc (051957769)