

# HEADACHE RELIEF EXTRA STRENGTH- acetaminophen, aspirin, caffeine tablet The Kroger Co.

-----  
**Drug Facts**

## Active ingredients (in each caplet)

**Acetaminophen 250 mg**

**Aspirin 250 mg (NSAID)\***

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

## Purposes

**Pain reliever**

**Pain reliever**

Pain reliever aid

## Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - a cold
  - arthritis
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

If a skin reaction, occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives

- facial swelling
- shock
- asthma (wheezing)

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have asthma
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

### **Stop use and ask doctor if**

- an allergic reaction occurs. Seek medical help right away
  - any new symptoms appear
- you experience any of the following signs of stomach bleeding
  - feel faint

- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- pains gets worse or last for more than 10 days
- fever gets worse or lasts for more than 3 days
- painful area is red or swollen

These could be signs of a serious condition

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; do not take more than 8 caplets in 24 hours
- children under 12 years: do not use unless directed by a doctor

### **Other information**

- store between 20°-25°C (68°-77°F)

### **Inactive ingredients**

carnauba wax\*\*, colloidal silicon dioxide, corn starch, hypromellose, microcrystalline cellulose, mineral oil\*\*, polyethylene glycol\*\*, povidone, sodium starch glycolate\*\*, stearic acid, talc\*\*, titanium dioxide

\*\*contains one or more of these ingredients

### **Questions or comments?**

Call **1-800-632-6900**

### **Principal Display Panel**

COMPARE TO the active ingredients in EXCEDRIN® EXTRA STRENGTH \*\*\*See back panel

## **Headache Relief Extra Strength**

Acetaminophen 250 mg - Pain Reliever

Aspirin (NSAID) 250 mg - Pain Reliever

Caffeine 65 mg - Pain Reliever Aid

CAPLETS

\*\*\*Excedrin is a registered trademark of GSK Consumer Healthcare, Warren, New Jersey 07059. GSK Consumer Healthcare is not affiliated with The Kroger Co. or this product

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

DISTRIBUTED BY THE KROGER CO.

CINCINNATI, OHIO 45202

[www.kroger.com](http://www.kroger.com)

**Package Label**

Exp. Date:  
Lot No.:

PLD-A1036 F0004964



DISTRIBUTED BY THE KROGER CO.  
CINCINNATI, OHIO 45207  
QUALITY GUARANTEE  
www.kroger.com

COMPARE TO the active ingredients in EXCEDRIN® EXTRA STRENGTH  
\*\*\*See back panel



**Headache Relief Extra Strength**  
Acetaminophen 250 mg - Pain Reliever  
Aspirin (NSAID) 250 mg - Pain Reliever  
Caffeine 65 mg - Pain Reliever Aid

NDC 30142-625-10



100 CAPLETS



**Drug Facts**  
**Active ingredients (in each caplet)**  
Acetaminophen 250 mg.....Pain reliever  
Aspirin 250 mg (NSAID)\*.....Pain reliever  
Caffeine 65 mg.....Pain reliever aid  
\*nonsteroidal anti-inflammatory drug

**Uses**  
temporarily relieves minor aches and pains due to  
■ headache ■ a cold ■ arthritis  
■ muscular aches ■ toothache  
■ premenstrual and menstrual cramps

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock  
■ asthma (wheezing)  
Allergy alert: Acetaminophen may cause severe skin reactions.  
Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:  
■ more than 8 caplets in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Drug Facts (continued)**  
**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.  
**Do not use**  
■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**  
■ you have liver disease  
■ stomach bleeding warning applies to you  
■ you have asthma ■ you are taking a diuretic  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
**Ask a doctor or pharmacist before use if you are**  
■ taking a prescription drug for diabetes, gout, or arthritis  
■ taking any other drug or are under a doctor's care for any serious condition

**Stop use and ask a doctor if**  
■ an allergic reaction occurs. Seek medical help right away.  
■ any new symptoms appear  
■ you experience any of the following signs of stomach bleeding  
■ feel faint ■ vomit blood  
■ have bloody or black stools  
■ have stomach pain that does not get better  
■ ringing in the ears or loss of hearing occurs  
■ pain gets worse or lasts for more than 10 days  
■ fever gets worse or lasts for more than 3 days  
■ painful area is red or swollen  
These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Drug Facts (continued)**  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ do not use more than directed  
■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 2 caplets every 6 hours; do not take more than 8 caplets in 24 hours  
■ children under 12 years: do not use unless directed by a doctor

**Other information**  
■ store between 20°-25°C (68°-77°F)

**Inactive ingredients** carnauba wax\*\*, colloidal silicon dioxide, corn starch, hypromellose, microcrystalline cellulose, mineral oil\*\*, polyethylene glycol\*\*, povidone, sodium starch glycolate\*\*, stearic acid, talc\*\*, titanium dioxide\*\*  
\*\*contains one or more of these ingredients

**Questions or comments?**  
Call 1-800-632-8600

Excedrin is a registered trademark of GSK Consumer Healthcare, Warren, New Jersey 07059. GSK Consumer Healthcare is not affiliated with The Kroger Co. or this product.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.  
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

# KROGER Extra Strength Headache Relief

## HEADACHE RELIEF EXTRA STRENGTH

acetaminophen, aspirin, caffeine tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:30142-625
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL370;S53
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-625-10	1 in 1 BOX	04/30/2018	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
	NDC:30142-			

2	NDC:30142-625-24	1 in 1 BOX	04/30/2018	
2		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013		04/30/2018	

**Labeler** - The Kroger Co. (006999528)

Revised: 6/2024

The Kroger Co.