

OXYTET SOLUBLE- oxytetracycline hydrochloride powder, for solution
Huvepharma, Inc

Oxytet[®] Soluble

280 GRAM Front Panel

Oxytet[®] Soluble

(oxytetracycline hydrochloride soluble powder)
Antibiotic

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

This packet contains 102.4 g of oxytetracycline hydrochloride and will make:

512 gallons (1936 liters) containing 200 mg oxytetracycline hydrochloride per gallon
128 gallons (484 liters) containing 800 mg oxytetracycline hydrochloride per gallon

This packet will treat 10,240 lbs of swine.

This packet will treat 4,096 lbs of turkey at 25 mg/lb body weight.

RECOMMENDED PACKET

STORAGE CONDITIONS:

Store between 20°C - 25°C
(68°F - 77°F) with excursions
permitted between 15°C - 40°C
(59°F - 104°F).

See back panel for instructions.

Net Weight: 9.87 oz (280 g)

For Use in Drinking Water Only

Not For Use in Liquid Medicated Feeds

Not For Use in Humans

Keep Out of Reach of Children

HAZARDOUS REMEDY

Restricted Drug (California) - Use only as directed.

HUVEPHARMA[®]

Manufactured for:
Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269

HUVEPHARMA and Oxytet
are registered trademarks of
Huvepharma EOOD

USE DIRECTIONS:

Mix fresh solutions daily. Use as the sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. As a generalization, 200 chickens will drink 1 gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer up to 5 days to swine and 7 to 14 days for chickens and turkeys. **NOTE:** The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperatures and humidity, each of which affects water consumption.

CAUTION

Use as the sole source of oxytetracycline. Not to be used for more than 5 consecutive days in swine or 14 consecutive days in chickens and turkeys.

WARNING -

Do not feed to birds producing eggs for human consumption.

MIXING DIRECTIONS FOR WATER MEDICATION CHICKENS — TURKEYS — SWINE

Dosage and indications for use in water proportioners - Add the following amount to 1 gallon of stock solution when proportioner is set to meter at the rate of 1 oz per gallon.

C Disease		Treatment Pack/Gallon	
		Level	Stock
H I C K E N S	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200 - 400 mg	¼ - ½ (70 - 140 g)
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400 - 800 mg	½ - 1 (140 - 280 g)
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400 - 800 mg	½ - 1 (140 - 280 g)
T U R K E Y S	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200 - 400 mg	¼ - ½ (70 - 140 g)
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	½ (140 g)
	GROWING TURKEYS — Control of complicating bacterial	25	Varies with

Y	organisms associated with bluecomb	mg/lb	age &
S	(transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	body weight	water consumption
FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE -			
S	Bacterial enteritis caused by <i>Escherichia coli</i> and		
W	<i>Salmonella choleraesuis</i> , susceptible to	10 mg/lb	Varies with
I	oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella</i>	body weight	age &
N	<i>multocida</i> , susceptible to oxytetracycline.		water consumption
E	FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.		

1400 GRAM Front Panel

Oxytet[®] Soluble

**(oxytetracycline hydrochloride soluble powder)
Antibiotic**

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

This packet contains 512 g of oxytetracycline hydrochloride and will make:

2560 gallons (9690 liters) containing 200 mg oxytetracycline hydrochloride per gallon

1280 gallons (4845 liters) containing 400 mg oxytetracycline hydrochloride per gallon

640 gallons (2422 liters) containing 800 mg oxytetracycline hydrochloride per gallon

This packet will treat 51,200 lbs of swine.

This packet will treat 20,480 lbs of turkey at 25 mg/lb body weight.

RECOMMENDED PACKET

STORAGE CONDITIONS:

Store between 20°C - 25°C

(68°F - 77°F) with excursions

permitted between 15°C - 40°C

(59°F - 104°F).

See back panel for instructions.

Net Weight

3.09 lbs (1400 g)

For Use in Drinking Water Only

Not For Use in Liquid Medicated Feeds

Not For Use in Humans

Keep Out of Reach of Children

HAZARDOUS REMEDY

Restricted Drug (California) - Use

only as directed.

HUVEPHARMA®

Manufactured for:
Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269

HUVEPHARMA and Oxytet
are registered trademarks of
Huvepharma EOOD

13924800 Rev. 02-2021
Approved by FDA under NADA #130-435

USE DIRECTIONS:

Mix fresh solutions daily. Use as the sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. As a generalization, 200 chickens will drink 1 gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer up to 5 days to swine and 7 to 14 days for chickens and turkeys.

NOTE: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperatures and humidity, each of which affects water consumption.

CAUTION:

Use as the sole source of oxytetracycline. Not to be used for more than 5 consecutive days in swine or 14 consecutive days in chickens and turkeys.

Warning -

Do not feed to birds producing eggs for human consumption.

MIXING DIRECTIONS FOR WATER MEDICATION CHICKENS — TURKEYS — SWINE

Dosage and indications for use in water proportioners - Add 1 pack to the following amount of water to make a stock solution for use in a proportioner set to meter at the rate of one ounce per gallon.

C	Disease	Treatment Gallons	Level	Stock/Pack
H	Control of infectious synovitis caused by <i>Mycoplasma</i>	200 - 400		20 -

C K E N S	<i>synoviae</i> , susceptible to oxytetracycline.	mg	10
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400 - 800 mg	10 - 5
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400 - 800 mg	10 - 5
T U R K E Y S	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200 - 400 mg	20 - 10
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	10
	GROWING TURKEYS — Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE -			
S W I N E	Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption
	Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.		
	FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.		

1772 GRAM Front Panel

Oxytet[®] Soluble

(oxytetracycline hydrochloride soluble powder) Antibiotic

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

This packet contains 648 g of oxytetracycline hydrochloride and will make:

3240 gallons (12,265 liters) containing 200 mg oxytetracycline hydrochloride per gallon
 1620 gallons (6,132 liters) containing 400 mg oxytetracycline hydrochloride per gallon
 810 gallons (3,066 liters) containing 800 mg oxytetracycline hydrochloride per gallon

This packet will treat 64,800 lbs of swine.

This packet will treat 25,920 lbs of turkey at 25 mg/lb body weight.

RECOMMENDED PACKET

STORAGE CONDITIONS:

Store between 20°C - 25°C
 (68°F - 77°F) with excursions
 permitted between 15°C - 40°C
 (59°F - 104°F).

See back panel for instructions.

Net Weight
3.91 lbs (1772 g)

For Use in Drinking Water Only

Not For Use in Liquid Medicated Feeds

Not For Use in Humans

Keep Out of Reach of Children

HAZARDOUS REMEDY

Restricted Drug (California) - Use only as directed.

HUVEPHARMA®

Manufactured for:
Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269

HUVEPHARMA and Oxytet are registered trademarks of Huvepharma EOOD.

13925200 (Rev. 02-2021)
Approved by FDA under #130-435

USE DIRECTIONS:

Mix fresh solutions daily. Use as the sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. As a generalization, 200 chickens will drink 1 gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer up to 5 days to swine and 7 to 14 days for chickens and turkeys.

NOTE: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperatures and humidity, each of which affects water consumption.

CAUTION:

Use as the sole source of oxytetracycline. Not to be used for more than 5 consecutive days in swine or 14 consecutive days in chickens and turkeys.

WARNING -


Do not feed to birds producing eggs for human consumption.

MIXING DIRECTIONS FOR WATER MEDICATION CHICKENS — TURKEYS — SWINE

Dosage and indications for use in water proportioners - Add 1 pack to the following amount of water to make a stock solution for use in a proportioner set to meter at the rate of one ounce per gallon.

	Disease	Treatment Level	Gallons Stock/Pack
C H I C K E N S	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200 - 400 mg	25.3 - 12.6
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400 - 800 mg	12.6 - 6.3
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400 - 800 mg	12.6 - 6.3
T U R K E Y S	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200 - 400 mg	25.3 - 12.6
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	12.6
	GROWING TURKEYS — Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
S W I N E	FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE – Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption
	Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption
	FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption

280 GRAM Packet Label Image



MIXING DIRECTIONS FOR WATER MEDICATION CHICKENS — TURKEYS — SWINE Dosage and indications for use in water proportioners - Add the following amount to 1 gallon of stock solution when proportioner is set to meter at the rate of 1 oz per gallon.

	Disease	Treatment Level	Pack/Gallon Stock
C H I C K E N S	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200 - 400 mg	¼ - ½ (70 - 140 g)
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400 - 800 mg	½ - 1 (140 - 280 g)
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400 - 800 mg	½ - 1 (140 - 280 g)
T U R K E Y S	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200 - 400 mg	¼ - ½ (70 - 140 g)
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	½ (140 g)
	GROWING TURKEYS — Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
S W I N E	FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE – Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption
	Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption
	FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption

CORNER RADIUS

Oxytet[®] Soluble

(oxytetracycline hydrochloride soluble powder)

Antibiotic

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

This packet contains 102.4 g of oxytetracycline hydrochloride and will make:
512 gallons (1936 liters) containing 200 mg oxytetracycline hydrochloride per gallon
128 gallons (484 liters) containing 800 mg oxytetracycline hydrochloride per gallon

This packet will treat 10,240 lbs of swine.

This packet will treat 4,096 lbs of turkey at 25 mg/lb body weight.

RECOMMENDED PACKET STORAGE CONDITIONS:
Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

See back panel for instructions.

Net Weight: 9.87 oz (280 g)

For Use in Drinking Water Only

Not For Use in Liquid Medicated Feeds

Not For Use in Humans

Keep Out of Reach of Children

HAZARDOUS REMEDY

Restricted Drug (California) - Use only as directed.



Manufactured for:
Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269

HUVEPHARMA and Oxytet are registered trademarks of Huvepharma EOOD

13925000 (Rev. 02-2021)
Approved by FDA under NADA # 130-435

CORNER RADIUS

USE DIRECTIONS: Mix fresh solutions daily. Use as the sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. As a generalization, 200 chickens will drink 1 gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer up to 5 days to swine and 7 to 14 days for chickens and turkeys. **NOTE:** The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperatures and humidity, each of which affects water consumption.

CAUTION: Use as the sole source of oxytetracycline. Not to be used for more than 5 consecutive days in swine or 14 consecutive days in chickens and turkeys.

WARNING - Do not feed to birds producing eggs for human consumption.

NO COPY

1400 GRAM Packet Label

NO COPY

MIXING DIRECTIONS FOR WATER MEDICATION CHICKENS — TURKEYS — SWINE

Dosage and indications for use in water proportioners - Add 1 pack to the following amount of water to make a stock solution for use in a proportioner set to meter at the rate of one ounce per gallon.

	Disease	Treatment Level	Gallons Stock/Pack
C H I C K E N S	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200 - 400 mg	20 - 10
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400 - 800 mg	10 - 5
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400 - 800 mg	10 - 5
	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200 - 400 mg	20 - 10
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	10

C H I C K E N S	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	10
	GROWING TURKEYS — Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronavirus enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
S W I N E	FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE — Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption
	FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.		

CORNER RADIUS

Oxytet[®] Soluble

(oxytetracycline hydrochloride soluble powder)

Antibiotic

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

This packet contains 512 g of oxytetracycline hydrochloride and will make:
 2560 gallons (9690 liters) containing 200 mg oxytetracycline hydrochloride per gallon
 1280 gallons (4845 liters) containing 400 mg oxytetracycline hydrochloride per gallon
 640 gallons (2422 liters) containing 800 mg oxytetracycline hydrochloride per gallon

This packet will treat 51,200 lbs of swine.

This packet will treat 20,480 lbs of turkey at 25 mg/lb body weight.

RECOMMENDED PACKET

STORAGE CONDITIONS:
 Store between 20°C - 25°C
 (68°F - 77°F) with excursions
 permitted between 15°C - 40°C
 (59°F - 104°F).

See back panel for instructions.

Net Weight
3.09 lbs (1400 g)

For Use in Drinking Water Only

Not For Use in Liquid Medicated Feeds

Not For Use in Humans

Keep Out of Reach of Children

HAZARDOUS REMEDY

Restricted Drug (California) - Use only as directed.



Manufactured for:
 Huvepharma, Inc.
 525 Westpark Drive, Suite 230
 Peachtree City, GA 30269

HUVEPHARMA and Oxytet
 are registered trademarks of
 Huvepharma EOOD

13924800 Rev. 02-2021
 Approved by FDA under NADA # 130-435

CORNER RADIUS

USE DIRECTIONS: Mix fresh solutions daily. Use as the sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. As a generalization, 200 chickens will drink 1 gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer up to 5 days to swine and 7 to 14 days for chickens and turkeys. **NOTE:** The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperatures and humidity, each of which affects water consumption. **CAUTION:** Use as the sole source of oxytetracycline. Not to be used for more than 5 consecutive days in swine or 14 consecutive days in chickens and turkeys.

WARNING - Do not feed to birds producing eggs for human consumption.

Oxytet[®] Soluble
 (oxytetracycline hydrochloride soluble powder)

Antibiotic

NO COPY

1772 GRAM Packet Label

NO COPY

MIXING DIRECTIONS FOR WATER MEDICATION CHICKENS — TURKEYS — SWINE
Dosage and indications for use in water proportioners - Add 1 pack to the following amount of water to make a stock solution for use in a proportioner set to meter at the rate of one ounce per gallon.

	Disease	Treatment Level	Gallons Stock/Pack
C H I C K E N S	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200 - 400 mg	25.3 - 12.6
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400 - 800 mg	12.6 - 6.3
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400 - 800 mg	12.6 - 6.3
T U R K E Y S	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200 - 400 mg	25.3 - 12.6
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	12.6
	GROWING TURKEYS — Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
S W I N E	FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE — Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline. FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption

CORNER RADIUS

Oxytet[®] Soluble

(oxytetracycline hydrochloride soluble powder)

Antibiotic

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

This packet contains 648 g of oxytetracycline hydrochloride and will make:
 3240 gallons (12,265 liters) containing 200 mg oxytetracycline hydrochloride per gallon
 1620 gallons (6,132 liters) containing 400 mg oxytetracycline hydrochloride per gallon
 810 gallons (3,066 liters) containing 800 mg oxytetracycline hydrochloride per gallon

This packet will treat 64,800 lbs of swine.

This packet will treat 25,920 lbs of turkey at 25 mg/lb body weight.

RECOMMENDED PACKET STORAGE CONDITIONS:
 Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

See back panel for instructions.

Net Weight
3.91 lbs (1772 g)

For Use in Drinking Water Only
Not For Use in Liquid Medicated Feeds
Not For Use in Humans
Keep Out of Reach of Children
HAZARDOUS REMEDY

Restricted Drug (California) - Use only as directed.



Manufactured for:
 Huvepharma, Inc.
 525 Westpark Drive, Suite 230
 Peachtree City, GA 30269

HUVEPHARMA and Oxytet are registered trademarks of Huvepharma EOOD

13925200 (Rev. 02-2021)
 Approved by FDA under NADA #130-435

CORNER RADIUS

USE DIRECTIONS: Mix fresh solutions daily. Use as the sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. As a generalization, 200 chickens will drink 1 gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer up to 5 days to swine and 7 to 14 days for chickens and turkeys. **NOTE:** The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperatures and humidity, each of which affects water consumption.

CAUTION: Use as the sole source of oxytetracycline. Not to be used for more than 5 consecutive days in swine or 14 consecutive days in chickens and turkeys.

WARNING - Do not feed to birds producing eggs for human consumption.

Oxytet[®] Soluble

(oxytetracycline hydrochloride soluble powder)

Antibiotic

NO COPY

oxytetracycline hydrochloride powder, for solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:23243-6759
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYTETRACYCLINE HYDROCHLORIDE (UNII: 4U7K4N52Z M) (OXYTETRACYCLINE ANHYDROUS - UNII:SLF0D9077S)	OXYTETRACYCLINE HYDROCHLORIDE	0.357 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-6759-1	50 in 1 PAIL		
1		280 g in 1 PACKET		
2	NDC:23243-6759-2	10 in 1 PAIL		
2		1400 g in 1 PACKET		
3	NDC:23243-6759-3	6 in 1 PAIL		
3		1772 g in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA130435	04/25/2017	

Labeler - Huvepharma, Inc (619153559)

Registrant - Huvepharma EOOD (552671651)

Establishment

Name	Address	ID/FEI	Business Operations
Huvepharma, Inc - St. Louis		883128204	manufacture, analysis, pack, label

Revised: 10/2023

Huvepharma, Inc