

MOTION SICKNESS RELIEF- dimenhydrinate tablet
Valu Merchandisers Company

Best Choice 44-198

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-

1222) right away.

Directions

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

Other information

- **each tablet contains:** calcium 35 mg
- see end flap for expiration date and lot number
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from moisture

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?

1-800-426-9391

Principal display panel

Best Choice®

Compare to the active ingredient in Dramamine® Original Formula*

Motion Sickness Relief

Dimenhydrinate 50 mg
Antiemetic

Prevents:

- Nausea
- Vomiting
- Dizziness

FOR CHILDREN & ADULTS

Actual Size

12 Tablets

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by
Medtech Products Inc., owner of the registered
trademark Dramamine® Original Formula.
50844 REV0518A19802

PROUDLY DISTRIBUTED BY:
ASSOCIATED WHOLESALE GROCERS, INC.
KANSAS CITY, KANSAS 66106

Best Choice®



Motion Sickness Relief

DIMENHYDRINATE 50 mg ANTIEMETIC

Compare to the active ingredient in
Dramamine®
Original Formula*

Prevents:
- Nausea
- Vomiting
- Dizziness

FOR CHILDREN & ADULTS



Actual Size



12 TABLETS

Best Choice®

Motion Sickness Relief

DIMENHYDRINATE 50 mg ANTIEMETIC

Prevents:
- Nausea
- Vomiting
- Dizziness

FOR CHILDREN & ADULTS



Actual Size



12 TABLETS

Best Choice®

Motion Sickness Relief

DIMENHYDRINATE 50 mg ANTIEMETIC

Compare to the active ingredient in
Dramamine®
Original Formula*

Prevents:
- Nausea
- Vomiting
- Dizziness

FOR CHILDREN & ADULTS



Actual Size



12 TABLETS

E7732RZ

PROUDLY DISTRIBUTED BY:
ASSOCIATED WHOLESALE GROCERS, INC.
KANSAS CITY, KANSAS 66106
Scan here for more product information.
Call 1-844-292-1112 for more product information.



*This product is not manufactured or distributed by Medmark/Dramamine® Original Formula.
50844 REV0518A19802

B-2115-198-02-RR
REV0518A19802

BEST CHOICE
58647 1123 LNK

No print/No varnish
Lot & Expiry date



0

one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours, do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	1/2 to 1 tablet every 6-8 hours, do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	1/2 tablet every 6-8 hours, do not exceed 1 1/2 tablets in 24 hours, or as directed by a doctor

Other information

- each tablet contains: calcium 35 mg
- see end flap for expiration date and lot number
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGING OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture
- protect from moisture

Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments? 1-800-426-9391

Uses for prevention and treatment of these symptoms associated with motion sickness: ■ nausea ■ vomiting ■ dizziness

Warnings

Do not use for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis ■ difficulty in urination due to enlargement of the prostate gland


Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product ■ marked drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic beverages ■ use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING



Best Choice 44-198

MOTION SICKNESS RELIEF

dimenhydrinate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-198
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, 8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;198
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-198-02	2 in 1 CARTON	02/15/2019	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination		

Product			
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	02/15/2019	

Labeler - Valu Merchandisers Company (868703513)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(63941-198)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63941-198) , pack(63941-198)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(63941-198)

Revised: 12/2023

Valu Merchandisers Company