

**DRX CHOICE FIBER THERAPY- methylcellulose powder, for solution**  
**RARITAN PHARMACEUTICALS INC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**DRx Choice Fiber Therapy Methylcellulose 2 g Bulk Forming Laxative**

***Active ingredient (in each rounded tablespoon)***

Methylcellulose (a non-allergenic fiber) 2g

***Purpose***

Bulk-forming fiber laxative

***Uses***

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 12-72 hours

***Warnings***

**Choking: taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.**

**Ask a doctor before use if you have**

- a sudden change in bowel habits that persists for two weeks
- abdominal pain, nausea or vomiting

**Stop use and ask a doctor if**

- constipation lasts more than 7 days
- you have rectal bleeding

These could be signs of a serious condition.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

## Directions

- **MIX THIS PRODUCT (CHILD OR ADULT DOSE) WITH AT LEAST 8 OUNCES (A FULL GLASS) OF WATER OR OTHER FLUID. TAKING THIS PRODUCT WITHOUT ENOUGH LIQUID MAY CAUSE CHOKING. SEE CHOKING WARNING**
- use product at the first sign of constipation or irregularity
- put one dose in a full glass of cold water
- stir briskly and drink promptly
- drinking another glass of water is helpful

Age	Dose
adults & children 12 years of age and over	start with 1 rounded <b>tablespoon</b> . Increase as needed, 1 rounded <b>tablespoon</b> at a time, up to 3 times per day.
children 6 - 11 years of age	start with 2 level <b>teaspoons</b> . Increase as needed, 2 level <b>teaspoons</b> at a time, up to 3 times per day.
children under 6 years of age	consult a physician

## Other information

- **each rounded tablespoon contains:** calcium 85mg and potassium 125mg
- each rounded tablespoon contributes 27 calories from maltodextrin
- store at room temperature
- protect contents from humidity
- keep tightly closed

## Inactive ingredients

Anhydrous citric acid, FD&C yellow #6 lake, flavor, maltodextrin, potassium citrate, silica, sucralose, tricalcium phosphate,

## Questions or comments?

**1-866-467-2748**

## Principal Display Panel

DRx Choice®

Compare to the Active Ingredient in Citrucel® Powder\*

**NDC 68163-517-16**

**Sugar free**

**Soluble fiber therapy**

**Methylcellulose Fiber Therapy for Irregularity**

- **Gentle and clinically proven effective\*\***
- **Gives you additional fiber to help relieve occasional constipation**

**Fiber for irregularity that won't cause excess gas\*\*\***

Orange Flavor

Naturally & Artificially Flavored

NET WT 16.9 OZ (479g)

**Directions for use:**

1. Fill glass with at least 8 oz. of cold water.
2. Measure rounded tablespoon of product (one dose) and stir product into water. Stir briskly and drink promptly.

\*\*When Used As Directed.

\*\*\*Based On Laboratory Testing. Individual Results may Vary

**TAMPER EVIDENT: DO NOT USE PRINTED INNER SEAL IS BROKEN OR MISSING**

GLUTEN FREE

\*This product is not manufactured or distributed by GSK Consumer Healthcare, the distributor of Citrucel® Powder.

**Manufactured By:**

**Raritan Pharmaceuticals**

**8 Joana Court,**

**East Brunswick, NJ 08816**

Compare to the active ingredient in Citrucel® Powder\*  
NDC 68163-517-16

# Dr. Choice®

**sugar free**  
**soluble**  
**fiber therapy**


**Methylcellulose**  
**Fiber Therapy for Irregularity**

Gentle and clinically proven effective\*\*  
Gives you additional fiber to help  
relieve occasional constipation  
Fiber for irregularity that won't cause excess gas\*\*\*


**Orange**  
(Natural Flavors and Citrus Oil Essence)

**NET WT. 16.9 OZ (479g)**

**DIRECTIONS FOR USE (NO SCOOP DOSING):**



1. Fill glass with at least 8 oz. of cold water.



2. Measure rounded tablespoon of product (one dose) and stir product into water. Stir briskly and drink promptly.

\*\*When Used As Directed.

\*\*\*Based On Laboratory Testing. Individual Results May Vary.

**TAMPER EVIDENT: DO NOT USE IF PRINTED INNER SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each rounded tablespoon)**  
Methylcellulose (a non-allergenic fiber) 2g **Purpose**  
Bulk-forming fiber laxative

**Uses**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 12-72 hours

**Warnings**

**Choking:** taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have a difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

**Ask a doctor before use if you have**

- a sudden change in bowel habits that persists for two weeks
- abdominal pain, nausea or vomiting

**Stop use and ask a doctor if**

- constipation lasts more than 7 days
- you have rectal bleeding.

These could be signs of a serious condition.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

**Directions**

- MIX THIS PRODUCT (CHILD OR ADULT DOSE) WITH AT LEAST 8 OUNCES (A FULL GLASS) OF WATER OR OTHER FLUID. TAKING THIS PRODUCT WITHOUT ENOUGH LIQUID MAY CAUSE CHOKING. SEE CHOKING WARNING.**
- use product at the first sign of constipation or irregularity
- put one dose in a full glass of cold water
- stir briskly and drink promptly
- drinking another glass of water is helpful

Age	Dose
adults and children 12 years of age and over	start with 1 rounded <b>tablespoon</b> . Increase as needed, 1 rounded <b>tablespoon</b> at a time, up to 3 times per day.
children 6-11 years of age	start with 2 level <b>teaspoons</b> . Increase as needed, 2 level <b>teaspoons</b> at a time, up to 3 times per day.
children under 6 years of age	consult a physician

**Other information**


- each rounded **tablespoon** contains: calcium 85 mg; potassium 125 mg
- each rounded **tablespoon** contributes 27 calories from maltodextrin
- store at room temperature
- protect contents from humidity
- keep tightly closed

**Inactive ingredients** anhydrous citric acid, FD&C Yellow No. 6 Lake, flavor, maltodextrin, potassium citrate, silica, sucralose, tricalcium phosphate

**Questions or comments? 1-866-467-2748**

\*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Citrucel® Powder.

Manufactured by:  
Raritan Pharmaceuticals  
8 Joanna Court,  
East Brunswick, NJ 08816



3 68163 51716 3

LOT: 5176DCLR

THIS IS TO DESIGNATE  
NON VARMISH AREA.  
DO NOT PRINT.

EXP:

## DRX CHOICE FIBER THERAPY

methylcellulose powder, for solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68163-517
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>METHYLCELLULOSE (4000 MPA.S)</b> (UNII: MRJ667KA5E) (METHYLCELLULOSE (4000 MPA.S) - UNII:MRJ667KA5E)	METHYLCELLULOSE (4000 MPA.S)	2 g in 10.2 g

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>POTASSIUM CITRATE</b> (UNII: EE90ONI6FF)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TRICALCIUM PHOSPHATE</b> (UNII: K4C08XP666)	

**Product Characteristics**

<b>Color</b>	ORANGE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68163-517-16	479 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/18/2022	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	07/18/2022	

**Labeler** - RARITAN PHARMACEUTICALS INC (127602287)

Revised: 9/2023

RARITAN PHARMACEUTICALS INC