DOCOSANOL- docosanol cream Chain Drug Consortium, LLC

Docosanol Cream, 10%

Active ingredient

Docosanol 10%

Purpose

Cold sore/fever blister treatment

Uses

- treats cold sores/fever blisters on the face or lips
- shortens healing time and duration of symptoms:
 - tingling, pain, burning, and/or itching

Warnings

Allergy Alert

This product may cause a severe allergic reaction. Symptoms may include: • hives • facial swelling • wheezing/difficulty breathing • shock • rash

If an allergic reaction occurs, stop use and seek medical help right away.

For external use only

Do not use

• if you are allergic to any ingredient in this product

When using this product

- apply only to the affected areas
- do not use in or near the eyes
- avoid applying directly inside your mouth
- do not share this product with anyone. This may spread the infection.

Stop use and ask a doctor if

• your cold sore gets worse or the cold sore is not healed within 10 days

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years or over:
 - wash hands before and after applying cream
 - apply to affected area on the face or lips at the first sign of cold sore/fever blister (tingle).
 - early treatment ensures the best results
 - rub in gently but completely
 - use 5 times a day until healed
- children under 12 years: ask a doctor

Other information

- store at 20° to 25°C (68° to 77°F)
- do not freeze

Inactive ingredients

benzyl alcohol, light mineral oil, propylene glycol, purified water, sucrose distearate, sucrose stearate

Questions?

1-888-838-2872 between 9 am and 5 pm ET, Monday-Friday.

Docosanol Cream, 10%

Thank you for purchasing Docosanol Cream. We hope that since Docosanol Cream shortens the healing time of cold sores, it helps you look and feel better faster. Please follow insert directions for proper use and refer to the information below.

You are not alone

An estimated one-in-five Americans is affected by cold sores each year. Most people who experience them get two to three cold sore outbreaks each year. Just before and during an outbreak, the cold sore infection can be easily passed on to others. It is especially important to be aware of early warning signs that a cold sore is coming, and to be sure to avoid close physical contact when you have one.

Getting to the Root of the Problem

A blister, or cluster of blisters around your mouth, is caused by the cold sore virus. After the cold sore has healed, the virus stays dormant in your body but can be reactivated by everyday triggers like stress, sunlight, extreme hot or cold weather, colds and flu, lack of sleep, or menstruation.

The Sooner, the Better

A typical cold sore outbreak can last as long as seven to ten days. The sooner you begin using Docosanol Cream, the more effective the medicine will be. So for best results, use Docosanol Cream at the very start of the cold sore - when you feel the first sign of the tingle, redness, bump or itch. Docosanol Cream is different from other over-the-counter cold sore products. Only Docosanol Cream contains 10% docosanol, which helps to shorten the healing time of cold sores and the duration of symptoms including tingling, pain, itching and burning.

Frequently Asked Questions:

How is Docosanol Cream different from the over-the-counter cold sore remedies I have already been using?

Docosanol Cream is a non-prescription cold sore medicine that can actually shorten healing time and the duration of symptoms. The sooner you start using Docosanol Cream, the sooner you can begin healing your cold sore.

Who can use Docosanol Cream?

Adults and children 12 years and over can use Docosanol Cream.

How often can I apply Docosanol Cream?

Docosanol Cream should be used five times a day until the cold sore is healed. Apply it directly to the cold sore at the first sign of a tingle, redness, bump or itch for best results.

Can I apply cosmetics on top of Docosanol Cream?

Yes, cosmetics such as lipstick may be applied over Docosanol Cream. However, use a separate applicator, like a cotton swab, to apply cosmetics over an unhealed cold sore to avoid spreading the infection. For best results, remove any cosmetics prior to applying Docosanol Cream.

Made in the USA Distributed by: Pharmacy Value Alliance, LLC 407 Lancaster Avenue, Wayne, PA 19087 Revised – March 2016

PRINCIPAL DISPLAY PANEL

Premier Value[®]

COMPARE TO THE ACTIVE INGREDIENT IN ABREVA®**

Cold Sore / Fever Blister Treatment


Docosanol Cream, 10%


Non-prescription Medicine to Shorten Healing Time*

Cold Sore Treatment

For Topical Use Only

Net Wt. 2g (.07 oz)





Active Ingredient/Active Moiety

Ingredient Name					Basis of Strength		Strength
DOCOSANOL (UNII:	9G10E216XY) ([OCOSANOL - UNII:9G10E216XY)			DOCOSANOL		0 mg in 1 g
Inactive Ingree	dients						
Ingredient Name						Stre	ength
BENZYL ALCOHOL (UNII: LKG8494WBH)							
LIGHT MINERAL OI	L (UNII: N6K5787	QVP)					
PROPYLENE GLYCO	DL (UNII: 6DC9Q	167V3)					
WATER (UNII: 059QF	F0KO0R)						
SUCROSE DISTEAR	ATE (UNII: 33X4	X4B90S)					
SUCROSE STEARAT	FE (UNII: 274KWC	0050M)					
Product Characteristics							
Color		white	Score				
Shape			Size				
Flavor			Imprint Code				
Contains							
Packaging							
# Item Code	Package Description			Mar	Marketing Start Ma Date		ting End ate
1 NDC:68016-052- 07	1 in 1 PACKAGE			04/02/2019			
1	2 g in 1 TUBE; Type 0: Not a Combination Product						
Marketing I	nformatio	on					
Marketing I Marketing Category		O N on Number or Citation	Monograph	ו M	arketing Start Date		eting End Date

Labeler - Chain Drug Consortium, LLC (101668460)

Revised: 12/2023

Chain Drug Consortium, LLC