

**GLYCERIN- glycerin suppository suppository**  
**MAJOR® PHARMACEUTICALS**

-----  
**MAJOR® Children's Glycerin**

**Drug Facts**

***Active Ingredient***

Glycerin 1.2 g

***Purpose***

Hyperosmotic Laxative

**Uses**

For relief of occasional constipation. This product usually produces a bowel movement in 1/4 to 1 hour.

***Warnings***

**For rectal use only.**

**Do not use**

laxative products when abdominal pain, nausea or vomiting are present unless directed by a doctor

**Ask a doctor before using any laxative if child has**

- abdominal pain, nausea or vomiting.
- a sudden change in bowel habits lasting more than 2 weeks.
- already used a laxative for more than 1 week.

**When using this product**

rectal discomfort or burning sensation may occur

**Stop use and consult a doctor if child has**

- rectal bleeding
- no bowel movement within 1 hour of using this product

This may indicate a serious condition.

**Keep out of reach of children.**

**If swallowed, get medical help or contact a Poison Control Center right**

**immediately (1-800-222-1222).**

## **Directions**

Single Daily dosage: 1 suppository or as directed by a doctor

Children 2 to 6 years:

- tear wrapper prior to immediate use
- remove the secondary wrapper before inserting into the rectum as follows:
  - hold suppository with pointed end up
  - carefully separate tabs with fingernail
  - gently pull apart by pulling tabs down on both sides
  - remove suppository from wrapper
  - insert one suppository well up into the rectum
- Children under 2 years of age: ask a doctor

## ***Other information***

- Only remove overwrap prior to immediate use
- Keep from excessive heat
- Do not use if overwrap is torn

## ***Inactive ingredients***

polyethylene glycol 400, sodium stearate

## ***Questions?***

**(855) 879-4368**

## **PRINCIPAL DISPLAY PANEL**

**MAJOR®** NDC 0904-7332-17

**Children's Glycerin**

**Suppositories**

**1.2 g**

**Laxative**

For fast, gentle relief of occasional constipation

Unit Dose

25 Individually Wrapped Child Size Suppositories

**MAJOR<sup>®</sup>**

NDC 0904-7332-17

# Children's Glycerin Suppositories

**1.2 g**

**Laxative**

**For fast, gentle relief of occasional constipation**

**Unit Dose**

**25 Individually Wrapped Child Size Suppositories**

**PRINCIPAL DISPLAY PANEL**

**MAJOR<sup>®</sup>** NDC 0904-7332-01

**Children's Glycerin**

**Suppositories**

**1.2 g**


**Laxative**

For fast, gentle relief of occasional constipation

DO NOT OPEN UNTIL IMMEDIATE USE

Unit Dose

1 Individually Wrapped Child Size Suppository



## GLYCERIN

glycerin suppository suppository

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-7332
Route of Administration	RECTAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>GLYCERIN</b> (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)	GLYCERIN	1.2 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>SODIUM STEARATE</b> (UNII: QU7E2XA9TG)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0904-7332-17	25 in 1 CARTON	07/13/2024	
1	NDC:0904-7332-01	1 in 1 POUCH; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M007	07/13/2024	

**Labeler** - MAJOR® PHARMACEUTICALS (191427277)

Revised: 7/2024

MAJOR® PHARMACEUTICALS