

ASPIRIN- aspirin tablet
Geri-Care Pharmaceuticals, Corp

Geri Care 44-249

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- pain and fever of colds
- headache
- muscle pain
- minor pain of arthritis
- toothache
- menstrual pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - feel faint
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredient

corn starch

Questions or comments?

1-800-540-3765

Principal Display Panel

NDC 57896-901-01

GERI CARE®

**ASPIRIN
325 mg (NSAID)**

PAIN RELIEVER / FEVER REDUCER

REGULAR STRENGTH

Compare to active ingredient
in **BAYER®** Aspirin **

100 TABLETS

Actual Size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING**

**This product is not manufactured or distributed by Bayer AG, owner
of the registered trademark Bayer® Aspirin.

50844

REV0122D24912

DISTRIBUTED BY: **GERI-CARE PHARMACEUTICAL CORP.**
1295 Towbin Ave., Lakewood, NJ 08701

ASPIRIN

325 mg (NSAID)

PAIN RELIEVER / FEVER REDUCER
REGULAR STRENGTH



Actual Size

100 Tablets

Compare to active ingredient
in **BAYER® Aspirin****



3 5789690101 1

**This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Aspirin.

DISTRIBUTED BY: GERI-CARE PHARMACEUTICAL CORP.
1295 Towbin Ave., Lakewood, NJ 08701

TAMPER EVIDENCE: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves:
 - pain and fever of colds
 - headache
 - muscle pain
 - minor pain of arthritis
 - toothache
 - menstrual pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

PEEL HERE FOR MORE DRUG FACTS

No print/No varnish
Lot & Exp date

Adhesive Only

STOP PEELING

Questions or comments? 1-800-540-3765

Inactive ingredient corn starch

Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

These could be signs of a serious condition:

- redness or swelling is present
- new symptoms occur

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - feel faint
 - ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Geri Care 44-249

ASPIRIN

aspirin tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:57896-901

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;249;ASPIRIN
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-901-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/1993	
2	NDC:57896-901-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/1993	
3	NDC:57896-901-27	250 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/1993	08/28/2020

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/04/1993	

Labeler - Geri-Care Pharmaceuticals, Corp (611196254)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(57896-901)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(57896-901)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(57896-901)

Revised: 7/2024

Geri-Care Pharmaceuticals, Corp