

**ASPIRIN- aspirin tablet**  
**Geri-Care Pharmaceuticals, Corp**

-----  
**Geri Care 44-249**

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves:

- pain and fever of colds
- headache
- muscle pain
- minor pain of arthritis
- toothache
- menstrual pain

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - feel faint
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

***Other information***

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)

***Inactive ingredient***

corn starch

***Questions or comments?***

**1-800-540-3765**

***Principal Display Panel***

NDC 57896-901-01

**GERICARE**

**ASPIRIN**

**325 mg (NSAID)**

**PAIN RELIEVER / FEVER REDUCER**

REGULAR STRENGTH

Compare to active ingredient  
in **BAYER®** Aspirin \*\*

**100 TABLETS**

Actual Size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY  
SEAL UNDER CAP IS BROKEN OR MISSING**


50844

REV0122D24912

\*\*This product is not manufactured or distributed by Bayer AG, owner  
of the registered trademark Bayer® Aspirin.

DISTRIBUTED BY: **GERI-CARE PHARMACEUTICAL CORP.**

1295 Towbin Ave., Lakewood, NJ 08701




NDC 57896-901-01

# ASPIRIN

## 325 mg (NSAID)


PAIN RELIEVER / FEVER REDUCER  
REGULAR STRENGTH



Actual Size

Compare to active ingredient  
in **BAYER® Aspirin™** \*\*

### 100 Tablets



3 57896 90101 1

\*\*\*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Aspirin.

REV012D24912

DISTRIBUTED BY: GERI-CARE PHARMACEUTICAL CORP.  
1295 Towbin Ave., Lakewood, NJ 08701

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 325 mg (NSAID)\* ..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves: ■ pain and fever of colds  
■ headache ■ muscle pain ■ minor pain of arthritis  
■ toothache ■ menstrual pain

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ facial swelling ■ shock ■ hives ■ asthma (wheezing)  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Drug Facts (continued)**  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer  
■ if you have ever had an allergic reaction to this product or any of its ingredients  
**Ask a doctor before use if** ■ you have asthma  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic  
**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ gout ■ diabetes ■ arthritis  
**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better ■ feel faint  
■ ringing in the ears or a loss of hearing occurs  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
These could be signs of a serious condition.  
**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than directed  
■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor  
■ children under 12 years: do not use unless directed by a doctor

**Other information** ■ use by expiration date on package  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredient** corn starch

**Questions or comments?** 1-800-540-3765

STOP PEELING

Adhesive Only

No print/No varnish  
Lot & Exp date

**PEEL HERE FOR MORE DRUG FACTS**

Geri Care 44-249

## ASPIRIN

aspirin tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-901
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;249;ASPIRIN
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-901-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/1993	
2	NDC:57896-901-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/1993	
3	NDC:57896-901-27	250 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/1993	08/28/2020

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/04/1993	

**Labeler** - Geri-Care Pharmaceuticals, Corp (611196254)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(57896-901)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(57896-901)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(57896-901)

Revised: 7/2023

Geri-Care Pharmaceuticals, Corp