

**TARON FORTE- iron, calcium ascorbate, cyanocobalamin, and folic acid capsule**

**Trigen Laboratories, LLC**

-----  
**Taron Forte  
Capsules**

**SUPPLEMENT FACTS**

<b>Supplement Facts</b>		
Serving Size: 1 Capsule		
Servings Per Container: 100		
	Amount Per Serving	% DV Adults Only
Vitamin C (as Calcium Ascorbate and Calcium Threonate)	60 mg	67%
Folate	1667 mcg DFE (1000 mcg Folic Acid)	417%
Vitamin B <sub>12</sub> (as Cyanocobalamin)	25 mcg	1042%
Iron (from Ferrous Bisglycinate Chelate and Polysaccharide Iron)	150 mg	833%

**Other Ingredients:** Orange Capsule (Gelatin, Titanium Dioxide, FD&C Yellow #6, FD&C Red #3), Black Ink (Propylene Glycol, FD&C Blue #1 Lake, FD&C Yellow #6 Lake, FD&C Red #40 Lake, Shellac), Microcrystalline Cellulose and Magnesium Stearate.

Taron Forte Multivitamin/Mineral with Iron capsules are indicated for the distinctive nutritional requirements of persons being treated for iron deficiencies by a physician.

**CONTRAINDICATIONS**

Taron Forte Capsules are contraindicated in patients with a known hypersensitivity to any of the components of this product.

Hemochromatosis and hemosiderosis are contraindications to iron therapy.

## WARNINGS

**WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. KEEP THIS PRODUCT OUT OF REACH OF CHILDREN. In case of accidental overdose, call a doctor or poison control center immediately.**

## PRECAUTIONS

**General:** The type of anemia and the underlying cause or causes should be determined before starting therapy with Taron Forte Capsules. Since the anemia may be a result of a systemic disturbance, such as recurrent blood loss, the underlying cause or causes should be corrected, if possible.

Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where B<sub>12</sub> is deficient.

Folic acid in doses above 0.1 mg daily may obscure pernicious anemia assessment, such that hematologic remission can occur while neurological manifestations remain progressive.

**Pediatric Use:** Safety and effectiveness of this product have not been established in pediatric patients.

## ADVERSE REACTIONS

Adverse reactions with iron therapy may include constipation, diarrhea, nausea, vomiting, dark stools, and abdominal pain. Adverse reactions with iron therapy are usually transient. Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

The clinical cause of acute iron overdosage can be variable. Initial symptoms may include: abdominal pain, nausea, vomiting, diarrhea, tarry stools, melena, hematemesis, hypotension, tachycardia, metabolic acidosis, hyperglycemia, dehydration, drowsiness, pallor, cyanosis, lassitude, seizures, shock, and coma.

## DESCRIPTION

Taron Forte Capsules are orange, imprinted with "TL042".

## DIRECTIONS FOR USE

One (1) capsule daily or as directed by a physician.

## HOW SUPPLIED

Taron Forte Capsules are packaged in child-resistant bottles of 100 capsules.

**PRODUCT CODE 13811-042-10.**

**STORAGE**

Store at 20°-25°C (68°-77°F), excursion permitted to 15°-30°C (59°-86°F) [See USP Controlled Room Temperature]. Protect from light and moisture. Dispense in a tight, light-resistant container with a child-resistant closure as defined in the USP/NF.

**KEEP OUT OF REACH OF CHILDREN.**

**For use on the order of a healthcare practitioner.**

**Call your doctor about side effects. To report side effects, call Trigen Laboratories at 1-770-509-4500 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**PLR-TARONFORTE-00001-1**

**Rev. 12/2021**

**Manufactured for:**

**Trigen Laboratories, LLC**

**Alpharetta, GA 30005**



**PRINCIPAL DISPLAY PANEL - 100 Capsule Bottle Label**

**13811-042-10**

**100 Capsules**

**Taron Forte  
CAPSULES**

**TRIGEN  
LABORATORIES**

STORAGE: Store at 20°-25°C (68°-77°F), excursions permitted to 15°-30°C (59°-86°F) [See USP Controlled Room Temperature]. Protect from light and moisture. Dispense in a tight, light-resistant container with a child-resistant closure as defined in the USP/NF.

13811-042-10  
100 Capsules

**WARNING:** Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. **KEEP THIS PRODUCT OUT OF REACH OF CHILDREN.** In case of accidental overdose, call a doctor or poison control center immediately.

Call your doctor about side effects. To report side effects, call Trigen Laboratories at 1-770-509-4500 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

**KEEP OUT OF THE REACH OF CHILDREN.**

For use on the order of a healthcare practitioner.

Manufactured for:  
Trigen Laboratories, LLC  
Alpharetta, GA 30005

PLR-TARONFORTE-00001-1

Rev. 12/2021

**Taron Forte**

CAPSULES

IRON SUPPLEMENT  
MULTIVITAMIN/MINERAL

**TRIGEN**  
LABORATORIES

Lot:  
Exp:



DESCRIPTION  
Taron Forte Capsules are orange, imprinted with "T042".

**Supplement Facts**

	Amount Per Serving	% DV Adults Only
Vitamin C (as Calcium Ascorbate and Calcium Threonate)	40 mg	67%
Folate	1667 mcg DFE (1000 mcg Folic Acid)	417%
Vitamin B12 (as Cyanocobalamin)	25 mcg	1041%
Iron (Iron Ferrous Bisglycinate Chelate and Polysaccharide Iron)	150 mg	833%

Other Ingredients: Orange Capsule (Gelatin, Titanium Dioxide, FD&C Yellow #6, FD&C Red #3), Black Ink (Polyethylene Glycol, FD&C Blue #1 Lake, FD&C Yellow #6 Lake, FD&C Red #40 Lake, Shellac), Microcrystalline Cellulose and Magnesium Stearate.

Taron Forte Multivitamin/Mineral with Iron capsules are indicated for the distinctive nutritional requirements of persons being treated for iron deficiencies by a physician.

**DIRECTIONS FOR USE**

Use (1) capsule daily or as directed by a physician.

Red label for prescribing and product information.

**TARON FORTE**

iron, calcium ascorbate, cyanocobalamin, and folic acid capsule

**Product Information**

<b>Product Type</b>	DIETARY SUPPLEMENT	<b>Item Code (Source)</b>	NHRIC:13811-042
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>IRON</b> (UNII: E1UOL152H7) (IRON - UNII:E1UOL152H7)	IRON	150 mg
<b>CALCIUM ASCORBATE</b> (UNII: 183E4W213W) (ASCORBIC ACID - UNII:PQ6CK8PD0R)	ASCORBIC ACID	60 mg
<b>CYANOCOBALAMIN</b> (UNII: P6YC3EG204) (CYANOCOBALAMIN - UNII:P6YC3EG204)	CYANOCOBALAMIN	25 ug
<b>FOLIC ACID</b> (UNII: 935E97BOY8) (FOLIC ACID - UNII:935E97BOY8)	FOLIC ACID	1 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>FD&amp;C RED NO. 3</b> (UNII: PN2ZH5LOQY)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>SHELLAC</b> (UNII: 46N107B71O)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NHRIC:13811-042-10	100 in 1 BOTTLE, PLASTIC		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
DIETARY SUPPLEMENT		05/01/2009	

### Supplement Facts

**Serving Size :** **Serving per Container :**

	Amount Per Serving	% Daily Value
color		
imprint		
scoring	1	
shape		
size (solid drugs)	22 mm	

**Labeler -** Trigen Laboratories, LLC (830479668)

Revised: 12/2021

Trigen Laboratories, LLC