

**COLD AND FLU DAY AND NIGHT, SEVERE- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, guaifenesin, phenylephrine hcl
ARMY AND AIR FORCE EXCHANGE SERVICE**

Exchange Select 44-503A473-08

Active ingredients (in each caplet) (Daytime Cold & Flu Severe)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Active ingredients (in each caplet) (Nighttime Cold & Flu Severe)

Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Antihistamine
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - headache
 - sore throat
 - nasal congestion
 - minor aches and pains
 - sinus congestion and pressure
 - sneezing and runny nose (**Nighttime only**)
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive (**Daytime only**)

- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- heart disease
- glaucoma (**Nighttime only**)
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- high blood pressure
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a breathing problem such as emphysema or chronic bronchitis (**Nighttime only**)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

- taking sedatives or tranquilizers (**Nighttime only**)

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children (**Nighttime only**)
- marked drowsiness may occur (**Nighttime only**)
- avoid alcoholic beverages (**Nighttime only**)
- use caution when driving a motor vehicle or operating machinery (**Nighttime only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Nighttime only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Do not take DAYTIME and NIGHTTIME products at the same time.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole – do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients (Daytime only)

corn starch, crospovidone, D&C yellow #10 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Inactive ingredients (Nighttime only)

corn starch, crospovidone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal display panel

exchange ✓ **select**™

Compare To The Active Ingredients of
Tylenol® COLD + FLU SEVERE Day & Night*

DAY & NIGHT
COLD & FLU
SEVERE

Acetaminophen Pain Reliever/Fever Reducer Dextromethorphan HBr Cough Suppressant Guaifenesin Expectorant Phenylephrine HCl Nasal Decongestant Actual Size 16 Caplets	Acetaminophen Pain Reliever/Fever Reducer Chlorpheniramine maleate Antihistamine Dextromethorphan HBr Cough Suppressant Phenylephrine HCl Nasal Decongestant Actual Size 8 Caplets
---	---

24 TOTAL CAPLETS

Do Not Take Daytime and Nighttime Products at the Same Time.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

*This product is not manufactured or distributed by
Johnson & Johnson Corporation, owner of the registered
trademark Tylenol® COLD + FLU SEVERE Day & Night.
50844 REV0922C50347308

“SATISFACTION GUARANTEED OR YOUR MONEY BACK”

Manufactured For Your Military Exchanges

Distributed by: LNK International, Inc.,
Hauppauge, NY 11788 1-800-426-9391

exchange select™ **DAY & NIGHT COLD & FLU SEVERE**

exchange select™ Compare To The Active Ingredients of Tylenol® COLD + FLU SEVERE Day & Night™

DAY & NIGHT COLD & FLU SEVERE

Acetaminophen  Pain Reliever/Fever Reducer
Dextromethorphan HBr
Cough Suppressant
Guaifenesin
Expectorant
Phenylephrine HCl
Nasal Decongestant

Acetaminophen  Pain Reliever/Fever Reducer
Chlorpheniramine maleate
Antihistamine
Dextromethorphan HBr
Cough Suppressant
Phenylephrine HCl
Nasal Decongestant

16 Caplets  Actual Size

8 Caplets  Actual Size

24 TOTAL CAPLETS

quality value

Do Not Take Daytime and Nighttime Products at the Same Time.

PARENTS: Learn about these medicines at www.StopMedicineAbuse.org

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Tylenol® COLD + FLU SEVERE Day & Night. 58844 REV0922C50347308

"SATISFACTION GUARANTEED OR YOUR MONEY BACK"
Manufactured For Your Military Exchanges
Distributed by: LNK International, Inc.,
Hauppauge, NY 11788 1-800-426-9391

TAMPER EVIDENT DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts (continued)

Active ingredients (in each caplet) (Daytime Cold & Flu Severe)

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Guaifenesin 200 mg Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Active ingredients (in each caplet) (Nighttime Cold & Flu Severe)

Acetaminophen 325 mg Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg Antihistamine
Dextromethorphan HBr 10 mg Cough suppressant
Phenylephrine HCl 5 mg Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - headache
 - minor aches and pains
 - nasal congestion
 - throat sore
- sinus congestion and pressure
- stinging and itchy nose (irritation only)
- helps loosen phlegm (mucus) (functional secretions to make coughs more productive (daytime only))

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- more than 3,000 mg of acetaminophen every day while using this product
- 3 or more alcoholic drinks every day while using this product

Warnings (continued)

- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

PEEL HERE FOR MORE DRUG FACTS

Lot & Exp ONLY

6 14299 40133 4

B-0066E-503A47308R
REV0922C50347308

ADHESIVE AREA

Drug Facts (continued)

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- heart disease
- glaucoma (Nighttime only)
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- high blood pressure
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a breathing problem such as emphysema or chronic bronchitis (Nighttime only)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (Nighttime only)

Drug Facts (continued)

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children (Nighttime only)
- marked drowsiness may occur (Nighttime only)
- avoid alcoholic beverages (Nighttime only)
- use caution when driving a motor vehicle or operating machinery (Nighttime only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

ADHESIVE AREA

ADHESIVE AREA

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Do not take DAYTIME and NIGHTTIME products at the same time.

Directions

- do not take more than directed
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole – do not crush, chew, or dissolve
- do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- each caplet contains: sodium 3 mg
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Drug Facts (continued)

Inactive ingredients (Daytime only)

corn starch, croscopolone, D&C yellow #10 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Inactive ingredients (Nighttime only)

corn starch, croscopolone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments? 1-800-426-9391

58844
REV0922C50347308
R-0066E-503A473-080F

ADHESIVE AREA

Exchange Select 44-503A473-08

COLD AND FLU DAY AND NIGHT, SEVERE

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, guaifenesin, phenylephrine hcl kit

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:55301-547

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-547-08	2 in 1 CARTON	03/31/2023	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1		8
Part 2		4

Part 1 of 2

COLD AND FLU DAY, SEVERE

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

Product Information

Item Code (Source)	NDC:55301-903
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	19mm
Flavor	MENTHOL	Imprint Code	44;503
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2023	

Part 2 of 2

COLD AND FLU NIGHT, SEVERE

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl tablet, film coated

Product Information

Item Code (Source)	NDC:55301-473
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	

FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	blue	Score	no score
Shape	OVAL	Size	17mm
Flavor	MENTHOL	Imprint Code	44;473
Contains			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2023	

Labeler - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55301-547) , pack(55301-547)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55301-547)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55301-547)

Revised: 4/2024

ARMY AND AIR FORCE EXCHANGE SERVICE