

**TELMISARTAN- telmisartan tablet**  
**Proficient Rx LP**

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use TELMISARTAN TABLETS safely and effectively. See full prescribing information for TELMISARTAN TABLETS.

TELMISARTAN tablets for oral use

Initial U.S. Approval: 1998

**WARNING: FETAL TOXICITY**

*See full prescribing information for complete boxed warning.*

- **When pregnancy is detected, discontinue telmisartan as soon as possible (5.1)**
- **Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus (5.1)**

**RECENT MAJOR CHANGES**

Warnings and Precautions

Dual Blockade of the Renin-Angiotensin-Aldosterone System (5.6)

12/2014

**INDICATIONS AND USAGE**

Telmisartan is an angiotensin II receptor blocker (ARB) indicated for:

- Treatment of hypertension, to lower blood pressure. Lowering blood pressure reduces the risk of fatal and nonfatal cardiovascular events, primarily strokes and myocardial infarctions. (1.1)

**DOSAGE AND ADMINISTRATION**

- May be administered with or without food (2.1)

<b>Indication</b>	<b>Starting Dose</b>	<b>Dose Range</b>
Hypertension (2.1)	40 mg once once daily	40 to 80 mg once daily

**DOSAGE FORMS AND STRENGTHS**

- Tablets: 20mg, 40mg, 80mg (3)

**CONTRAINDICATIONS**

- Known hypersensitivity (e.g., anaphylaxis or angioedema) to telmisartan or any other component of this product (4)
- Do not co-administer aliskiren with telmisartan in patients with diabetes (4)

**WARNINGS AND PRECAUTIONS**

- Avoid fetal or neonatal exposure (5.1)
- Hypotension: Correct any volume or salt depletion before initiating therapy. Observe for signs and symptoms of hypotension (5.2)
- Monitor carefully in patients with impaired hepatic (5.4) or renal function (5.5)
- Avoid concomitant use of an ACE inhibitor and angiotensin receptor blocker (5.6)

**ADVERSE REACTIONS**

- Hypertension: The most common adverse events ( $\geq 1\%$ ) reported in hypertension trials are back pain, sinusitis, and diarrhea (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Ascend Laboratories, LLC, Parsippany, NJ 07054 at 1-877-272-7901 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

**DRUG INTERACTIONS**

- NSAID : Increased risk of renal impairment and loss of anti-hypertensive effect (7)
- Do not co-administer aliskiren with telmisartan in patients with diabetes (7)

----- **USE IN SPECIFIC POPULATIONS** -----

- Nursing Mothers: Choose to discontinue nursing or drug (8.3)
- Geriatric Patients: No overall difference in efficacy or safety vs younger patients, but greater sensitivity of some older individuals cannot be ruled out (8.5)

See 17 for **PATIENT COUNSELING INFORMATION**.

**Revised: 6/2020**

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**WARNING: FETAL TOXICITY**

**1 INDICATIONS & USAGE**

1.1 Hypertension

**2 DOSAGE & ADMINISTRATION**

2.1 Hypertension

**3 DOSAGE FORMS & STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

5.1 Fetal Toxicity

5.2 Hypotension

5.3 Hyperkalemia

5.4 Impaired Hepatic Function

5.5 Impaired Renal Function

5.6 Dual Blockade of the Renin-Angiotensin-Aldosterone System

**6 ADVERSE REACTIONS**

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

**7 DRUG INTERACTIONS**

**8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Insufficiency

**10 OVERDOSAGE**

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis & Mutagenesis & Impairment Of Fertility

13.3 Developmental Toxicity

**14 CLINICAL STUDIES**

14.1 Hypertension

**16 HOW SUPPLIED/STORAGE AND HANDLING**

**17 PATIENT COUNSELING INFORMATION**

17.1 Pregnancy

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## FULL PRESCRIBING INFORMATION

### WARNING: FETAL TOXICITY

- **When pregnancy is detected, discontinue telmisartan as soon as possible [see Warnings and Precautions (5.1)].**
- **Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus [see Warnings and Precautions (5.1)]**

## 1 INDICATIONS & USAGE

### 1.1 Hypertension

Telmisartan is indicated for the treatment of hypertension, to lower blood pressure. Lowering blood pressure reduces the risk of fatal and nonfatal cardiovascular events, primarily strokes and myocardial infarctions. These benefits have been seen in controlled trials of antihypertensive drugs from a wide variety of pharmacologic classes including the class to which this drug principally belongs.

Control of high blood pressure should be part of comprehensive cardiovascular risk management, including, as appropriate, lipid control, diabetes management, antithrombotic therapy, smoking cessation, exercise, and limited sodium intake. Many patients will require more than one drug to achieve blood pressure goals. For specific advice on goals and management, see published guidelines, such as those of the National High Blood Pressure Education Program's Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC).

Numerous antihypertensive drugs, from a variety of pharmacologic classes and with different mechanisms of action, have been shown in randomized controlled trials to reduce cardiovascular morbidity and mortality, and it can be concluded that it is blood pressure reduction, and not some other pharmacologic property of the drugs, that is largely responsible for those benefits. The largest and most consistent cardiovascular outcome benefit has been a reduction in the risk of stroke, but reductions in myocardial infarction and cardiovascular mortality also have been seen regularly.

Elevated systolic or diastolic pressure causes increased cardiovascular risk, and the absolute risk increase per mmHg is greater at higher blood pressures, so that even modest reductions of severe hypertension can provide substantial benefit. Relative risk reduction from blood pressure reduction is similar across populations with varying absolute risk, so the absolute benefit is greater in patients who are at higher risk independent of their hypertension (for example, patients with diabetes or hyperlipidemia), and such patients would be expected to benefit from more aggressive treatment to a lower blood pressure goal.

Some antihypertensive drugs have smaller blood pressure effects (as monotherapy) in black patients, and many antihypertensive drugs have additional approved indications and effects (e.g., on angina, heart failure, or diabetic kidney disease). These considerations may guide selection of therapy.

It may be used alone or in combination with other antihypertensive agents [see Clinical Studies (14.1)].

Use of telmisartan and an ACE inhibitor is not recommended [see Warnings and Precautions (5.6)].

## 2 DOSAGE & ADMINISTRATION

## 2.1 Hypertension

Dosage must be individualized. The usual starting dose of telmisartan tablets USP is 40 mg once a day. Blood pressure response is dose-related over the range of 20 to 80 mg [see *Clinical Studies (14.1)*].

Most of the antihypertensive effect is apparent within 2 weeks and maximal reduction is generally attained after 4 weeks. When additional blood pressure reduction beyond that achieved with 80 mg telmisartan is required, a diuretic may be added.

No initial dosage adjustment is necessary for elderly patients or patients with renal impairment, including those on hemodialysis. Patients on dialysis may develop orthostatic hypotension; their blood pressure should be closely monitored.

Telmisartan tablets USP may be administered with other antihypertensive agents.

Telmisartan tablets USP may be administered with or without food.

## 3 DOSAGE FORMS & STRENGTHS

- 20 mg, white to off-white, uncoated round shaped tablets, flat with beveled edges, plain on one side and debossed "036" on other side.
- 40 mg, white to off-white, uncoated oval shaped tablets, biconvex with beveled edges, plain on one side and debossed "037" on other side.
- 80 mg, white to off-white, uncoated oval shaped tablets, biconvex with beveled edges, plain on one side and debossed "038" on other side.

## 4 CONTRAINDICATIONS

Telmisartan is contraindicated in patients with known hypersensitivity (e.g., anaphylaxis or angioedema) to telmisartan or any other component of this product [see *Adverse Reactions (6.2)*].

Do not co-administer aliskiren with telmisartan in patients with diabetes. [see *Drug interactions(7)*].

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Fetal Toxicity

#### Pregnancy Category D

Use of drugs that act on the renin-angiotensin system during the second and third trimesters of pregnancy reduces fetal renal function and increases fetal and neonatal morbidity and death. Resulting oligohydramnios can be associated with fetal lung hypoplasia and skeletal deformations. Potential neonatal adverse effects include skull hypoplasia, anuria, hypotension, renal failure, and death. When pregnancy is detected, discontinue telmisartan as soon as possible [see *Use in Specific Populations (8.1)*].

### 5.2 Hypotension

In patients with an activated renin-angiotensin system, such as volume- or salt-depleted patients (e.g., those being treated with high doses of diuretics), symptomatic hypotension may occur after initiation of therapy with telmisartan. Either correct this condition prior to administration of telmisartan, or start treatment under close medical supervision with a reduced dose.

If hypotension does occur, the patient should be placed in the supine position and, if necessary, given an intravenous infusion of normal saline. A transient hypotensive response is not a contraindication to further treatment, which usually can be continued without difficulty once the blood pressure has stabilized.

### 5.3 Hyperkalemia

Hyperkalemia may occur in patients on ARBs, particularly in patients with advanced renal impairment, heart failure, on renal replacement therapy, or on potassium supplements, potassium-sparing diuretics, potassium-containing salt substitutes or other drugs that increase potassium levels. Consider periodic determinations of serum electrolytes to detect possible electrolyte imbalances, particularly in patients at risk.

### 5.4 Impaired Hepatic Function

As the majority of telmisartan is eliminated by biliary excretion, patients with biliary obstructive disorders or hepatic insufficiency can be expected to have reduced clearance. Initiate telmisartan at low doses and titrate slowly in these patients [*see Use in Specific Populations (8.6) and Clinical Pharmacology (12.3)*].

### 5.5 Impaired Renal Function

As a consequence of inhibiting the renin-angiotensin-aldosterone system, anticipate changes in renal function in susceptible individuals. In patients whose renal function may depend on the activity of the renin-angiotensin-aldosterone system (e.g., patients with severe congestive heart failure or renal dysfunction), treatment with angiotensin-converting enzyme (ACE) inhibitors and angiotensin receptor antagonists has been associated with oliguria and/or progressive azotemia and (rarely) with acute renal failure and/or death. Similar results have been reported with telmisartan [*see Clinical Pharmacology (12.3)*].

In studies of ACE inhibitors in patients with unilateral or bilateral renal artery stenosis, increases in serum creatinine or blood urea nitrogen were observed. There has been no long term use of telmisartan in patients with unilateral or bilateral renal artery stenosis, but anticipate an effect similar to that seen with ACE inhibitors.

### 5.6 Dual Blockade of the Renin-Angiotensin-Aldosterone System

Dual blockade of the RAS with angiotensin-receptor blockers, ACE inhibitors, or aliskiren is associated with increased risks of hypotension, hyperkalemia, and changes in renal function (including acute renal failure) compared to monotherapy.

Patients receiving the combination of telmisartan and ramipril (*in ONTARGET trial that enrolled 25,620 patients ≥55 years old with atherosclerotic disease or diabetes with endorgan damage – added for clarification*) did not obtain any additional benefit compared to monotherapy, but experienced an increase incidence of renal dysfunction (e.g., acute renal failure) compared with groups receiving telmisartan alone or ramipril alone.

Concomitant use of telmisartan and ramipril is not recommended.

In most patients no benefit has been associated with using two RAS inhibitors concomitantly. In general, avoid combined use of RAS inhibitors. Closely monitor blood pressure, renal function, and electrolytes in patients on telmisartan and other agents that affect the RAS.

Do not co-administer aliskiren with telmisartan in patients with diabetes. Avoid concomitant use of aliskiren with telmisartan in patients with renal impairment (GFR <60 mL/min/1.73 m<sup>2</sup>).

## 6 ADVERSE REACTIONS

The following adverse reaction is described elsewhere in labeling:

Renal dysfunction upon use with ramipril [*see Warnings and Precautions (5.6)*]

### 6.1 Clinical Trials Experience

Because clinical studies are conducted under widely varying conditions, adverse reactions rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

### *Hypertension*

Telmisartan has been evaluated for safety in more than 3700 patients, including 1900 treated for over 6 months and more than 1300 for over one year. Adverse experiences have generally been mild and transient in nature and have infrequently required discontinuation of therapy.

In placebo-controlled trials involving 1041 patients treated with various doses of telmisartan (20 to 160 mg) monotherapy for up to 12 weeks, the overall incidence of adverse events was similar to that in patients treated with placebo.

Adverse events occurring at an incidence of  $\geq 1\%$  in patients treated with telmisartan and at a greater rate than in patients treated with placebo, irrespective of their causal association, are presented in Table 1.

**Table 1 Adverse Events Occurring at an Incidence of  $\geq 1\%$  in Patients Treated with Telmisartan and at a Greater Rate Than Patients Treated with Placebo**

	<b>Telmisartan n=1455 %</b>	<b>Placebo n=380 %</b>
Upper respiratory tract infection	7	6
Back pain	3	1
Sinusitis	3	2
Diarrhea	3	2
Pharyngitis	1	0

In addition to the adverse events in the table, the following events occurred at a rate of  $\geq 1\%$  but were at least as frequent in the placebo group: influenza-like symptoms, dyspepsia, myalgia, urinary tract infection, abdominal pain, headache, dizziness, pain, fatigue, coughing, hypertension, chest pain, nausea, and peripheral edema. Discontinuation of therapy because of adverse events was required in 2.8% of 1455 patients treated with telmisartan tablets USP and 6.1% of 380 placebo patients in placebo-controlled clinical trials.

The incidence of adverse events was not dose-related and did not correlate with gender, age, or race of patients.

The incidence of cough occurring with telmisartan in 6 placebo-controlled trials was identical to that noted for placebo-treated patients (1.6%).

In addition to those listed above, adverse events that occurred in more than 0.3% of 3500 patients treated with telmisartan monotherapy in controlled or open trials are listed below. It cannot be determined whether these events were causally related to telmisartan tablets USP:

*Autonomic Nervous System:* impotence, increased sweating, flushing; *Body as a Whole:* allergy, fever, leg pain, malaise; *Cardiovascular:* palpitation, dependent edema, angina pectoris, tachycardia, leg edema, abnormal ECG; *CNS:* insomnia, somnolence, migraine, vertigo, paresthesia, involuntary muscle contractions, hypoesthesia; *Gastrointestinal:* flatulence, constipation, gastritis, vomiting, dry mouth, hemorrhoids, gastroenteritis, enteritis, gastroesophageal reflux, toothache, nonspecific gastrointestinal disorders; *Metabolic:* gout, hypercholesterolemia, diabetes mellitus; *Musculoskeletal:* arthritis, arthralgia, leg cramps; *Psychiatric:* anxiety, depression, nervousness; *Resistance Mechanism:* infection, fungal infection, abscess, otitis media; *Respiratory:* asthma, bronchitis, rhinitis, dyspnea, epistaxis; *Skin:* dermatitis, rash, eczema, pruritus; *Urinary:* micturition frequency, cystitis; *Vascular:* cerebrovascular disorder; and *Special Senses:* abnormal vision, conjunctivitis, tinnitus, earache.

During initial clinical studies, a single case of angioedema was reported (among a total of 3781 patients treated).

#### *Clinical Laboratory Findings*

In placebo-controlled clinical trials, clinically relevant changes in standard laboratory test parameters were rarely associated with administration of telmisartan tablets USP.

**Hemoglobin:** A greater than 2 g/dL decrease in hemoglobin was observed in 0.8% telmisartan patients compared with 0.3% placebo patients. No patients discontinued therapy because of anemia.

**Creatinine:** A 0.5 mg/dL rise or greater in creatinine was observed in 0.4% telmisartan patients compared with 0.3% placebo patients. One telmisartan-treated patient discontinued therapy because of increases in creatinine and blood urea nitrogen.

**Liver Enzymes:** Occasional elevations of liver chemistries occurred in patients treated with telmisartan; all marked elevations occurred at a higher frequency with placebo. No telmisartan-treated patients discontinued therapy because of abnormal hepatic function.

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during post-approval use of telmisartan. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to estimate reliably their frequency or establish a causal relationship to drug exposure. Decisions to include these reactions in labeling are typically based on one or more of the following factors: (1) seriousness of the reaction, (2) frequency of reporting, or (3) strength of causal connection to telmisartan.

The most frequent spontaneously reported events include: headache, dizziness, asthenia, coughing, nausea, fatigue, weakness, edema, face edema, lower limb edema, angioneurotic edema, urticaria, hypersensitivity, sweating increased, erythema, chest pain, atrial fibrillation, congestive heart failure, myocardial infarction, blood pressure increased, hypertension aggravated, hypotension (including postural hypotension), hyperkalemia, syncope, dyspepsia, diarrhea, pain, urinary tract infection, erectile dysfunction, back pain, abdominal pain, muscle cramps (including leg cramps), myalgia, bradycardia, eosinophilia, thrombocytopenia, uric acid increased, abnormal hepatic function/liver disorder, renal impairment including acute renal failure, anemia, increased CPK, anaphylactic reaction, tendon pain (including tendonitis, tenosynovitis), drug eruption (toxic skin eruption mostly reported as toxicoderma, rash, and urticaria), hypoglycemia (in diabetic patients), and angioedema (with fatal outcome).

Rare cases of rhabdomyolysis have been reported in patients receiving angiotensin II receptor blockers, including telmisartan.

## **7 DRUG INTERACTIONS**

**Aliskiren:** Do not co-administer aliskiren with telmisartan in patients with diabetes. Avoid use of aliskiren with telmisartan in patients with renal impairment (GFR <60 mL/min).

**Digoxin:** When telmisartan was co-administered with digoxin, median increases in digoxin peak plasma concentration (49%) and in trough concentration (20%) were observed. Therefore, monitor digoxin levels when initiating, adjusting, and discontinuing telmisartan for the purpose of keeping the digoxin level within the therapeutic range.

**Lithium:** Reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with angiotensin II receptor antagonists including telmisartan. Therefore, monitor serum lithium levels during concomitant use.

**Non-Steroidal Anti-Inflammatory Agents including Selective Cyclooxygenase-2 Inhibitors (COX-2 Inhibitors):** In patients who are elderly, volume-depleted (including those on diuretic therapy), or with compromised renal function, co-administration of NSAIDs, including selective COX-2 inhibitors, with angiotensin II receptor antagonists, including telmisartan, may result in deterioration of renal function,

including possible acute renal failure. These effects are usually reversible. Monitor renal function periodically in patients receiving telmisartan and NSAID therapy.

The antihypertensive effect of angiotensin II receptor antagonists, including telmisartan may be attenuated by NSAIDs including selective COX-2 inhibitors.

*Ramipril and Ramiprilat:* Co-administration of telmisartan 80 mg once daily and ramipril 10 mg once daily to healthy subjects increases steady-state C<sub>max</sub> and AUC of ramipril 2.3- and 2.1-fold, respectively, and C<sub>max</sub> and AUC of ramiprilat 2.4- and 1.5-fold, respectively. In contrast, C<sub>max</sub> and AUC of telmisartan decrease by 31% and 16%, respectively. When co-administering telmisartan and ramipril, the response may be greater because of the possibly additive pharmacodynamic effects of the combined drugs, and also because of the increased exposure to ramipril and ramiprilat in the presence of telmisartan. Concomitant use of telmisartan and ramipril is not recommended.

*Other Drugs:* Co-administration of telmisartan did not result in a clinically significant interaction with acetaminophen, amlodipine, glyburide, simvastatin, hydrochlorothiazide, warfarin, or ibuprofen. Telmisartan is not metabolized by the cytochrome P450 system and had no effects in vitro on cytochrome P450 enzymes, except for some inhibition of CYP2C19. Telmisartan is not expected to interact with drugs that inhibit cytochrome P450 enzymes; it is also not expected to interact with drugs metabolized by cytochrome P450 enzymes, except for possible inhibition of the metabolism of drugs metabolized by CYP2C19.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

*Pregnancy Category D. [See Warnings and Precautions (5.1)].*

Use of drugs that act on the renin-angiotensin system during the second and third trimesters of pregnancy reduces fetal renal function and increases fetal and neonatal morbidity and death. Resulting oligohydramnios can be associated with fetal lung hypoplasia and skeletal deformations. Potential neonatal adverse effects include skull hypoplasia, anuria, hypotension, renal failure, and death. When pregnancy is detected, discontinue telmisartan as soon as possible. These adverse outcomes are usually associated with use of these drugs in the second and third trimester of pregnancy. Most epidemiologic studies examining fetal abnormalities after exposure to antihypertensive use in the first trimester have not distinguished drugs affecting the renin-angiotensin system from other antihypertensive agents. Appropriate management of maternal hypertension during pregnancy is important to optimize outcomes for both mother and fetus.

In the unusual case that there is no appropriate alternative to therapy with drugs affecting the renin-angiotensin system for a particular patient, apprise the mother of the potential risk to the fetus. Perform serial ultrasound examinations to assess the intra-amniotic environment. If oligohydramnios is observed, discontinue telmisartan, unless it is considered lifesaving for the mother. Fetal testing may be appropriate, based on the week of pregnancy. Patients and physicians should be aware, however, that oligohydramnios may not appear until after the fetus has sustained irreversible injury. Closely observe infants with histories of in utero exposure to telmisartan for hypotension, oliguria, and hyperkalemia [*see Use in Specific Populations (8.4)*].

### **8.3 Nursing Mothers**

It is not known whether telmisartan is excreted in human milk, but telmisartan was shown to be present in the milk of lactating rats. Because of the potential for adverse effects on the nursing infant, decide whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

### **8.4 Pediatric Use**



## Neonates with a history of in utero exposure to telmisartan tablets USP:

If oliguria or hypotension occurs, direct attention toward support of blood pressure and renal perfusion. Exchange transfusions or dialysis may be required as a means of reversing hypotension and/or substituting for disordered renal function.

Safety and effectiveness in pediatric patients have not been established [see *Clinical Pharmacology (12.3)*].

### **8.5 Geriatric Use**

Of the total number of patients receiving telmisartan tablets USP in hypertension clinical studies, 551 (19%) were 65 to 74 years of age and 130 (4%) were 75 years or older. No overall differences in effectiveness and safety were observed in these patients compared to younger patients and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

### **8.6 Hepatic Insufficiency**

Monitor carefully and uptitrate slowly in patients with biliary obstructive disorders or hepatic insufficiency [see *Warnings and Precautions (5.4)*].


## **10 OVERDOSAGE**

Limited data are available with regard to overdosage in humans. The most likely manifestation of overdosage with telmisartan tablets USP would be hypotension, dizziness and tachycardia; bradycardia could occur from parasympathetic (vagal) stimulation. If symptomatic hypotension should occur, supportive treatment should be instituted. Telmisartan is not removed by hemodialysis.

## **11 DESCRIPTION**

Telmisartan tablets USP is a non-peptide angiotensin II receptor (type AT<sub>1</sub>) antagonist.

Telmisartan is chemically described as 4'-[(1,4'-dimethyl-2'-propyl [2,6'-bi-1H-benzimidazol]-1'-yl)methyl]-[1,1'-biphenyl]-2-carboxylic acid. Its empirical formula is C<sub>33</sub>H<sub>30</sub>N<sub>4</sub>O<sub>2</sub>, its molecular weight is 514.63, and its structural formula is:



Telmisartan USP is a white to slightly yellowish solid. It is practically insoluble in water and in the pH range of 3 to 9, sparingly soluble in strong acid (except insoluble in hydrochloric acid), and soluble in strong base.

Telmisartan is available as tablets for oral administration, containing 20 mg, 40 mg or 80 mg of telmisartan USP. The tablets contain the following inactive ingredients: magnesium stearate, mannitol, meglumine, povidone, sodium hydroxide and sodium strearyl fumarate. Telmisartan tablets USP are hygroscopic and require protection from moisture.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Angiotensin II is formed from angiotensin I in a reaction catalyzed by angiotensin-converting enzyme (ACE, kininase II). Angiotensin II is the principal pressor agent of the reninangiotensin system, with effects that include vasoconstriction, stimulation of synthesis and release of aldosterone, cardiac stimulation, and renal reabsorption of sodium. Telmisartan blocks the vasoconstrictor and aldosterone-secreting effects of angiotensin II by selectively blocking the binding of angiotensin II to the AT1 receptor in many tissues, such as vascular smooth muscle and the adrenal gland. Its action is therefore independent of the pathways for angiotensin II synthesis.

There is also an AT2 receptor found in many tissues, but AT2 is not known to be associated with cardiovascular homeostasis. Telmisartan has much greater affinity (>3,000 fold) for the AT1 receptor than for the AT2 receptor.

Blockade of the renin-angiotensin system with ACE inhibitors, which inhibit the biosynthesis of angiotensin II from angiotensin I, is widely used in the treatment of hypertension. ACE inhibitors also inhibit the degradation of bradykinin, a reaction also catalyzed by ACE. Because telmisartan does not inhibit ACE (kininase II), it does not affect the response to bradykinin. Whether this difference has clinical relevance is not yet known. Telmisartan does not bind to or block other hormone receptors or ion channels known to be important in cardiovascular regulation.

Blockade of the angiotensin II receptor inhibits the negative regulatory feedback of angiotensin II on renin secretion, but the resulting increased plasma renin activity and angiotensin II circulating levels do not overcome the effect of telmisartan on blood pressure.

### **12.2 Pharmacodynamics**

In normal volunteers, a dose of telmisartan 80 mg inhibited the pressor response to an intravenous infusion of angiotensin II by about 90% at peak plasma concentrations with approximately 40% inhibition persisting for 24 hours.

Plasma concentration of angiotensin II and plasma renin activity (PRA) increased in a dose-dependent manner after single administration of telmisartan to healthy subjects and repeated administration to hypertensive patients. The once-daily administration of up to 80 mg telmisartan to healthy subjects did not influence plasma aldosterone concentrations. In multiple dose studies with hypertensive patients, there were no clinically significant changes in electrolytes (serum potassium or sodium), or in metabolic function (including serum levels of cholesterol, triglycerides, HDL, LDL, glucose, or uric acid).

In 30 hypertensive patients with normal renal function treated for 8 weeks with telmisartan 80 mg or telmisartan 80 mg in combination with hydrochlorothiazide 12.5 mg, there were no clinically significant changes from baseline in renal blood flow, glomerular filtration rate, filtration fraction, renovascular resistance, or creatinine clearance.

### **12.3 Pharmacokinetics**

Following oral administration, peak concentrations ( $C_{max}$ ) of telmisartan are reached in 0.5 to 1 hour

after dosing. Food slightly reduces the bioavailability of telmisartan, with a reduction in the area under the plasma concentration-time curve (AUC) of about 6% with the 40 mg tablet and about 20% after a 160 mg dose. The absolute bioavailability of telmisartan is dose dependent. At 40 and 160 mg the bioavailability was 42% and 58%, respectively. The pharmacokinetics of orally administered telmisartan are nonlinear over the dose range 20 to 160 mg, with greater than proportional increases of plasma concentrations ( $C_{max}$  and AUC) with increasing doses. Telmisartan shows bi-exponential decay kinetics with a terminal elimination half life of approximately 24 hours. Trough plasma concentrations of telmisartan with once daily dosing are about 10% to 25% of peak plasma concentrations. Telmisartan has an accumulation index in plasma of 1.5 to 2.0 upon repeated once daily dosing.

#### *Distribution*

Telmisartan is highly bound to plasma proteins (>99.5%), mainly albumin and  $\alpha_1$  – acid glycoprotein. Plasma protein binding is constant over the concentration range achieved with recommended doses. The volume of distribution for telmisartan is approximately 500 liters indicating additional tissue binding.

#### *Metabolism and Elimination*

Following either intravenous or oral administration of  $^{14}C$ -labeled telmisartan, most of the administered dose (>97%) was eliminated unchanged in feces via biliary excretion; only minute amounts were found in the urine (0.91% and 0.49% of total radioactivity, respectively).

Telmisartan is metabolized by conjugation to form a pharmacologically inactive acyl glucuronide; the glucuronide of the parent compound is the only metabolite that has been identified in human plasma and urine. After a single dose, the glucuronide represents approximately 11% of the measured radioactivity in plasma. The cytochrome P450 isoenzymes are not involved in the metabolism of telmisartan.

Total plasma clearance of telmisartan is >800 mL/min. Terminal half-life and total clearance appear to be independent of dose.

#### Specific Populations

##### Renal Insufficiency

No dosage adjustment is necessary in patients with decreased renal function. Telmisartan is not removed from blood by hemofiltration [see *Warnings and Precautions (5.5) and Dosage and Administration (2.1)*].

##### Hepatic Insufficiency

In patients with hepatic insufficiency, plasma concentrations of telmisartan are increased, and absolute bioavailability approaches 100% [see *Warnings and Precautions (5.4) and Use in Specific Populations (8.6)*].

##### Gender

Plasma concentrations of telmisartan are generally 2 to 3 times higher in females than in males. In clinical trials, however, no significant increases in blood pressure response or in the incidence of orthostatic hypotension were found in women. No dosage adjustment is necessary.

##### Geriatric Patients

The pharmacokinetics of telmisartan do not differ between the elderly and those younger than 65 years [see *Dosage and Administration (2.1)*].

##### Pediatric Patients

Telmisartan pharmacokinetics have not been investigated in patients <18 years of age.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis & Mutagenesis & Impairment Of Fertility**

There was no evidence of carcinogenicity when telmisartan was administered in the diet to mice and rats for up to 2 years. The highest doses administered to mice (1000 mg/kg/day) and rats (100 mg/kg/day) are, on a mg/m<sup>2</sup> basis, about 59 and 13 times, respectively, the maximum recommended human dose (MRHD) of telmisartan. These same doses have been shown to provide average systemic exposures to telmisartan >100 times and >25 times, respectively, the systemic exposure in humans receiving the MRHD (80 mg/day).

Genotoxicity assays did not reveal any telmisartan-related effects at either the gene or chromosome level. These assays included bacterial mutagenicity tests with *Salmonella* and *E. coli* (Ames), a gene mutation test with Chinese hamster V79 cells, a cytogenetic test with human lymphocytes, and a mouse micronucleus test.

No drug-related effects on the reproductive performance of male and female rats were noted at 100 mg/kg/day (the highest dose administered), about 13 times, on a mg/m<sup>2</sup> basis, the MRHD of telmisartan. This dose in the rat resulted in an average systemic exposure (telmisartan AUC as determined on day 6 of pregnancy) at least 50 times the average systemic exposure in humans at the MRHD (80 mg/day).

### **13.3 Developmental Toxicity**

There is no clinical experience with the use of telmisartan tablets USP in pregnant women. No teratogenic effects were observed when telmisartan was administered to pregnant rats at oral doses of up to 50 mg/kg/day and to pregnant rabbits at oral doses up to 45 mg/kg/day. In rabbits, embryoletality associated with maternal toxicity (reduced body weight gain and food consumption) was observed at 45 mg/kg/day [about 12 times the maximum recommended human dose (MRHD) of 80 mg on a mg/m<sup>2</sup> basis]. In rats, maternally toxic (reduction in body weight gain and food consumption) telmisartan doses of 15 mg/kg/day (about 1.9 times the MRHD on a mg/m<sup>2</sup> basis), administered during late gestation and lactation, were observed to produce adverse effects in neonates, including reduced viability, low birth weight, delayed maturation, and decreased weight gain. Telmisartan has been shown to be present in rat fetuses during late gestation and in rat milk. The no observed effect doses for developmental toxicity in rats and rabbits, 5 and 15 mg/kg/day, respectively, are about 0.64 and 3.7 times, on a mg/m<sup>2</sup> basis, the maximum recommended human dose of telmisartan (80 mg/day).

## **14 CLINICAL STUDIES**

### **14.1 Hypertension**

The antihypertensive effects of telmisartan have been demonstrated in six principal placebo controlled clinical trials, studying a range of 20 to 160 mg; one of these examined the antihypertensive effects of telmisartan and hydrochlorothiazide in combination. The studies involved a total of 1773 patients with mild to moderate hypertension (diastolic blood pressure of 95 to 114 mmHg), 1031 of whom were treated with telmisartan. Following once daily administration of telmisartan, the magnitude of blood pressure reduction from baseline after placebo subtraction was approximately (SBP/DBP) 6-8/6 mmHg for 20 mg, 9-13/6-8 mmHg for 40 mg, and 12-13/7-8 mmHg for 80 mg. Larger doses (up to 160 mg) did not appear to cause a further decrease in blood pressure.

Upon initiation of antihypertensive treatment with telmisartan, blood pressure was reduced after the first dose, with a maximal reduction by about 4 weeks. With cessation of treatment with telmisartan tablets USP, blood pressure gradually returned to baseline values over a period of several days to one week. During long term studies (without placebo control) the effect of telmisartan appeared to be maintained for up to at least one year. The antihypertensive effect of telmisartan is not influenced by patient age, gender, weight, or body mass index. Blood pressure response in black patients (usually a low-renin population) is noticeably less than that in Caucasian patients. This has been true for most, but not all, angiotensin II antagonists and ACE inhibitors.

In a controlled study, the addition of telmisartan to hydrochlorothiazide produced an additional dose-related reduction in blood pressure that was similar in magnitude to the reduction achieved with

telmisartan monotherapy. Hydrochlorothiazide also had an added blood pressure effect when added to telmisartan.

The onset of antihypertensive activity occurs within 3 hours after administration of a single oral dose. At doses of 20, 40, and 80 mg, the antihypertensive effect of once daily administration of telmisartan is maintained for the full 24-hour dose interval. With automated ambulatory blood pressure monitoring and conventional blood pressure measurements, the 24-hour trough-to peak ratio for 40 to 80 mg doses of telmisartan was 70 to 100% for both systolic and diastolic blood pressure. The incidence of symptomatic orthostasis after the first dose in all controlled trials was low (0.04%).

There were no changes in the heart rate of patients treated with telmisartan in controlled trials.

There are no trials of telmisartan demonstrating reductions in cardiovascular risk in patients with hypertension, but at least one pharmacologically similar drug has demonstrated such benefits.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **Telmisartan Tablets, USP are supplied as below:**

80 mg, white to off-white uncoated oval shaped tablets, biconvex with beveled edges, plain on one side and debossed "038" on other side.

NDC 63187-951-30 bottle of 30 units

NDC 63187-951-60 bottle of 60 units

NDC 63187-951-90 bottle of 90 units

#### *Storage*

**Store at 20° to 25°C (68° to 77°F)** [see USP Controlled Room Temperature]. see USP Controlled Room Temperature]. Tablets should not be removed from blisters until immediately before administration.

Dispense in a tightly closed container.

## **17 PATIENT COUNSELING INFORMATION**

*See FDA-approved Patient Labeling.*

### **17.1 Pregnancy**

Female patients of childbearing age should be told about the consequences of exposure to telmisartan during pregnancy. Discuss treatment options with women planning to become pregnant. Patients should be asked to report pregnancies to their physicians as soon as possible [*see Warnings and Precautions (5.1)*].

Manufactured By:

#### **Inventia Healthcare Private Limited**

Plot No.F1 & F-1/1, Additional Ambernath M.I.D.C.,

Ambernath (East)-421506,

Dist. Thane, Maharashtra, India

Distributed by:

Ascend Laboratories, LLC

Parsippany, NJ 07054

Revised: March 2017

Relabeled by:

Proficient Rx LP  
Thousand Oaks, CA 91320  
ATE0215L/02

**Patient Information**  
**Telmisartan Tablets USP**  
**(TEL mi SAR tan)**

**Read this Patient Information before you start taking telmisartan tablets USP and each time you get a refill. There may be new information. This information does not take the place of talking to your doctor about your medical condition or your treatment.**

**What is the most important information I should know about telmisartan tablets USP?**

Telmisartan tablets USP can cause harm or death to an unborn baby. Talk to your doctor about other ways to lower your blood pressure if you plan to become pregnant. If you get pregnant while taking telmisartan, tell your doctor right away.

**What is telmisartan tablets USP?**

Telmisartan tablets USP is a prescription medicine used:

- to treat high blood pressure (hypertension)

It is not known if telmisartan tablets USP is safe and effective in children.

**Who should not take telmisartan tablets USP?**

You should not take telmisartan tablets USP if you are allergic (hypersensitive) to the active ingredient (telmisartan) or any of the other ingredients listed at the end of this leaflet.

For patients with diabetes, if you are taking telmisartan you should not take aliskiren.

**What should I tell my doctor before taking telmisartan tablets USP?**

Before you take telmisartan tablets USP, tell your doctor if you:

- have liver problems
- have kidney problems
- have heart problems
- have any other medical conditions
- are pregnant or are planning to become pregnant. See **“What is the most important information I should know about telmisartan tablets USP?”**
- are breast-feeding or plan to breast-feed. It is not known if telmisartan tablets USP passes into your breast milk. You and our doctor should decide if you will take telmisartan tablets USP or breast-feed. You should not do both. Talk with your doctor about the best way to feed your baby if you take telmisartan tablets USP.

**Tell your doctor about all the medicines you take, including**

prescription and non-prescription medicines, vitamins, and herbal supplements.

For patients with diabetes, if you are taking telmisartan you should not take aliskiren.

Telmisartan tablets USP may affect the way other medicines work, and other medicines may affect how telmisartan works. Especially tell your doctor if you take:

- aliskiren
- digoxin (Lanoxin<sup>®</sup>, Lanoxicaps<sup>®</sup>)

- lithium (Eskalith<sup>®</sup>, Lithobid<sup>®</sup>)
- medicines used to treat pain and arthritis, called non-steroidal anti-inflammatory drugs (NSAIDs), including COX-2 inhibitors
- ramipril (Altace<sup>®</sup>) or other medicines used to treat your high blood pressure or heart problem
- water pills (diuretic)

Know the medicines you take. Keep a list of them and show it to your doctor or pharmacist when you get a new medicine.

### **How should I take telmisartan tablets USP?**

- Take telmisartan tablets USP exactly as your doctor tells you to take it.
- Your doctor will tell you how much telmisartan tablets USP to take and when to take it.
- Do not change your dose unless your doctor tells you to.
- Take telmisartan tablets USP one time each day at the same time.
- Take telmisartan tablets USP with or without food.
- If you miss a dose, take it as soon as you remember. If it is close to your next dose, do not take the missed dose. Take the next dose at your regular time.
- If you take too much telmisartan tablets USP, call your doctor, or go to the nearest hospital emergency room right away.
- Read the **“How to Open the Blister”** at the end of this leaflet before you use telmisartan tablets USP. Talk with your doctor if you do not understand the instructions.

### **What are the possible side effects of telmisartan tablets USP?**

#### **Telmisartan tablets USP may cause serious side effects, including:**

- **Injury or death to your unborn baby.** See **“What is the most important information I should know about telmisartan tablets USP?”**
- **Low blood pressure (hypotension)** is most likely to happen if you also:
  - take water pills (diuretics)
  - are on a low-salt diet
  - get dialysis treatments
  - have heart problems
  - get sick with vomiting or diarrhea

#### **If you feel faint or dizzy, lie down and call your doctor right away.**

- Kidney problems, which may get worse if you already have kidney disease. You may have changes in your kidney test results, and you may need a lower dose of telmisartan tablets USP. Call your doctor if you get:
  - swelling in your feet, ankles, or hands
  - unexplained weight gain

#### **Call your doctor right away if you get any of the symptoms listed above.**

- High potassium in the blood (hyperkalemia). Your doctor may check your potassium levels as needed.

Rare, serious allergic reactions may happen. Tell your doctor right away if you get any of these symptoms:

- swelling of the face, tongue, throat
- difficulty breathing
- skin rash

**The most common side effects of telmisartan tablets USP include:**

- sinus pain and congestion (sinusitis)
- back pain
- diarrhea

These are not all the possible side effects with telmisartan tablets USP. Tell your doctor if you have any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store telmisartan tablets USP tablets?**

- Store telmisartan tablets USP between 59°F to 86°F (15°C to 30°C).
- Do not remove telmisartan tablets USP from blisters until right before you take them.

**Keep telmisartan tablets USP and all medicines out of the reach of children.**

**General information about telmisartan tablets USP**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use telmisartan tablets USP for a condition for which it was not prescribed. Do not give telmisartan tablets USP to other people, even if they have the same condition you have. It may harm them.

This Patient Information leaflet summarizes the most important information about telmisartan tablets USP. If you would like more information, talk with your doctor. You can ask your pharmacist or doctor for information about telmisartan tablets USP that is written for health professionals.

For more information, call Ascend Laboratories, LLC, Parsippany, NJ 07054 at 1-877-272-7901

**What are the ingredients in telmisartan tablets USP?**

**Active Ingredient:** telmisartan USP

**Inactive Ingredients:** magnesium stearate, mannitol, meglumine, povidone, sodium hydroxide and sodium steryl fumarate.

**What is High Blood Pressure (Hypertension)?**


Blood pressure is the force in your blood vessels when your heart beats and when your heart rests. You have high blood pressure when the force is too much. Telmisartan tablets USP can help your blood vessels relax so your blood pressure is lower. Medicines that lower your blood pressure lower your chance of having a stroke or heart attack.

High blood pressure makes the heart work harder to pump blood throughout the body and causes damage to the blood vessels. If high blood pressure is not treated, it can lead to stroke, heart attack, heart failure, kidney failure, and vision problems.


**How to open the blister:**

**1. Tear (You may also use scissors to tear the blister apart)**





**2. Peel (Peel off the paper layer from the aluminum foil)**



Manufactured By:

**Inventia Healthcare Private Limited**

Plot No.F1 & F-1/1, Additional Ambernath M.I.D.C.,

Ambernath (East)-421506,

Dist. Thane, Maharashtra, India

Distributed by:

Ascend laboratories, LLC

Parsippany, NJ 07054

Revised: March 2017

Relabeled by:

Proficient Rx LP

Thousand Oaks, CA 91320


Trademarks are the property of their respective owners.


## PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

**Telmisartan tablets, USP 80 mg**

**30 Tablets**

**NDC : 63187-951-30**

Scan Here 

 NDC 63187-951-30 Relabeled By: Proficient Rx LP  
Thousand Oaks, CA 91320

**RX Only**

**Telmisartan 80mg**

**#30 Tablets**

Each tablet contains: 80 mg of telmisartan USP.

*White to off-white uncoated oval shaped tablets, biconvex with beveled edges, plain on one side and debossed "038" on other side.*


Product ID: RT095130

Mfr. By: Inventia Healthcare Limited, Additional Ambemath M.I.D.C., Ambemath (East) - 421506, INDIA.  
Store at 20°-25°C (68°-77°F) Keep medication out of the reach of children

Telmisartan 80mg #30 Tablets Lot # 00000 NDC 63187-951-30 SN# MASTER Exp. 00/00/00

Telmisartan 80mg #30 Tablets Lot # 00000 NDC 63187-951-30 SN# MASTER Exp. 00/00/00

Telmisartan 80mg #30 Tablets Lot # 00000 NDC 63187-951-30 SN# MASTER Exp. 00/00/00

 GTIN: 00363187951306  
SN# MASTER  
Exp. 00/00/00  
Lot # 00000

## TELMISARTAN

telmisartan tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63187-951(NDC:67877-484)
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TELMISARTAN (UNII: U5S YW473RQ) (TELMISARTAN - UNII:U5S YW473RQ)	TELMISARTAN	80 mg

### Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
MEGLUMINE (UNII: 6HG8UB2MUY)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	

## Product Characteristics

<b>Color</b>	WHITE (white or off-white)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	038
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63187-951-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2018	
2	NDC:63187-951-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2018	
3	NDC:63187-951-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2018	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA205150	06/12/2016	

**Labeler** - Proficient Rx LP (079196022)

## Establishment

Name	Address	ID/FEI	Business Operations
Proficient Rx LP		079196022	REPACK(63187-951) , RELABEL(63187-951)

Revised: 6/2020

Proficient Rx LP