

**EXTRA PAIN RELIEF- acetaminophen, aspirin and caffeine tablet, film coated
TIME CAP LABS**

Acetaminophen 250 mg, Aspirin 250 mg (NSAID), and Caffeine 65 mg Tablets

DRUG FACTS

Active ingredient (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:

headache

a cold

arthritis

muscular aches

toothache

premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause

severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout, or arthritis
- any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - o feel faint
 - o vomit blood
 - o have bloody or black stools
 - o have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or

complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

DIRECTIONS

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

OTHER INFORMATION

Other information

- store at 20°- 25°C (68°- 77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients carnauba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

***Questions or comments?* Call 1-877-290-4008**

NDC 49483-370-31




Compare to the active ingredients in Excedrin® Extra Strength†

Extra Strength **PAIN RELIEVER** Acetaminophen 250 mg, Aspirin 250 mg (NSAID), and Caffeine 65 mg

Pain Reliever/Pain Reliever Aid

300 CAPLETS** (**Capsule-Shaped Tablets)

Actual Size



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active Ingredients (in each caplet)

- Acetaminophen 250 mg... Pain reliever
Aspirin 250 mg (NSAID)*... Pain reliever
Caffeine 65 mg... Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses

- Temporarily relieves minor aches and pains due to: headache, a cold, arthritis, muscular aches, toothache, premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product...

Allergy alert: Acetaminophen may cause severe skin reactions...

Allergy alert: Aspirin may cause a severe allergic reaction...

Liver warning: This product contains acetaminophen...

Stomach bleeding warning: This product contains an NSAID...

PREPARE HERE

LOT:

EXP.:



3 4 9 4 8 3 1 3 7 0 3 1 7

Varnish Omit Area

Drug Facts (continued under label)

This product is not manufactured or distributed by GSK Consumer Healthcare, owner of the registered trademark Excedrin® Extra Strength.

Manufactured by: TimelyCaplets, Inc. 7 Micbad Avenue Farmingdale, NY 11735 370R 0324

Compare to the active ingredients in Excedrin® Extra Strength†

LOT:

EXP.:

PREPARE HERE

Varnish Omit Area

LOT:



NDC 49483-370-20

Compare to the active ingredients in Excedrin® Extra Strength†

Extra Strength **PAIN RELIEVER**
Acetaminophen 250 mg,
Aspirin 250 mg (NSAID),
and Caffeine 65 mg

Pain Reliever / Fever Reducer

200 CAPLETS**
(**Capsule-Shaped Tablets)

Actual Size



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active Ingredients (in each caplet)

Acetaminophen 250 mg.....Pain reliever
Aspirin 250 mg (NSAID).....Pain reliever
Caffeine 65 mg.....Pain reliever and
*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains due to:
■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache
■ premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

■ more than 8 caplets in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may

cause severe stomach bleeding. The chance is higher if you

■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use
■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if
■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are
■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away; ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

Other information
■ store at 20° - 25°C (68° - 77°); excursions permitted between 15° - 30°C (59° - 86°F) ■ use by expiration date on package

Inactive ingredients: camellia wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, polydioxane, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-877-290-4008

Adhesive Area

Drug Facts (continued)

cause severe stomach bleeding. The chance is higher if you

■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use
■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if
■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are
■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away; ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

Other information
■ store at 20° - 25°C (68° - 77°); excursions permitted between 15° - 30°C (59° - 86°F) ■ use by expiration date on package

Inactive ingredients: camellia wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, polydioxane, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-877-290-4008



Varnish Omit Area

Lot No.:

Exp Date:



Drug Facts (continued under label)

*This product is not manufactured or distributed by Healeon group of companies, owner of the registered trademark Excedrin® Extra Strength.

Manufactured by:
Time-Can Labs, Inc.
7, Michael Avenue Farmingdale,
NY 11735

370R 0823

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Exp. Date:
Lot No.:
3 49483 37042 3

Drug Facts

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Active ingredient (in each caplet) **Purpose**
Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID)* Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - cold
 - arthritis
 - muscular aches
 - toothache
 - premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: hives; facial swelling; shock; asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening; blisters; rash. If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: more than 8 caplets in 24 hours, which is the maximum daily amount; with other drugs containing acetaminophen; 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. For more than 10 days for pain unless directed by a doctor; for more than 3 days for fever unless directed by a doctor.

Ask a doctor before use if you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis or kidney disease; you are taking a diuretic.

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis; taking any other drug or are under a doctor's care for any serious condition

Drug Facts (continued)

Stop using and ask a doctor if symptoms do not improve; new symptoms occur; pain or fever persists or gets worse; you experience any of the following signs of stomach bleeding: feel faint; have bloody or black stools; vomit blood; have stomach pain that does not get better; ringing in the ears or loss of hearing occurs, consult a doctor before taking any more of this product.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions do not use more than directed; drink a full glass of water with each dose; adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours; children under 12 years: ask a doctor

Other information store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F); use by expiration date on package

Inactive ingredients carnauba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose,

NDC 49483-370-42

Compare to the active ingredients in Excedrin® Extra Strength



Extra Strength

PAIN RELIEVER
Acetaminophen 250 mg, Aspirin 250 mg (NSAID)
and Caffeine 65 mg

Pain Reliever / Pain Reliever Aid

24 CAPLETS
(Caplet-Shaped Tablets)

Actual Size

E3144R

Drug Facts (continued)
mineral oil, polydimethylsiloxane, polydioxane, sodium stearoyl phosphate, stearic acid, titanium dioxide

Questions or comments? call 1-877-298-0008

This product is not manufactured or distributed by GlaxoSmithKline plc, owner of the registered trademark Excedrin® Extra Strength. © 3708-1220 Distributed by: Time-Cap, Ltd., Inc., 7 Michael Avenue, Farmingdale, NY 11735



NDC 49483-370-42

Extra Strength PAIN RELIEVER

Acetaminophen 250 mg,
Aspirin 250 mg (NSAID),
and Caffeine 65 mg

Pain Reliever /
Pain Reliever Aid

24 CAPLETS**
(**Capsule-Shaped Tablet)

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Uses ■ temporarily relieves minor aches and pains due to: headache, a cold, arthritis, muscular aches, toothache ■ premenstrual and menstrual cramps

Warnings See carton for complete warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription (nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed ■ drink a full glass of water every 6 hours, not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

Other information ■ TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package **Questions or comments?** call 1-877-290-4008

Manufactured by: Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale, NY 11735
Lot No.:

Exp. Date:

370R 1120

Exp Date: _____
Lot No.: _____
49483 37001 3
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each caplet)	Purpose
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
■ headache ■ a cold ■ arthritis ■ muscular aches
■ toothache ■ premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 8 caplets in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop using and ask a doctor if

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood ■ have stomach pain that does not get better

Drug Facts (continued)

- ringing in the ears or loss of hearing occurs, consult a doctor before taking any more of this product.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours, not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

caruba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, pregelatinized starch, povidone, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? call 1-877-290-4008

*This product is not manufactured or distributed by GlaxoSmithKline plc, owner of the registered trademark Excedrin® Extra Strength, 370R 1020
Distributed by: Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale, NY 11735

NDC 49483-370-01
Compare to the active ingredients in Excedrin® Extra Strength!



Extra Strength
PAIN RELIEVER
Acetaminophen 250 mg, Aspirin 250 mg (NSAID),
and Caffeine 65 mg
Pain Reliever / Pain Reliever Aid
100 CAPLETS**
(Capsule-Shaped Tablet)

Actual Size

E6011AR5

Extra Strength
PAIN RELIEVER
Acetaminophen 250 mg,
Aspirin 250 mg (NSAID),
and Caffeine 65 mg
Pain Reliever / Pain Reliever Aid
100 CAPLETS**
(Capsule-Shaped Tablet)

NDC 49483-370-01



**Extra Strength
PAIN RELIEVER**
Acetaminophen 250 mg,
Aspirin 250 mg (NSAID),
and Caffeine 65 mg

Pain Reliever / Pain Reliever Aid

100 CAPLETS**
(*Capsule-Shaped Tablet)

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Uses ■ temporarily relieves minor aches and pains due to: headache, a cold, arthritis, muscular aches, toothache ■ premenstrual and menstrual cramps

Warnings See carton for complete warnings

Revers syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 8 capsules in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 2 caplets every 6 hours, not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

Other information

■ **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

■ use by expiration date on package

Questions or comments? call 1-877-390-4008

Manufactured by: Time-Cap Labs, Inc.
7 Michael Avenue
Farmingdale, NY 11735

370R 1020

Lot No.:

Exp. Date:

EXTRA PAIN RELIEF

acetaminophen, aspirin and caffeine tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49483-370
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg	
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg	

Inactive Ingredients		
Ingredient Name		Strength
STEARIC ACID (UNII: 4ELV7Z65AP)		
CARNAUBA WAX (UNII: R12CBM0EIZ)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
HYPROMELLOSES (UNII: 3NXW29V3WO)		
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
MINERAL OIL (UNII: T5L8T28FGP)		
POVIDONE (UNII: FZ989GH94E)		
STARCH, CORN (UNII: O8232NY3SJ)		
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE ((CAPSULE SHAPED TABLET))	Size	18mm
Flavor		Imprint Code	TCL370
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49483-370-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/16/2021	
2	NDC:49483-370-20	200 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2013	
3	NDC:49483-370-00	20000 in 1 CARTON; Type 0: Not a Combination Product	02/22/2016	
4	NDC:49483-370-42	1 in 1 CARTON	07/16/2021	
4		24 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:49483-370-31	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/01/2013	

Labeler - TIME CAP LABS (037052099)

Registrant - TIME CAP LABS (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABS		037052099	manufacture(49483-370)

Revised: 3/2024

TIME CAP LABS