BISMUTH SUBSALICYLATE- bismuth subsalicylate capsule, liquid filled Meijer, Inc.

Stomach Relief

Drug Facts

Active ingredient(in each softgel)
Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and anti-diarrheal

Uses

relieves

- travelers' diarrhea diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn indigestion nausea gas belching fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not Use if you have

- an ulcer
- a bleeding problem
- bloody or black stoo

Drug Facts

Ask a doctor before use if you have

fever • mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for • anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- swallow with water, do not chew
- adults and children 12 years and over:
- 2 softgels every $\frac{1}{2}$ hour or 4 softgels every hour as needed for diarrhea
- 2 softgels every $\frac{1}{2}$ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 softgels) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other Information

each softgel contains:salicylate 102 mg

- very low sodium
- do not exceed 25°C

Inactive ingredients

aerosil, beeswax, D&C Red#33, FD&C Yellow-6, gelatin, glycerin, light liquid paraffin,

povidone, purified water, sodium carboxy methyl cellulose, sorbitol sorbitan solution, soya lecithin, titanium dioxide

Ouestions or comments? 1-888-333-9792

Principle Display Panel

















"This product is not marufactured or distributed by Proctor & Gembé, owner of the tradements Pepto-Banol" and Liquicapa".

Questions or comments? + 88 83 9 92

ebit domunistit, nint sels yes, mo tube snistit res, toldine s liquit per stillin, po Michine, pur fledwater, s columics rhony m ethylos filos e, as rost, beas wax, D. &C. Reder 33, FD&C Yelowe & gelothn,gly cell n,lglit inactive ingredients

> 1. C. (66000 101 0) m n SOF elikyllez anisiere elestrese en everylow sodim everylower noitemnoini redtO

CHILES

 of nok pleny of clear fluids to halp proved dehydralb no aus ed by n children under 12 years a sk a do do r ■ use utilidarmee stops but not more than? do ys a do not esce de sagin es di Sa digeis) n 24 hou s fresneu unasafipu trinque ar to auros ■ 2s offgelse very 14 frounds 1 frounds needed for overlind utgence (upper in Zsofigelse very % four of 4 sofigels every hourses need bit der mee

■ sonce and children 12 years and over Directions - swallow withward, donotones

contact a Policon Control Certer rights way. a distribiti sissi u die dan Sosk a too opin set to sol to as sentinign bin in

Drug Facts (confined)

■ symptom sgat worse or be tmone the n≥d ays linetad eske knassugelö

apor su qo c qui dina u si cc ca. ent b grins Ausb, assim ei but "Ne rogm et et au bo q sidt g nieu es dW

e antico aguation (friming the dood) e date de god e a ritrius Naka doctor or phasmacint before uself you are lakeliga ny drug Tor IOOS BUILDI SECTIUI ■ JB ABL ■

we due to belone use if you have

m su npst m s pesquib brogeu m picoqù a pisx s poi eval ucy tesub ne d

■ pay yiết quete siph ja pa tron nga early sign of 18.94 5 syntrome, a sa ea buts efrous ily on ene Allengy a leaf. Combains selleybate. Do neafable ily on ene a allengic fiz sicy bias finculating septin) м шридосот с со изпра досрогрества дляя вущо ше сопр рачи When us ingthis product, to fanges in behavior with nausea and Rejer's syndion o e Childre nand beenages who have orase noovering from chicken poxorfluilke symptoms should nd use this product. Marnings

профессования профессования

38 AB IB J sasn

Leath dibitne bine gm 595 ediskydlasdua Upost stomach is leve r Active ingredient (in each softgel) Purpose

Drug Facts

meijer **Stomach Relief**

BISMUTH SUBSALICYLATE 262 mg / UPSET STOMACH RELIEVER • ANTIDIARRHEAL

meijer

Stomach Relief

5 SYMPTOM DIGESTIVE RELIEF

- HEARTBURN INDIGESTION UPSET STOMACH
- NAUSEA DIARRHEA

BISMUTH SUBSALICYLATE 262 mg UPSET STOMACH RELIEVER • ANTIDIARRHEAL NDC 79481-0058-2

COMPARE TO PEPTO-BISMOL® LIQUICAPSTM **ACTIVE INGREDIENT***

> 24 SOFTGELS



DATE ä 5

BVIDENT: DO NOT USE IF PACKAGE IS TORN, BROKEN OR SHOWS ANY

BISMUTH SUBSALICYLATE 262 mg / UPSETSTOMACH RELIEVER • ANTIDIARRHEA Relief

BISMUTH SUBSALICYLATE

bismuth subsalicylate capsule, liquid filled

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0058
--------------	----------------	--------------------	----------------

Route of Administration ORAL

Active Ingredient/Active Moiety Ingredient Name BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII: 0414PZ4LPZ) BISMUTH SUBSALICYLATE BISMUTH SUBSALICYLATE BISMUTH SUBSALICYLATE

Inactive Ingredients			
Ingredient Name	Strength		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
SYNTHETIC BEESWAX (UNII: 08MNR5YE2R)			
D&C RED NO. 33 (UNII: 9DBA0SBB0L)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
GELATIN (UNII: 2G86QN327L)			
GLYCERIN (UNII: PDC6A3C0OX)			
PARAFFIN (UNII: 1900E3H2ZE)			
POVIDONE (UNII: FZ989GH94E)			
WATER (UNII: 059QF0KO0R)			
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)			
SORBITOL SOLUTION (UNII: 8KW3E207O2)			
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			

Product Characteristics				
Color	pink	Score	no score	
Shape	OVAL	Size	18mm	
Flavor		Imprint Code	854	
Contains				

P	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481- 0058-2	2 in 1 CARTON	02/01/2024	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	02/01/2024	

Labeler - Meijer, Inc. (006959555)

Revised: 4/2024 Meijer, Inc.