

COCCIAID- amprolium solution
Aurora Pharmaceutical, Inc.

CocciAid®

(amprolium)

9.6% Oral Solution

DIN 02521040

Net Contents:

3.8 L

FOR VETERINARY USE ONLY

Active Ingredient:

Each mL contains 96 mg of amprolium.

Benzoic acid 0.1% added as preservative.

INDICATIONS:

CHICKENS

As an aid in the treatment of caecal coccidiosis in growing chickens and laying birds.

CALVES

As an aid in the treatment of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves.

USE DIRECTIONS:

CHICKENS

As soon as caecal coccidiosis is diagnosed, give 0.024% amprolium in the drinking water for 5 to 7 days.

Continue treatment with 0.006% amprolium medicated water for an additional one to two weeks.

No other source of drinking water should be available to the birds during this time.

Use as the sole source of amprolium.

CALVES

0.012% amprolium in drinking water for 5 days. At the usual rate of water consumption, this will provide a daily intake of approximately 10 mg of amprolium per kg of body weight. Give as the sole source of water during the treatment period.

NOTE

When one or more calves show signs of coccidiosis, it is likely that the rest of the group have been exposed, and all calves in the group should be treated.

WARNING:

Treated calves must not be slaughtered for use in food for at least 7 days after the latest treatment with this drug.

No withdrawal period is required for meat or eggs when chickens are treated according to the label.

When handling the product, avoid oral exposure and direct contact with skin or eyes.

Keep out of reach of children.

CAUTIONS:

CHICKENS

If no improvement is noted within 3 days, have the diagnosis reconfirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

CALVES

For an appropriate diagnosis, a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

Do not use in calves intended for future breeding

MIXING DIRECTIONS:

To prepare 200 L of medicated water

DOSAGE MIXING DIRECTIONS

0.024% Add 500 mL of CocciAid[®] (amprolium) 9.6% Solution to about 25 L of water in a 200 L medication barrel. Stir, then add water to the 200 L mark. *Stir thoroughly.*

0.012% Follow same directions as above but use 250 mL of CocciAid[®] 9.6% Solution.

0.006% Follow same directions as above but use 125 mL of CocciAid[®] 9.6% Solution.

STORAGE:

Store at 15° to 25°C. Protect from freezing.

REORDER NO: 24003

Manufactured by:

Aurora Pharmaceutical, Inc.
Northfield, MN 55057

1-888-215-1256

Distributed by:

Partner Animal Health Inc.

Ilderton, Ontario NDM 2A0

IN 50-1628 09/2021

Container Label

PRINCIPAL DISPLAY PANEL - 3.8L bottle label



DIN 02521040

Net Contents
Contenu net
3.8 L

CocciAid®

Amprolium Oral Solution 9.6 % /
Solution orale d'amprolium à 9,6 %**FOR
VETERINARY
USE ONLY****USAGE
VÉTÉRINAIRE
SEULEMENT****Active ingredient:**Each mL contains 96 mg of amprolium.
Benzoic acid 0.1% added as preservative.**WARNINGS**Treated calves must not be slaughtered for use in
food for at least 7 days after the latest treatment
with this drug.No withdrawal period is required for meat or eggs
when chickens are treated according to the label.When handling the product, avoid oral exposure
and direct contact with skin or eyes.

Keep out of reach of children.

® Trademark AURORA PHARMACEUTICAL, INC.

® Marque déposée de AURORA
PHARMACEUTICAL, INC.**Ingrédient actif:**Chaque mL renferme 96 mg d'amprolium.
L'acide benzoïque 0,1% est ajouté comme agent
de conservation.**MISES EN GARDE**Les veaux traités ne doivent pas être abattus à des
fins alimentaires dans un délai d'au moins 7 jours
après la dernière administration de ce médicament.Un délai d'attente n'est pas requis pour la viande
ou les œufs quand les poulets sont traités
conformément aux directives de l'étiquette.Lors de la manipulation du produit, éviter
l'exposition orale et le contact direct
avec la peau ou les yeux.Garder hors de la portée
des enfants.**COCCIAID**

amprolium solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:51072-111
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMPROLIUM (UNII: 95CO6N199Q) (AMPROLIUM ION - UNII:H2T307KMZR)	AMPROLIUM	96 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51072-111-00	3785 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
export only		03/15/2023	

Labeler - Aurora Pharmaceutical, Inc. (832848639)

Establishment

Name	Address	ID/FEI	Business Operations
Aurora Pharmaceutical, Inc.		832848639	manufacture

Revised: 3/2023

Aurora Pharmaceutical, Inc.