

WOMENS GENTLE LAXATIVE- bisacodyl tablet, delayed release
CHAIN DRUG MARKETING ASSOCIATION INC

Quality Choice 44-676A

Active ingredient (in each tablet)

Bisacodyl USP, 5 mg

Purpose

Stimulant laxative

Uses

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children under 6 years	ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- avoid excessive humidity
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

carmine, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal display panel

**QC®
Quality
Choice**

NDC 63868-573-30

Women’s Gentle Laxative

Bisacodyl USP, 5 mg - Stimulant Laxative

Gentle, Dependable
Constipation Relief

30 Comfort Coated Tablets

Actual
Size

50844 ORG011967601

**TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING**

Distributed by: C.D.M.A., Inc, ©
43157 W 9 Mile Rd
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362

**SATISFACTION
GUARANTEED
100%
QC**



NDC 63868-573-30

Women's Gentle Laxative

Bisacodyl USP, 5 mg - Stimulant Laxative

Gentle, Dependable Constipation Relief



30 Comfort Coated Tablets

Distributed by C.D.M.A., Inc.®
43157 W 9 Mile Rd
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362



Drug Facts (continued)
polydextrose, polyethylene glycol, stibac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, tracefin, triethyl citrate
Questions or comments? 1-800-426-9391

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844 ORG011967601

No Print / No Varnish Lot no. & Exp. date

Drug Facts
KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION
Active ingredient (in each tablet) Bisacodyl USP, 5 mg.....Stimulant laxative
Purpose ■ for relief of occasional constipation (irregularity) ■ this product generally produces bowel movement in 6 to 12 hours
Uses
Drug Facts (continued)

B-0220-676A-01-R
ORG011967601

Drug Facts (continued)
Warnings
Do not use if you cannot swallow without chewing.
Ask a doctor before use if you have
■ a sudden change in bowel habits that lasts more than 2 weeks
■ stomach pain, nausea or vomiting
When using this product
■ do not chew or crush tablet(s)
■ do not use within 1 hour after taking an antacid or milk
■ you may have stomach discomfort, faintness and cramps
Stop use and ask a doctor if
■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
■ you need to use a laxative for more than 1 week
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Drug Facts (continued)
Directions
■ take with a glass of water
adults and children 12 years and over take 1 to 3 tablets in a single daily dose
children 6 to under 12 years take 1 tablet in a single daily dose
children under 6 years ask a doctor
Other information
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
■ avoid excessive humidity
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number
Inactive ingredients carmine, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose.

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-573
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	B
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-573-30	2 in 1 CARTON	02/17/2020	
1		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	02/17/2020	

Labeler - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(63868-573)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63868-573) , pack(63868-573)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63868-573)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(63868-573)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(63868-573)

Revised: 1/2024

CHAIN DRUG MARKETING ASSOCIATION INC