# **TANDEM PLUS NPE FREE- iodine, dodecylbenzenesulfonic acid liquid IBA**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

#### **Tandem Plus NPE Free**

#### **ACTIVE INGREDIENTS**

Linear Dodecyl Benzene Sulfonic Acid (DDBSA) 1.90%	ó
Iodine 0.25%	
(Provides a full 0.25% Available Iodine as used)	

## **INERT INGREDIENTS:**

Glycerine (as a humectant and skin	conditioner) 10.00%
Other inerts:	87.85%
(Includes Iodine Stabilizers, Bufferin	<i>,</i>

This product contains highly effective levels of free iodine for fast kill of mastitis causing organisms.

## RECOMMENDED PROCEDURES FOR TEAT SANITATION

## A. Pre-Milking (Pre-dipping)

- 1. Remove visible solids from the teats.
- 2. Observe foremilk by stripping two or three streams of milk into a strip cup
- 3. Dip, foam or spray the cow's teats full length with **TANDEM + Teat Dip**
- 4. Wait approximately 30 seconds
- 5. Remove all teat dip thoroughly dryin the teats, teat orifices and udder (if foaming or spraying) with a clean, single service paper towel(s) to avoid contamination of milk.
- 6. Attach milking unit.

## **B. POST-MILKING (POST-DIPPING)**

Immediately after milking dip or spray all teats full length with **TANDEM + Teat Dip**. Ensure good coverage of all sides of the teats and teat orifices.

## **GENERAL RECOMMENDATIONS**

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.

- 2. When solution in the teat dip cup becomes dirty, discard remaining contents and refill cup with fresh teat dip.
- 3. Do not use this product for cleaning or sanitizing equipment
- 4. Wash Teat Dip cups after each milking.

## **WARNING**

DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.
PROTECT THIS PRODUCT FROM FREEZING.

NOT FOR HUMAN USE.

CAUTION: KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

## **FIRST AID PROCEDURES**

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR READY-TO-USE PRODUCTS. WASH ANY SENSITIVE AREAS. INGESTION:DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES CALL 1-800-424-9300

## **Tanden Plus NPE Free**



## NPE FREE SANITIZING TEAT DIP

A Highly Effective and Advanced Formula Contains **TWO** Germicides and 10% Glycerin

## ACTIVE INGREDIENTS: Linear Dodecyl Benzene Sulfonic Acid (DDBSA).... (Provides A Full .25% Available lodine As Used) INERT INGREDIENTS: Glycerin (As A Humectant And Skin Conditioner)... Other Inerts

This product contains highly effective levels of free iodine for fast kill of mastitis causing organisms.

#### **GENERAL RECOMMENDATIONS**

- Use teat dip applicators made of plastic or other non-corrosive materials.
   Do not use cups made of aluminum.
   When solution in the teat dip cup becomes dirty, discard remaining contents and refill cup with fresh teat dip.
   Do not use this product 'or cleaning or sanitizing equipment.
   Wash teat dip cups after each milking.

AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

## FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID ITSSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES CALL 1-800-424-9300



**PATENT PENDING** 

## RECOMMENDED PROCEDURES FOR TEAT SANITATION

#### A. PRE-MILKING (PRE-DIPPING)

- 1. Remove visible soils from the teat
- 2. Observe foremilk by stripping two or three streams of milk into a strip cup.
- 3. Dip, foam or spray the cow's teats full length with TANDEM +
- 4. Wait approximately 30 seconds.
- Remove all teat dip by thoroughly drying the teats, teat orifices and udder (if foaming or spraying) with a clean, single service paper towel(s) to avoid contamination of milk..
- 6. Attach milking unit.

#### B. POST-MILKING (POST-DIPPING)

Immediately after milking, dip or spray all teats full length with TANDEM + Teat Dip. Ensure good coverage of all sides of the teats and teat orifices.

#### WARNING

DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

PROTECT THIS PRODUCT FROM FREEZING

## A PART OF THE COMPLETE IBA SANITATION PROGRAM

**EXP. DATE:** 

**NET CONTENTS: 275 GALLONS** 

**IBA STOCK # 260175** 

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

## **TANDEM PLUS NPE FREE**

iodine, dodecylbenzenesulfonic acid liquid

<b>Product Type</b>	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-1076
---------------------	-----------------	--------------------	----------------

Route of Administration TOPICAL

Active Ingredient/Active Moiety				
Ingredient Name	<b>Basis of Strength</b>	Strength		
DODECYLBENZENESULFONIC ACID (UNII: 60NSK897G9) (DODECYLBENZENESULFONIC ACID - UNII:60NSK897G9)	DODECYLBENZ ENES ULFONIC ACID	1.9 kg in 100 kg		
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	0.25 kg in 100 kg		

Packaging				
#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
1	NDC:29983-1076-9	1072.50 kg in 1 CONTAINER		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		08/17/2020	

## **Labeler -** IBA (019494160)

## **Registrant - IBA** (019494160)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
Morgan Gallacher Inc DBA Custom Chemical Formulators Inc		028311595	manufacture, api manufacture, pack

Revised: 12/2023 IBA