


TRIAMCINOLONE ACETONIDE - triamcinolone acetonide cream
Cosette Pharmaceuticals, Inc.

Triamcinolone Acetonide Cream USP

FOR EXTERNAL USE ONLY
NOT FOR OPHTHALMIC USE
Rx Only

DESCRIPTION

Triamcinolone Acetonide Cream USP contains triamcinolone acetonide [Pregna-1,4-diene-3,20-dione, 9-fluoro-11,21-dihydroxy-16,17-[(1-methylethylidene)bis- (oxy)]-, (11 β ,16 α)-], with the empirical formula C₂₄H₃₁FO₆ and molecular weight 434.50. CAS 76-25-5.



Triamcinolone Acetonide Cream USP 0.025% contains: 0.25 mg of triamcinolone acetonide USP per gram in a water washable cream base consisting of mineral oil (and) lanolin alcohol, isopropyl palmitate NF, propylene glycol stearate, propylene glycol USP, cetyl alcohol NF, sorbitan monostearate NF, polysorbate 60 NF, sorbic acid NF, polyoxyl (40) stearate NF, propylparaben NF, methylparaben NF and purified water USP.

Triamcinolone Acetonide Cream USP 0.1% contains: 1 mg of triamcinolone acetonide USP per gram in a water washable cream base consisting of mineral oil (and) lanolin alcohol, isopropyl palmitate NF, propylene glycol stearate, propylene glycol USP, cetyl alcohol NF, sorbitan monostearate NF, polysorbate 60 NF, sorbic acid NF, polyoxyl (40) stearate NF, propylparaben NF, methylparaben NF and purified water USP.

CLINICAL PHARMACOLOGY

Topical corticosteroids share anti-inflammatory, anti-pruritic and vasoconstrictive actions. The mechanism of anti-inflammatory activity of the topical corticosteroids is unclear. Various laboratory methods, including vasoconstrictor assays, are used to compare and predict potencies and/or clinical efficacies of the topical corticosteroids. There is some evidence to suggest that a recognizable correlation exists between vasoconstrictor potency and therapeutic efficacy in man.

Pharmacokinetics

The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle, the integrity of the epidermal barrier, and the use of occlusive dressings. Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin increase percutaneous absorption. Occlusive dressings substantially increase the percutaneous absorption of topical corticosteroids. Thus, occlusive dressings may be a valuable therapeutic adjunct for treatment of resistant dermatoses (See **DOSAGE AND ADMINISTRATION**). Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to systemically administered corticosteroids. Corticosteroids are bound to plasma proteins in varying degrees. Corticosteroids are metabolized primarily in the liver and are then excreted by the kidneys. Some of the topical corticosteroids and their metabolites are also excreted into the bile.

INDICATIONS AND USAGE

Topical corticosteroids are indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses.

CONTRAINDICATIONS

Topical corticosteroids are contraindicated in those patients with a history of hypersensitivity to any of the components of the preparation.

PRECAUTIONS

General

Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia, and glucosuria in some patients. Conditions which augment systemic absorption include the application of the more potent steroids, use over large surface areas, prolonged use, and the addition of occlusive dressings. Therefore, patients receiving a large dose of a potent topical steroid applied to a large surface area or under an occlusive dressing should be evaluated periodically for evidence of HPA axis suppression by using the urinary free cortisol and ACTH stimulation tests. If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent steroid. Recovery of HPA axis function is generally prompt and complete upon discontinuation of the drug.

Infrequently, signs and symptoms of steroid withdrawal may occur, requiring supplemental systemic corticosteroids. Children may absorb proportionally larger amounts of topical corticosteroids and thus be more susceptible to systemic toxicity (See **PRECAUTIONS-Pediatric Use**). If irritation develops, topical corticosteroids should be discontinued and appropriate therapy instituted. In the presence of dermatological infections, the use of an appropriate anti-fungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, the corticosteroid should be discontinued until the infection has been adequately controlled.

Information for the Patient

Patients using topical corticosteroids should receive the following information and instructions.

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. Patients should be advised not to use this medication for any disorder other than for which it was prescribed.
3. The treated skin area should not be bandaged or otherwise covered or wrapped as to be occlusive unless directed by the physician.
4. Patients should report any signs of local adverse reactions especially under occlusive dressing.

5. Parents of pediatric patients should be advised not to use tight fitting diapers or plastic pants on a child being treated in the diaper area, as these garments may constitute occlusive dressings.

Laboratory Tests

The following tests may be helpful in evaluating the HPA axis suppression: Urinary free cortisol test; ACTH stimulation test.

Carcinogenesis, Mutagenesis, and Impairment of Fertility

Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of topical corticosteroids. Studies to determine mutagenicity with prednisolone and hydrocortisone have revealed negative results.

Pregnancy: Teratogenic Effects -

Pregnancy Category C

Corticosteroids are generally teratogenic in laboratory animals when administered systemically at relatively low dosage levels. The more potent corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. There are no adequate and well-controlled studies in pregnant women on teratogenic effects from topically applied corticosteroids. Therefore, topical corticosteroids should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Drugs of this class should not be used extensively on pregnant patients, in large amounts, or for prolonged periods of time.

Nursing Mothers

It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids are secreted into breast milk in quantities not likely to have a deleterious effect on the infant. Nevertheless, caution should be exercised when topical corticosteroids are administered to a nursing woman.

Pediatric Use

Pediatric patients may demonstrate greater susceptibility to topical corticosteroid-induced HPA axis suppression and Cushing's syndrome than mature patients because of a larger skin surface area to body weight ratio. Hypothalamic-pituitary-adrenal (HPA) axis suppression, Cushing's syndrome, and intracranial hypertension have been reported in children receiving topical corticosteroids. Manifestations of adrenal suppression in children include linear growth retardation, delayed weight gain, low plasma cortisol levels, and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema. Administration of topical corticosteroids to children should be limited to the least amount compatible with an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of children.

ADVERSE REACTIONS

The following local adverse reactions are reported infrequently with topical corticosteroids, but may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae and miliaria.

To report SUSPECTED ADVERSE REACTIONS, contact Cosette Pharmaceuticals, Inc. at 1-800-922-1038 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects (See **PRECAUTIONS**).

DOSAGE AND ADMINISTRATION

Apply Triamcinolone Acetonide Cream USP 0.025% to the affected area two to four times daily. Rub in gently.

Apply the 0.1% Triamcinolone Acetonide Cream, as appropriate, to the affected area two to three times daily. Rub in gently.

Occlusive dressings may be used for the management of psoriasis or recalcitrant conditions. If an infection develops, the use of occlusive dressings should be discontinued and appropriate antimicrobial therapy instituted.

HOW SUPPLIED

Triamcinolone Acetonide Cream USP 0.025%
15 gram tubes NDC 0713-0226-15
80 gram tubes NDC 0713-0226-80

Triamcinolone Acetonide Cream USP 0.1%
15 gram tubes NDC 0713-0225-15
30 gram tubes NDC 0713-0225-31
80 gram tubes NDC 0713-0225-80

STORAGE

**Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].
Keep out of reach of children.
Avoid excessive heat. Protect from freezing.**

Manufactured by:

Cosette Pharmaceuticals, Inc.
111 Coolidge Street,
South Plainfield, NJ 07080

Iss. 07/2019
8-TRIACRCP1

PRINCIPAL DISPLAY PANEL

NDC 0713-0225-15

Triamcinolone Acetonide Cream, USP 0.1%

15 g

Rx only

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

Cosette Pharmaceuticals, Inc.





**Triamcinolone Acetonide
Cream, USP**

NDC 0713-0225-15

Rx only

0.1%

15 g

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

Usual Dose: Apply a small amount to the affected area two to three times daily.
For complete information read accompanying insert.

WARNING: Keep out of reach of children.

Each gram contains: 1 mg triamcinolone acetonide USP in a water washable cream base consisting of mineral oil (and) lanolin alcohol, isopropyl palmitate NF, propylene glycol stearate, propylene glycol USP, cetyl alcohol NF, sorbitan monostearate NF, polysorbate 60 NF, sorbic acid NF, polyoxyl (40) stearate NF, propylparaben NF, methylparaben NF and purified water USP.

Store at 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature].

Avoid excessive heat. Protect from freezing.

Manufactured by:

Cosette Pharmaceuticals, Inc.
111 Coolidge Street
South Plainfield, NJ 07080

Iss. 02/2020
11-22515CP2



N
3

0713-0225-15

1

NDC 0713-0225-31

Triamcinolone Acetonide Cream, USP 0.1%

30 g

Rx only

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

Cosette Pharmaceuticals, Inc.

Glue Flap



**Triamcinolone Acetonide
Cream, USP**

NDC 0713-0225-31

Rx only

0.1%

30 g

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.



Usual Dose: Apply a small amount to the affected area two to three times daily.
For complete information read accompanying insert.

WARNING: Keep out of reach of children.

Manufactured by:
Cosette Pharmaceuticals, Inc.
111 Coolidge Street
South Plainfield, NJ 07080 10-22531CP1 Iss. 06/2019

GTIN 00307130225311



N 3 0713-0225-31 1



**Triamcinolone Acetonide
Cream, USP**

NDC 0713-0225-31

Rx only

0.1%

30 g

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.



Each Gram Contains: 1 mg triamcinolone acetonide USP in a water washable cream base consisting of mineral oil (and) lanolin alcohol, isopropyl palmitate NF, propylene glycol stearate, propylene glycol USP, cetyl alcohol NF, sorbitan monostearate NF, polysorbate 60 NF, sorbic acid NF, polyoxyl (40) stearate NF, propylparaben NF, methylparaben NF and purified water USP.

**Store at 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature].
Avoid excessive heat. Protect from freezing.**



Bend Line

NDC 0713-0225-31

Triamcinolone Acetonide Cream, USP

Rx only

0.1%

30 g

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

Usual Dose: Apply a small amount to the affected area two to three times daily.
For complete information read accompanying insert.

WARNING: Keep out of reach of children.

Each gram contains: 1 mg triamcinolone acetonide USP in a water washable cream base consisting of mineral oil (and) lanolin alcohol, isopropyl palmitate NF, propylene glycol stearate, propylene glycol USP, cetyl alcohol NF, sorbitan monostearate NF, polysorbate 60 NF, sorbic acid NF, polyoxyl (40) stearate NF, propylparaben NF, methylparaben NF and purified water USP.

Store at 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature].

Avoid excessive heat. Protect from freezing.

Manufactured by:

Cosette Pharmaceuticals, Inc.

111 Coolidge Street

South Plainfield, NJ 07080

11-22531CP1

Iss. 06/2019



N
3

0713-0225-31

1

0713-0225-80


Triamcinolone Acetonide Cream, USP 0.1%

80 g

Rx only

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

Cosette Pharmaceuticals, Inc.



DC 0713-0226-15

Triamcinolone Acetonide Cream, USP 0.1%

15 g

Rx only

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

Cosette Pharmaceuticals, Inc.





**Triamcinolone Acetonide
Cream, USP**

NDC 0713-0226-15

Rx only

0.025%

15 g

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

Usual Dose: Apply a small amount to the affected area two to four times daily.
For complete information read accompanying insert.

WARNING: Keep out of reach of children.

Each gram contains: 0.25 mg triamcinolone acetonide USP in a water washable cream base consisting of mineral oil (and) lanolin alcohol, isopropyl palmitate NF, propylene glycol stearate, propylene glycol USP, cetyl alcohol NF, sorbitan monostearate NF, polysorbate 60 NF, sorbic acid NF, polyoxyl (40) stearate NF, propylparaben NF, methylparaben NF and purified water USP.

Store at 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature].

Avoid excessive heat. Protect from freezing.

Manufactured by:

Cosette Pharmaceuticals, Inc.
111 Coolidge Street
South Plainfield, NJ 07080

Rev. 04/2020
11-22615CP2



N
3

0713-0226-15

8

NDC 0713-0226-80

Triamcinolone Acetonide Cream, USP 0.1%

80 g

Rx only

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

Cosette Pharmaceuticals, Inc.

Glue Flap



**Triamcinolone Acetonide
Cream, USP**

NDC 0713-0226-80
Rx only

0.025%

80 g

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

Usual Dose: Apply a small amount to the affected area two to four times daily.
For complete information read accompanying insert.

WARNING: Keep out of reach of children.

Manufactured by:
Cosette Pharmaceuticals, Inc.
111 Coolidge Street
South Plainfield, NJ 07080 10-22680CP1 Iss. 06/2019

GTIN 00307130226806



N 3 0713-0226-80 6



**Triamcinolone Acetonide
Cream, USP**

NDC 0713-0226-80
Rx only

0.025%

80 g

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

Each Gram Contains: 0.25 mg triamcinolone acetonide USP in a water washable cream base consisting of mineral oil (and) lanolin alcohol, isopropyl palmitate NF, propylene glycol stearate, propylene glycol USP, cetyl alcohol NF, sorbitan monostearate NF, polysorbate 60 NF, sorbic acid NF, polyoxyl (40) stearate NF, propylparaben NF, methylparaben NF and purified water USP.

**Store at 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature].
Avoid excessive heat. Protect from freezing.**

Bend Line



**Triamcinolone Acetonide
Cream, USP**

NDC 0713-0226-80

Rx only

0.025%

80 g

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

Usual Dose: Apply a small amount to the affected area two to four times daily.
For complete information read accompanying insert.

WARNING: Keep out of reach of children.

Each gram contains: 0.25 mg triamcinolone acetonide USP in a water washable cream base consisting of mineral oil (and) lanolin alcohol, isopropyl palmitate NF, propylene glycol stearate, propylene glycol USP, cetyl alcohol NF, sorbitan monostearate NF, polysorbate 60 NF, sorbic acid NF, polyoxyl (40) stearate NF, propylparaben NF, methylparaben NF and purified water USP.

Store at 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature].

Avoid excessive heat. Protect from freezing.

Manufactured by:

Cosette Pharmaceuticals, Inc.
111 Coolidge Street
South Plainfield, NJ 07080
11-22680CP1 Iss. 06/2019



N 3 0713-0226-80 6

TRIAMCINOLONE ACETONIDE

triamcinolone acetonide cream

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0713-0226
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRIAMCINOLONE ACETONIDE (UNII: F446C597KA) (TRIAMCINOLONE ACETONIDE - UNII:F446C597KA)	TRIAMCINOLONE ACETONIDE	0.25 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
MINERAL OIL (UNII: T5L8T28FGP)	
LANOLIN ALCOHOLS (UNII: 884C3FA9HE)	
ISOPROPYL PALMITATE (UNII: 8CRQ2TH63M)	
PROPYLENE GLYCOL STEARATE (UNII: MZM1I680W0)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
SORBITAN MONOSTEARATE (UNII: NVZ4I0H58X)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
SORBIC ACID (UNII: X045WJ989B)	
POLYOXYL 40 STEARATE (UNII: 13A4J4NH9I)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0713-0226-15	1 in 1 CARTON	04/14/2015	
1		15 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0713-0226-80	1 in 1 CARTON	04/14/2015	
2		80 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA089797	04/14/2015	

TRIAMCINOLONE ACETONIDE

triamcinolone acetonide cream

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0713-0225
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRIAMCINOLONE ACETONIDE (UNII: F446C597KA) (TRIAMCINOLONE ACETONIDE - UNII:F446C597KA)	TRIAMCINOLONE ACETONIDE	1 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
MINERAL OIL (UNII: T5L8T28FGP)	
LANOLIN ALCOHOLS (UNII: 884C3FA9HE)	
ISOPROPYL PALMITATE (UNII: 8CRQ2TH63M)	
PROPYLENE GLYCOL STEARATE (UNII: MZM1I680W0)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
SORBITAN MONOSTEARATE (UNII: NVZ4I0H58X)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
SORBIC ACID (UNII: X045WJ989B)	
POLYOXYL 40 STEARATE (UNII: 13A4J4NH9I)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
WATER (UNII: 059QF0K00R)	

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0713-0225-15	1 in 1 CARTON	04/14/2015	
1		15 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0713-0225-31	1 in 1 CARTON	04/14/2015	
2		30 g in 1 TUBE; Type 0: Not a Combination Product		
3	NDC:0713-0225-80	1 in 1 CARTON	04/14/2015	
3		80 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA089798	04/14/2015	

Labeler - Cosette Pharmaceuticals, Inc. (116918230)

Registrant - Cosette Pharmaceuticals, Inc. (116918230)

Establishment

Name	Address	ID/FEI	Business Operations
Cosette Pharmaceuticals, Inc.		116918230	analysis(0713-0225, 0713-0226) , label(0713-0225, 0713-0226) , manufacture(0713-0225, 0713-0226) , pack(0713-0225, 0713-0226)

Revised: 10/2020

Cosette Pharmaceuticals, Inc.