

**DESENFRIOL-D- chlorpheniramine maleate, phenylephrine hydrochloride,
acetaminophen tablet**
Salimex, S.A.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Desenfriol-D

Active ingredient (in each caplet)

CHLORPHENIRAMINE MALEATE 2.0 mg

PHENYLEPHRINE HYDROCHLORIDE 5.0 mg

ACETAMINOPHEN 500 mg

Purpose

Pain reliever

Antihistamine

Nasal decongestant

Uses

- temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
- headache
- sinus congestion and pressure
- nasal congestion
- runny nose and sneezing
- minor aches and pains
- temporarily relieves these additional symptoms of hay fever:
- itching of the nose or throat
- itchy, watery eyes
- helps clear nasal passages
- helps decongest sinus openings and passages

Warnings

- Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product
- Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
- blisters
- rash
- If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure

whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- to make a child sleepy

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.etc

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 caplets every 4-6 hours
- swallow whole – do not crush, chew or dissolve
- do not take more than 10 caplets in 24 hours

children under 12 years

- ask a doctor

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C red #33, FD&C blue #1, FD&C red 40, flavors, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

Desfriol-D

Tablets

DESENFRIOL D

CHLORPHENIRAMINE, PHENYLEPHRINE, ACETAMINOPHEN
2 mg / 5 mg / 500 mg

anti-flu

For the temporary relief of discomfort due to the common cold

- Headache
- Fever
- Nasal congestion
- Aching body
- Sneezing
- Runny nose
- Crying eyes
- Sore throat

~~24~~ **30** Tables



DESENFRIOL D

Active ingredients Purpose

MALEATE 2.0 mg.....	Antihistamine
PHENYLEPHRINE HYDROCHLORIDE 5.0 mg	Nasal decongestant
ACETAMINOPHEN 500 mg.....	Pain reliever and a fever reducer

Indications

Temporarily relieves runny nose and sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever. For the temporary relief of nasal congestion due to the common cold.

Warnings

- Do not give to children under 3 years of age or use for more than 10 days unless directed by a physician.
- May cause excitability especially in children.
- Do not take this product, unless directed by a doctor, if you have a breathing problem such as emphysema or chronic

bronchitis, or if you have glaucoma or difficulty in urination due to enlargement of the prostate gland.

May cause drowsiness; alcohol, sedatives, and tranquilizers may increase the drowsiness effect. Avoid alcoholic beverages while taking this product. Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor. Use caution when driving a motor vehicle or operating machinery. Sedatives and tranquilizers may increase the drowsiness effect. Do not give this product to children who are taking sedatives or tranquilizers, without first consulting the child's doctor.

Do not exceed recommended dosage. If nervousness, dizziness, or sleeplessness occur, discontinue use and consult a doctor. If symptoms do not improve within 7 days or are accompanied by fever, consult a doctor. Do not give this product to a child who has heart disease, high blood pressure, thyroid disease, or diabetes unless directed by a doctor. Do not use this product for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen. If symptoms persist, consult a doctor. Do not take this product if you have heart disease, high blood pressure, thyroid disease, diabetes, or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor. Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Directions

Adults and children 12 years of age and over: oral dosage is 4 milligrams every 4 to 6 hours, not to exceed 24 milligrams in 24 hours, or as directed by a doctor. Children 6 to under 12 years of age: oral dosage is 2 milligrams every 4 to 6 hours, not to exceed 12 milligrams in 24 hours, or as directed by a doctor. Children 2 to under 6 years of age: oral dosage is 1 milligram every 4 to 6 hours, not to exceed 6 milligrams in 24 hours.

CHLORPHENIRAMINE MALEATE, PHENYLEPHRINE HYDROCHLORIDE, ACETAMINOPHEN.

MADE IN MEXICO BY:

SCHERING-PLOUGH, S.A. DE C.V.
AV. 16 DE SEPTIEMBRE NO. 301
COL. XALTOCAN, MEXICO D.F.
16090 MEXICO. REG. NO. 55563 SSA VI



DESENFRIOL-D

chlorpheniramine maleate, phenylephrine hydrochloride, acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53666-617
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	0.002 g in 1 g
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	0.005 g in 1 g
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	0.5 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	

GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
XANTHAN GUM (UNII: TTV12P4NEE)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	18mm
Flavor		Imprint Code	PME
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53666-617-01	30 in 1 BOX	10/18/2017	
1		0.05 g in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/18/2017	

Labeler - Salimex, S.A. (589201581)

Registrant - Salimex, S.A. (589201581)

Establishment

Name	Address	ID/FEI	Business Operations
Bayer de México, S.A. de C.V.		588165709	manufacture(53666-617)

Revised: 12/2020

Salimex, S.A.