MAX TUSSIN COUGH AND CHEST CONGESTION DM SUGAR FREEdextromethorphan hbr, guaifenesin liquid Akron Pharma Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr 20 mg
Guaifenesin 200 mg

Purposes

Cough suppressant

Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

If pregnant or breast-feeding,

ask a health professional before use.

Stop use and ask a doctor if

cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL= milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

- Phenylketonurics: contains Phenylalanine 30 mg per 10 mL
- store between 20-25°C(68-77°F). Do not refrigerate

Inactive ingredients

anhydrous citric acid, flavor cherry, glycerin, menthol, purified water, saccharin sodium, sodium benzoate, sorbitol solution, sodium citrate, sucralose.

Questions or comments?

toll-free 1-877-225-6999

Manufactured for:

Akron Pharma, Inc.

Fairfield, NJ 07004

www.akronpharma.com



Expectorant (Guaifenesin)

SUGAR-FREE

estio ✓ Controls Coughs √ Loosens & Relieves **Chest Congestion**

Cherry FlavoredAlcohol Free

• Gluten Free • Dye Free Non-Drowsy

Ages 12+





4 FL OZ . (118 mL)

Drug Facts

Active ingredients (in each 10 mL) Jextromethorphan HBr, USP 20 mg

Suaifenesin, USP 200 mg

Sough suppressant

Active ingredients (in each 10 mL) Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg Expectorant Uses temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold ■ helps loosen phlegm

emotional conditions, or Parkinson's disease), or for 2 weeks after stopping

oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or

Warnings Do not use ■if you are now taking a prescription monoamine

ritation as may occur with a cold helps loosen phlegm (mucus) and

hin bronchial secretions to drain bronchial tubes

ISes = temporarily relieves cough due to minor throat and bronchial

before use if you have
cough that occurs with too much phlegm (mucus

MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor

the MAOI drug. If you do not know if your prescription drug contains an

pregnant or breast-feeding, ask a health professional before use. Keep out

Purposes

. Cough suppressant

of reach of children. In case of overdose, get medical help or contact

Poison Control Center right away (1-800-222-1222)

asts for more than 7 days, comes back, or is accompanied by fever, rash

or persistent headache. These could be signs of a serious condition. If

chronic bronchitis, or emphysema. Stop use and ask a doctor if cough

cough that lasts or is chronic such as occurs with smoking, asthma,

Directions ■ do not take more than 6 doses in any 24-hour period ■ measure only with dosing cup provided ■ keep dosing cup with product ■ mL = milliliter ■ this adult product is not intended for use in

(mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ cough that occurs with too much phlegm (mucus) cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be a signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Drug Facts (Continued)

children under 12 years of age

If pregnant or breast-feeding, ask a health professional before use.

Other information - Phenylketonurics: contains Phenylalanine 30 mg

per 10 mL ■ store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, flavor cherry, glycerin, menthol nronvlene alvcol. purified water, saccharin sodium, sodium

10 mL every 4 hours

adults and children 12 years and over

children under 12 years

do not use

Directions ■ do not take more than 6 doses in any 24-hour period ■ measure only with dosing cup provided ■ keep dosing cup with product ■mL = milliliter ■ this adult product is not intended for use in children under 12 years of age

age	dose	
adults and children 12 years and over	10 mL every 4 hours	
children under 12 years	do not use	

Other information Phenylketonurics: contains Phenylalanine 30 mg per 10 mL ■ store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, flavor cherry, glycerin, menthol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution, sodium citrate, sucralose.

Questions or comments? toll-free 1-877-225-6999

Tamper Evident: Do Not Use If Printed Safety Seal Around Bottle or Under Cap is Broken or Missing *This Product is not manufactured or distributed by Pfizer, distributor DRUGS/AP/122/97

Akran Pharma Manufactured for:

Akron Pharma, Inc. Fairfield, NJ 07004 www.akronpharma.com



MTD-1/23/L-01M

This Product is not manufactured or distributed by Pfizer, distributor of Robitussin® Sugar-Free Cough+Chest Congestion DM DRUGS/AP/122/97 | Tamper Fudent, Do Not Use If Printed Safery Seal Around Bottle or Under Can is Boxien or Missing I

Manufactured for: Akron Pharma, Inc. Fairfield, NJ 07004 www.akronpharma.com

de Pharma

Questions or comments? toll-free 1-877-225-6999

sodium citrate, sucralose.

benzoate, sorbitol solution.

NDC 71399-0026-08 MAXT COUGH & CHEST CONGESTION

Cough Suppressant (Dextromethorphan HBr) Expectorant (Guaifenesin)

SUGAR-FREE

✓ Controls Coughs ✓ Loosens & Relieves **Chest Congestion**

- Cherry Flavored Alcohol Free
- Gluten Free Dve Free

Ages 12+


Akr&n Pharma

8 FL OZ . (236 mL)



Cough Suppressant (Dextromethorphan HBr) Expectorant (Guaifenesin)

SUGAR-FREE



Ages 12+

Akr&n Pharma

16 FL OZ . (473 mL)

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg .. Cough suppressant Guaifenesin, USP 200 mg Expectorant

Purposes

Uses ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These cloud be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (Continued)

Directions ■ do not take more than 6 doses in any 24-hour period ■ mL = milliliter ■ this adult product is not intended for use in children under 12 years of age

, , , , , , , , , , , , , , , , , , , ,		
age	dose	
adults and children 12 years and over	10 mL every 4 hours	
children under 12 years	do not use	

Other information ■ Phenylketonurics: contains Phenylalanine 30 mg per 10 mL ■ store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid. flavor cherry, glycerin, menthol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution, sodium citrate, sucralose.

Questions or comments? toll-free 1-877-225-6999

*This Product is not manufactured or distributed by Pfizer, distributor of Robitussin® Sugar-Free Cough+Chest Congestion DM

Tamper Evident: Do Not Use If foil seal over bottle opening is torn broken or missing.

DRUGS/AP/122/97

Akr@n Pharma

Manufactured for: Akron Pharma, Inc. Fairfield, NJ 07004 www.akronpharma.com



MAX TUSSIN COUGH AND CHEST CONGESTION DM SUGAR FREE

dextromethorphan hbr, guaifenesin liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:71399-0026

Route of Administration ORAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength Strength **DEXTROMETHORPHAN HYDROBROMIDE** (UNII: 9D2RTI9KYH) **DEXTROMETHORPHAN** 20 mg **HYDROBROMIDE** (DEXTROMETHORPHAN - UNII:7355X3ROTS) in 10 mL 200 mg GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) **GUAIFENES IN** in 10 mL

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
GLYCERIN (UNII: PDC6A3C0OX)		
MENTHOL (UNII: L7T10EIP3A)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		

SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71399- 0026-4	1 in 1 PACKAGE	05/15/2023	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:71399- 0026-8	1 in 1 PACKAGE	05/15/2023	
2		236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:71399- 0026-6	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/15/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	05/15/2023	

Labeler - Akron Pharma Inc. (067878881)

Revised: 5/2023 Akron Pharma Inc.