

PREMIER VALUE POVIDONE IODINE- povidone iodine 10% liquid
Chain Drug Consortium

Premier Value Povidone Iodine

Drug Facts

Active Ingredient

Povidone-Iodine 10%

(Equivalent to 1% titrable iodine)

Purpose

Antiseptic

Use

First aid antiseptic to prevent infection in minor cuts and burns.

Warnings

For External Use Only

Ask a doctor before use if you have

- deep puncture wounds
- animal bites
- serious burns

Stop use and consult a doctor if

- the condition persists or gets worse
- irritation and redness develop and persists for more than 72 hours

When using this product do not

- use in eyes
- use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor
- apply over large areas of the body

Keep out of reach of children.

In case of accidental ingestion, seek professional assistance or contact a Poison Control

Center immediately.

Directions

Clean the affected area. Apply a small amount on the area 1 to 3 times daily. May be covered with sterile bandage. If badaged, let it dry first.

Other information

store at room temperature, 20-25C (68-77F)

Inactive Ingredients

Citric Acid, Disodium Phosphate, Glycerin, nonoxynol-9. Sodium Hydroxide and Purified Water.

Label

<table border="1"><tr><td colspan="2">Drug Facts</td></tr><tr><td>Active ingredient</td><td>Purpose</td></tr><tr><td>Povidone-iodine 10%.....</td><td>Antiseptic</td></tr><tr><td colspan="2">(*Equivalent to 1% tetratable iodine)</td></tr><tr><td colspan="2">Use</td></tr><tr><td colspan="2">First aid antiseptic to prevent infection in minor cuts and burns</td></tr><tr><td colspan="2">Warnings For external use only.</td></tr><tr><td colspan="2">Ask a doctor before use if you have</td></tr><tr><td colspan="2">■ deep or puncture wounds</td></tr><tr><td colspan="2">■ animal bites</td></tr><tr><td colspan="2">■ serious burns</td></tr><tr><td colspan="2">Stop use and consult a doctor if</td></tr><tr><td colspan="2">■ the condition persists or gets worse</td></tr><tr><td colspan="2">■ irritation and redness develop and persists for more than 72 hours</td></tr><tr><td colspan="2">When using this product do not:</td></tr><tr><td colspan="2">■ use in the eyes</td></tr><tr><td colspan="2">■ use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor</td></tr><tr><td colspan="2">■ apply over large areas of the body</td></tr><tr><td colspan="2">Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.</td></tr><tr><td colspan="2" style="text-align: right;"><i>Continued</i> ▶</td></tr></table>	Drug Facts		Active ingredient	Purpose	Povidone-iodine 10%.....	Antiseptic	(*Equivalent to 1% tetratable iodine)		Use		First aid antiseptic to prevent infection in minor cuts and burns		Warnings For external use only.		Ask a doctor before use if you have		■ deep or puncture wounds		■ animal bites		■ serious burns		Stop use and consult a doctor if		■ the condition persists or gets worse		■ irritation and redness develop and persists for more than 72 hours		When using this product do not:		■ use in the eyes		■ use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor		■ apply over large areas of the body		Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.		<i>Continued</i> ▶		<p style="text-align: right;">NDC 68086-415-01 **Compare to Active Ingredient in Betadine®</p>  <h1>Povidone-Iodine</h1> <p>10%* TOPICAL SOLUTION USP</p> <p>Topical Antiseptic Microbicide</p> <p>NET WT 8 FL OZ (227 mL)</p> 	<table border="1"><tr><td>Drug Facts (continued)</td></tr><tr><td>Directions</td></tr><tr><td>■ clean the affected area ■ apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first</td></tr><tr><td>Other information</td></tr><tr><td>■ store at room temperature, 20-25° C (68-77° F)</td></tr><tr><td>Inactive ingredients</td></tr><tr><td>Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water</td></tr><tr><td colspan="2">**This product is not manufactured or distributed by Parke-Products L.P., owner of the registered trademark Betadine®.</td></tr></table> <p style="text-align: right;">8127813RLE</p>  <p>8 40986 01294 5</p> <p><small>If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.</small></p> <p>DISTRIBUTED BY: CHAM-DRUG CONSORTIUM 3301 NW BOCARATON BLVD SUITE 101 BOCARATON, FL 33431 MADE IN U.S.A.</p>	Drug Facts (continued)	Directions	■ clean the affected area ■ apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first	Other information	■ store at room temperature, 20-25° C (68-77° F)	Inactive ingredients	Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water	**This product is not manufactured or distributed by Parke-Products L.P., owner of the registered trademark Betadine®.	
Drug Facts																																																			
Active ingredient	Purpose																																																		
Povidone-iodine 10%.....	Antiseptic																																																		
(*Equivalent to 1% tetratable iodine)																																																			
Use																																																			
First aid antiseptic to prevent infection in minor cuts and burns																																																			
Warnings For external use only.																																																			
Ask a doctor before use if you have																																																			
■ deep or puncture wounds																																																			
■ animal bites																																																			
■ serious burns																																																			
Stop use and consult a doctor if																																																			
■ the condition persists or gets worse																																																			
■ irritation and redness develop and persists for more than 72 hours																																																			
When using this product do not:																																																			
■ use in the eyes																																																			
■ use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor																																																			
■ apply over large areas of the body																																																			
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.																																																			
<i>Continued</i> ▶																																																			
Drug Facts (continued)																																																			
Directions																																																			
■ clean the affected area ■ apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first																																																			
Other information																																																			
■ store at room temperature, 20-25° C (68-77° F)																																																			
Inactive ingredients																																																			
Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water																																																			
**This product is not manufactured or distributed by Parke-Products L.P., owner of the registered trademark Betadine®.																																																			

Drug Facts

Active ingredient **Purpose**
 Povidone-iodine 10%.....Antiseptic
 (*Equivalent to 1% titratable iodine)

Use
 First aid antiseptic to prevent infection in minor cuts and burns

Warnings For external use only.
Ask a doctor before use if you have
 ■ deep or puncture wounds
 ■ animal bites
 ■ serious burns

Stop use and consult a doctor if
 ■ the condition persists or gets worse
 ■ irritation and redness develop and persists for more than 72 hours

When using this product do not
 ■ do not use in the eyes
 ■ use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor
 ■ do not apply over large areas of the body

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.
Continued ▶



NDC 68016-415-00
 **Compare to Active Ingredient in Betadine®

Povidone-Iodine

10%* TOPICAL SOLUTION USP

Topical Antiseptic Microbicide

NET WT. 8 FL OZ (237 mL)



Drug Facts (continued)

Directions
 ■ clean the affected area ■ apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first

Other information
 ■ store at room temperature, 20-25° C (68-77° F)

Inactive ingredients
 Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water

**This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine®.

R110113.RLG



8 40986 01294 5

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

DISTRIBUTED BY: CHAIN DRUG CONSORTIUM
 3301 NW BOCA RATON BLVD
 SUITE 101, BOCA RATON, FL 33431
 MADE IN U.S.A.

PREMIER VALUE POVIDONE IODINE

povidone iodine 10% liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-415
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POVIDONE-IODINE (UNII: 85H0HZ U99M) (IODINE - UNII:9679TC07X4)	IODINE	100 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
NONOXYNOL-9 (UNII: 48Q180SH9T)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	NDC:68016-415-00	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/30/2017	
----------	------------------	----------------------------------------------------------------	------------	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	01/01/2008	

Labeler - Chain Drug Consortium (101668460)

Registrant - Pharma Nobis, LLC (118564114)

Establishment

Name	Address	ID/FEI	Business Operations
Pharma Nobis, LLC		118564114	manufacture(68016-415) , analysis(68016-415) , pack(68016-415) , label(68016-415)

Revised: 12/2023

Chain Drug Consortium