

LANSOPRAZOLE- lansoprazole capsule, delayed release
Safeway, Inc.

Drug Facts

Active ingredient (in each capsule)

Lansoprazole 15 mg

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

WARNINGS

Clostridium difficile-associated diarrhea (CDAD): A diagnosis of CDAD should be considered for patients taking Proton Pump Inhibitors (PPIs) who develop diarrhea that does not improve. Symptoms include:

- watery stool
- abdominal pain
- fever

Patients should seek immediate care from a healthcare professional if these symptoms do not go away while taking this medication.

Allergy alert: Do not use if you are allergic to lansoprazole.

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

taking

- warfarin (blood-thinning medicine)

- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus (immune system medicine)
- atazanavir (medicine for HIV infection)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor. Patients should use the lowest dose and shortest duration of this therapy.**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20°-25°C (68°-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients

colloidal silicon dioxide, D&C Red No. 33, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, gelatin, hypromellose, magnesium carbonate, methacrylic acid copolymer dispersion, polyethylene glycol, polysorbate 80, sucrose, sugar spheres, talc, titanium dioxide

Questions or comments?

Call **1-888-SAFEWAY**

Monday-Friday 7AM-6PM PST

Principal Display Panel

Compare to Active Ingredient in Prevacid®24HR*

24 Hour

Lansoprazole

Delayed-Release Capsules, USP 15 mg

Acid Reducer

- May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- Clinically proven to treat frequent heartburn

Sodium Free**SEE NEW WARNINGS INFORMATION**

CAPSULES 14-DAY COURSE OF TREATMENT

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE

Do not use if inner foil seal imprinted with "Sealed For Your Protection" or dark blue to black gelatin band around the center of each capsule is missing or broken.

DISTRIBUTED BY SAFEWAY INC.

P.O. BOX 99, PLEASANTON, CA 94566-0009

1-888-SAFEWAY / www.safeway.com

Product of India

*This product is not manufactured or distributed by Takeda Pharmaceuticals North America, Inc., owner of the registered trademark Prevacid®, or by Novartis Consumer Health, Inc., distributor of the Prevacid®24HR product.

Product label

Drug Facts	Drug Facts (continued)
Active ingredient (in each capsule) Lansoprazole 15 mg	Purpose Acid reducer
Use ● treats frequent heartburn (occurs 2 or more days a week) ● not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect	Warnings Allergy alert: Do not use if you are allergic to lansoprazole. Do not use ● if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have ● liver disease ● had heartburn over 3 months. This may be a sign of a more serious condition. ● heartburn with lightheadedness, sweating, or dizziness ● chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck, or shoulders, or lightheadedness ● frequent chest pain ● frequent wheezing, particularly with heartburn ● unexplained weight loss ● nausea or vomiting ● stomach pain Ask a doctor or pharmacist before use if you are taking ● warfarin (blood thinning medicine) ● prescription antifungal or anti-yeast medicines ● digoxin (heart medicine) ● theophylline (asthma medicine) ● tacrolimus (immune system medicine) ● zalcitabine (medicine for HIV infection) Step use and ask a doctor if ● your heartburn continues or worsens
Warnings Allergy alert: Do not use if you are allergic to lansoprazole. Do not use ● if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have ● liver disease ● had heartburn over 3 months. This may be a sign of a more serious condition. ● heartburn with lightheadedness, sweating, or dizziness ● chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck, or shoulders, or lightheadedness ● frequent chest pain ● frequent wheezing, particularly with heartburn ● unexplained weight loss ● nausea or vomiting ● stomach pain Ask a doctor or pharmacist before use if you are taking ● warfarin (blood thinning medicine) ● prescription antifungal or anti-yeast medicines ● digoxin (heart medicine) ● theophylline (asthma medicine) ● tacrolimus (immune system medicine) ● zalcitabine (medicine for HIV infection) Step use and ask a doctor if ● your heartburn continues or worsens	Directions ● adults 18 years of age and older ● this product is to be used once a day (every 24 hours), every day for 14 days ● it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours 14-Day Course of Treatment ● swallow 1 capsule with a glass of water before eating in the morning ● take every day for 14 days ● do not take more than 1 capsule a day ● swallow whole. Do not crush or chew capsules. ● do not use for more than 14 days unless directed by your doctor Repeated 14-Day Courses (if needed) ● you may repeat a 14-day course every 4 months ● do not take for more than 14 days or more often than every 4 months unless directed by a doctor Patients should use the lowest dose and shortest duration of this therapy. ● children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. Other information ● read the directions, warnings, and package insert before use ● keep the carton and package insert. They contain important information. ● store at 20°-25°C (68°-77°F) ● keep product out of high heat and humidity ● protect product from moisture


Drug Facts (continued)
Inactive ingredients: coloidal silicon dioxide, D&C Red No. 33, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, gelatin, hypromellose, magnesium carbonate, methacrylic acid copolymer dispersion, polyethylene glycol, polyoxane 68, sucrose, sugar spheres, talc, titanium dioxide
Questions or comments? (Call 1-888-725-9929) Monday-Friday 7AM-6PM PST

TAMPER-EVIDENT BOTTLE: DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR DARK BLUE TO BLACK GELATIN BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

24 Hour

Lansoprazole

Delayed-Release Capsules, USP 15 mg Acid Reducer



ACTUAL SIZE

Sodium Free

14 CAPSULES ONE 14-DAY COURSE OF TREATMENT

NDC 21130-019-14

Compare to Active Ingredient in Prevacid®24HR*

- May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- Clinically proven to treat frequent heartburn

SAFEMAY

KEEP OUTER CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

*This product is not manufactured or distributed by Takeda Pharmaceuticals North America, Inc., owner of the registered trademark Prevacid®, or by Rowland Consumer Health, Inc., distributor of the Prevacid®24HR product.


DISTRIBUTED BY SAFEMAY INC.
P.O. BOX 99, PLEASANTON, CA 94568-0099
QUALITY & SATISFACTION GUARANTEED OR YOUR MONEY BACK
1-888-SAFEMAY / www.safemay.com

PRODUCT OF INDIA 1 Bottle Inside

PLD-D
FCD00558

Lot No.:
Exp. Date:

RD 13296 52801



3 21130 77900 1

Safeway Lansoprazole Capsule

LANSOPRAZOLE			
lansoprazole capsule, delayed release			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-019(NDC:64679-140)
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
LANSOPRAZOLE (UNII: 0K5C5T2QPG) (LANSOPRAZOLE - UNII:0K5C5T2QPG)	LANSOPRAZOLE	15 mg	
Inactive Ingredients			
Ingredient Name	Strength		
D&C RED NO. 33 (UNII: 9DBA0SBB0L)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GELATIN (UNII: 2G86QN327L)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			
MAGNESIUM CARBONATE (UNII: 0E53J927NA)			
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)			
POLYSORBATE 80 (UNII: 6OZP39ZG8H)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
SUCROSE (UNII: C151H8M554)			
TALC (UNII: 7SEV7J4R1U)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			

METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)

STARCH, CORN (UNII: O8232NY3SJ)

Product Characteristics

Color	PINK, GREEN	Score	no score
Shape	CAPSULE	Size	16mm
Flavor		Imprint Code	W140
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-019-14	1 in 1 BOX		
1		14 in 1 BOTTLE		
2	NDC:21130-019-42	3 in 1 BOX		
2		14 in 1 BOTTLE		
3	NDC:21130-019-28	2 in 1 BOX		
3		14 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202727	12/31/2012	

Labeler - Safeway, Inc. (009137209)

Registrant - P and L Development of New York Corporation (800014821)

Revised: 1/2013

Safeway, Inc.