POLYETHYLENE GLYCOL 3350- polyethylene glycol 3350 powder, for solution Marlex Pharmaceuticals Inc

Polyethylene Glycol 3350

MARLEX PHARMACEUTICALS, INC

Polyethylene Glycol 3350 powder, for solution

Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity) generally
- produces a bowel movement in 1 to 3 days

Warnings

Allergy alert:

Do not use if you are allergic to polyethylene glycol

Do not use

if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These
 may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (clear section in cap)
- adults and children 17 years of age and older:
 - o fill to top of clear section in cap which is marked to indicate the correct dose (17 g)
 - o stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - o use once a day
 - o use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° 25°C (68° 77°F)
- tamper- evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredient

none

Questions or comments?

1-800-FDA-1088

Manufactured for & Distributed by:


Marlex Pharmaceuticals, Inc.

New Castle, DE 19720

Rev. 04/19IT


PRINCIPAL DISPLAY PANEL

NDC 10135-0683-04 Polyethylene Glycol 3350 Powder for Solution, Osmotic Laxative Net Wt 4.1 oz (119 g) 7 Once-Daily Doses




PRINCIPAL DISPLAY PANEL

NDC 10135-0683-08 Polyethylene Glycol 3350 Powder for Solution, Osmotic Laxative Net Wt 8.3 oz (238 g) 14 Once-Daily Doses



PRINCIPAL DISPLAY PANEL

NDC 10135-0683-12 Polyethylene Glycol 3350 Powder for Solution, Osmotic Laxative Net Wt 17.9 oz (510 g) 30 Once-Daily Doses



POLYETHYLENE GLYCOL 3350

polyethylene glycol 3350 powder, for solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:10135-683
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g	

Product Characteristics			
Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

F	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10135-683-04	119 g in 1 BOTTLE; Type 0: Not a Combination Product	0 4/0 1/20 19	
2	NDC:10135-683-08	238 g in 1 BOTTLE; Type 0: Not a Combination Product	0 4/0 1/20 19	
3	NDC:10135-683-12	510 g in 1 BOTTLE; Type 0: Not a Combination Product	0 4/0 1/20 19	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA202071	04/01/2019		

Labeler - Marlex Pharmaceuticals Inc (782540215)

Revised: 5/2019 Marlex Pharmaceuticals Inc