

**AMPROMED-P FOR POULTRY- amprolium powder**  
**Bimeda, Inc., Division of Cross Vetpharm Group**

-----  
**AmproMed™-P FOR POULTRY**

**(amprolium)**

**20% Soluble Powder**  
**Coccidiostat**

**NOT FOR HUMAN USE**

**KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN**

**Water Soluble Treatment for Coccidiosis**

**Contents of each packet will treat 128 gallons at 0.012% level.**

**Restricted Drug (California) – Use Only as Directed**

ANADA 200-488, Approved by FDA

**INDICATIONS:** AmproMed™ P FOR POULTRY is intended for the treatment of coccidiosis in growing chickens, turkeys and laying hens.

**Net Weight: 10 oz (283.5 g)**

**ACTIVE INGREDIENT: amprolium.....20%**

**Use Directions:** 1 oz = 3 1/2 measuring tablespoons

**PREMEASURED FOR AUTOMATIC WATER PROPORTIONERS:** For use in automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.

<b>To prepare dosage levels of</b>	Dissolve AmproMed™ P FOR POULTRY in one gallon of water. (Makes 1 gallon of stock solution.)
0.024%	2 bags (20 oz) in 1 gallon
0.012%	1 bag (10 oz) in 1 gallon
0.006%	1/2 bag (5 oz) in 1 gallon

**To prepare 50 Gallons of Medicated Water:**

<b>DOSAGE LEVEL</b>	<b>MIXING DIRECTIONS</b>
0.024%	Dissolve 8 ounces of AmproMed™ P FOR POULTRY in about 5 gallons of water in a 50 gallon medication barrel. Stir, then add water to the 50 gallon mark. Stir thoroughly.
0.012%	Follow same directions as above but use 4 ounces of AmproMed™ P FOR POULTRY
0.006%	Follow same directions as above but use 2 ounces of AmproMed™ P FOR POULTRY

**Note:** Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

**USE DIRECTIONS:** Give amprolium at the 0.012% level (10 oz AmproMed™ P FOR POULTRY per 128 Gallons) as soon as coccidiosis is diagnosed and continue for three to five days. (In severe outbreaks, give amprolium at the 0.024% level). Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time.

Use as sole source of amprolium.

**CAUTION:** If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

**STORAGE:** Store at 20°C - 25° C (68° F - 77 ° F), excursions permitted between 15° C - 30° C (59° F - 86°F).

To obtain an MSDS or for assistance contact Bimeda, Inc. at 1-888-524-6332.

Take Time Observe Label Directions

**Bimeda**

**AmproMed™ P FOR POULTRY**  
(amprolium)

**20% Soluble Powder**  
**Coccidiostat**

**NOT FOR HUMAN USE**  
**KEEP THIS AND ALL DRUGS OUT**  
**OF REACH OF CHILDREN**

**Water-Soluble Treatment for Coccidiosis**  
Contents of each packet will treat 128 gallons at 0.012% level

**Restricted Drug (California) – Use Only as Directed**  
ANADA 200-488, Approved by FDA

**INDICATIONS:** AmproMed™ P FOR POULTRY is intended for the treatment of coccidiosis in growing chickens, turkeys and laying hens.

**Net Weight: 10 oz (283.5 g)**

ACTIVE INGREDIENT: amprolium.....20%

Manufactured by:  
Bimeda, Inc.  
Le Sueur, MN 56058  
www.bimeda.com

**AmproMed™ P FOR POULTRY** 20% Soluble Powder  
(amprolium) Coccidiostat

**Use Directions:** 1 oz = 3 1/2 measuring teaspoonfuls

**PREMEASURED FOR AUTOMATIC WATER PROPORTIONERS:** For use in automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.

<b>To prepare dosage levels of:</b>	Dissolve AmproMed™ P FOR POULTRY in one gallon of water. (Makes 1 gallon of stock solution.)
0.024%	2 bags (20 oz) in 1 gallon
0.012%	1 bag (10 oz) in 1 gallon
0.006%	1/2 bag (5 oz) in 1 gallon

**To prepare 50 Gallons of Medicated Water:**

DOSAGE LEVEL	MIXING DIRECTIONS
0.024%	Dissolve 8 ounces of AmproMed™ P FOR POULTRY in about 5 gallons of water in a 50 gallon medication barrel. Stir, then add water to the 50 gallon mark. <i>Stir thoroughly.</i>
0.012%	Follow same directions as above but use 4 ounces of AmproMed™ P FOR POULTRY.
0.006%	Follow same directions as above but use 2 ounces of AmproMed™ P FOR POULTRY.

**Note:** Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

**USE DIRECTIONS:** Give amprolium at the 0.012% level (10 oz AmproMed™ P FOR POULTRY per 128 Gallons) as soon as coccidiosis is diagnosed and continue for three to five days. (In severe outbreaks, give amprolium at the 0.024% level). Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time.

Use as sole source of amprolium.

**CAUTION:** If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

**STORAGE:** Store at 20°C-25°C (68°F-77°F), excursions permitted between 15°C-30°C (59°F-86°F).  
To obtain an MSDS or for assistance contact Bimeda, Inc. at 1-888-524-6332.

TAKE TIME OBSERVE LABEL DIRECTIONS

**Bimeda**

Product No. 1AMP022  
8AMP014 Rev. 11/11

## AMPROMED-P FOR POULTRY

amprolium powder

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:61133-8252
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
AMPROLIUM (UNII: 95CO6N199Q) (AMPROLIUM ION - UNII:H2T307KMZR)	AMPROLIUM ION	56.7 g in 283.5 g

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61133-8252-1	283.5 g in 1 POUCH		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200488	05/23/2011	

**Labeler** - Bimeda, Inc., Division of Cross Vetpharm Group (060492923)

**Registrant** - Bimeda, Inc., Division of Cross Vetpharm Group (060492923)

**Establishment**

Name	Address	ID/FEI	Business Operations
Bimeda, Inc., Division of Cross Vetpharm Group		060492923	manufacture

**Establishment**

Name	Address	ID/FEI	Business Operations
Zhejiang Juizhou		654682061	api manufacture

Revised: 8/2013

Bimeda, Inc., Division of Cross Vetpharm Group