

**ANTACID ANTIGAS NTA- aluminum hydroxide, magnesium hydroxide,
simethicone liquid
AMERISOURCEBERGEN DRUG CORPORATION**

GNP Antacid Anti-Gas max

Active ingredients (in each 10 mL dose)

Aluminum hydroxide (equivalent to dried gel, USP) 800 mg

Magnesium hydroxide 800 mg

Simethicone 80 mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- symptoms of gas

Warnings

Do not take more than 60 mL (6 doses) in a 24-hour period or use the maximum dosage for more than 2 weeks.

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug.

Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use

- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses) between meals as needed, at bedtime, or as directed by a doctor
- do not take more than 60 mL (6 doses) in any 24 hour period or use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor

Other information

- **each 10 mL dose contains:** magnesium 340 mg, sodium 10 mg
- do not freeze
- store at room temperature tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or Comments?

1-800-540-3765

package label



Compare to Mylanta®
Maximum Strength
active ingredients*

NDC 46122-431-40

MAXIMUM STRENGTH

Antacid and Anti-Gas Liquid

**Alumina, Magnesia, and
Simethicone Oral Suspension**

Fast Relief of:

- Acid Indigestion
- Heartburn
- Sour Stomach
- Pressure and Bloating

ORIGINAL FLAVOR

Contains: Alcohol 0.5%

12 fl oz (355 mL)

297-06122-0 REV0917

Drug Facts

TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY
BAND ON CAP IS BROKEN OR MISSING.

Active ingredients (in each 10 mL dose)

Aluminum hydroxide (equivalent to dried gel, USP) 800 mg	Antacid
Magnesium hydroxide 800 mg	Antacid
Simethicone 80 mg	Antigas

Purposes

Uses relieves • heartburn • sour stomach • acid indigestion
• the symptoms referred to as gas

Warnings

Do not take more than 60 mL (6 doses) in a 24-hour period or use
the maximum dosage for more than 2 weeks.

Ask a doctor before use if you have

- kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a
prescription drug. Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 10 mL to 20 mL
(1 to 2 doses) between meals as needed, at bedtime, or as directed
by a doctor
- do not take more than 60 mL (6 doses) in any 24-hour period or
use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor

Other information

- each 10 mL dose contains: magnesium 340 mg, sodium 10 mg
- do not freeze • store at room temperature tightly closed

Inactive ingredients benzyl alcohol, butylparaben,
carboxymethylcellulose sodium, flavor (contains alcohol),
hypromellose, microcrystalline cellulose, propylparaben,
purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765



Distributed By
Amersource Bergen
1 West First Avenue
Conshohocken, PA 19428
Questions or Concerns?
www.mygnp.com

10181208 997-06122-1 REV0521



ANTACID ANTIGAS NTA

aluminum hydroxide, magnesium hydroxide, simethicone liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-431
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	800 mg in 10 mL

MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)		MAGNESIUM HYDROXIDE	800 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)		DIMETHICONE	80 mg in 10 mL

Inactive Ingredients	
Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-431-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/31/2018	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	09/01/2017	

Labeler - AMERISOURCEBERGEN DRUG CORPORATION (007914906)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(46122-431)

Revised: 11/2023

AMERISOURCEBERGEN DRUG CORPORATION